

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/85, 9/10, C07K 16/30, 16/40,</b> <b>C12Q 1/68, A01K 67/027</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 99/32644</b> <b>(43) International Publication Date:</b> 1 July 1999 (01.07.99)
<b>(21) International Application Number:</b> PCT/IB98/02133 <b>(22) International Filing Date:</b> 22 December 1998 (22.12.98)  <b>(30) Priority Data:</b> 08/996,306 22 December 1997 (22.12.97) US 60/099,658 9 September 1998 (09.09.98) US  <b>(71) Applicant (for all designated States except US):</b> GENSET [FR/FR]; 24, rue Royale, F-75008 Paris (FR).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> COHEN, Daniel [FR/FR]; 5, avenue Odette, F-94210 Fontenay-sous-Bois (FR). BLUMENFELD, Marta [FR/FR]; 5, rue Tagore, F-75013 Paris (FR). CHUMAKOV, Ilya [FR/FR]; 196, rue des Chèvrefeuilles, F-77000 Vaux-le-Pénil (FR). BOUGUEL- ERET, Lydie [FR/FR]; 108, avenue Victor-Hugo, F-92170 Vanves (FR).  <b>(74) Agents:</b> MARTIN, Jean-Jacques et al.; Cabinet Regimbeau, 26, avenue Kléber, F-75116 Paris (FR).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> PROSTATE CANCER GENE  <b>(57) Abstract</b>  The present invention relates to PG1, a gene associated with prostate cancer. The invention provides polynucleotides including biallelic markers derived from PG1 and from flanking genomic regions. Primers hybridizing to these biallelic markers and regions flanking are also provided. This invention provides polynucleotides and methods suitable for genotyping a nucleic acid containing sample for one or more biallelic markers of the invention. Further, the invention provides methods to detect a statistical correlation between a biallelic marker allele and prostate cancer and between a haplotype and prostate cancer. The invention also relates to diagnostic methods of determining whether an individual is at risk for developing prostate cancer, and whether an individual suffers from prostate cancer as a result of a mutation in the PG1 gene.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## PROSTATE CANCER GENE

### Background of the Invention

5 A cancer is a clonal proliferation of cells produced as a consequence of cumulative genetic damage that finally results in unrestrained cell growth, tissue invasion and metastasis (cell transformation). Regardless of the type of cancer, transformed cells carry damaged DNA in many forms: as gross chromosomal translocations or, more subtly, as DNA amplification, rearrangement or even point mutations.

10 Some oncogenic mutations is inherited in the germline, thus predisposing the mutation carrier to an increased risk of cancer. However, in a majority of cases, cancer does not occur as a simple monogenic disease with clear Mendelian inheritance. There is only a two- or threefold increased risk of cancer among first-degree relatives for many cancers (Mulvihill JJ, Miller RW & Fraumeni JF, 1977, Genetics of human cancer Vol 3, New York Raven Press). Alternatively, DNA damage is  
15 acquired somatically, probably induced by exposure to environmental carcinogens. Somatic mutations are generally responsible for the vast majority of cancer cases.

Studies of the age dependence of cancer have suggested that several successive mutations are needed to convert a normal cell into an invasive carcinoma. Since human mutation rates are typically  $10^{-6}$ /gene/cell, the chance of a single cell undergoing many independent mutations is very low (Loeb  
20 LA, Cancer Res 1991, 51: 3075-3079). Cancer nevertheless happens because of a combination of two mechanisms. Some mutations enhance cell proliferation, increasing the target population of cells for the next mutation. Other mutations affect the stability of the entire genome, increasing the overall mutation rate, as in the case of mismatch repair proteins (reviewed in Arnheim N & Shibata D, Curr. Op. Genetics & Development, 1997, 7:364-370).

25 An intricate process known as the cell cycle drives normal proliferation of cells in an organism. Regulation of the extent of cell cycle activity and the orderly execution of sequential steps within the cycle ensure the normal development and homeostasis of the organism. Conversely, many of the properties of cancer cells - uncontrolled proliferation, increased mutation rate, abnormal translocations and gene amplifications - can be attributed directly to perturbations of the normal  
30 regulation or progression of the cycle. In fact, many of the genes that have been identified over the past several decades as being involved in cancer, can now be appreciated in terms of their direct or indirect role in either regulating entry into the cell cycle or coordinating events within the cell cycle.

Recent studies have identified three groups of genes which are frequently mutated in cancer. The first group of genes, called oncogenes, are genes whose products activate cell proliferation. The  
35 normal non-mutant versions are called protooncogenes. The mutated forms are excessively or inappropriately active in promoting cell proliferation, and act in the cell in a dominant way in that a

single mutant allele is enough to affect the cell phenotype. Activated oncogenes are rarely transmitted as germline mutations since they may probably be lethal when expressed in all the cells. Therefore oncogenes can only be investigated in tumor tissues.

Oncogenes and protooncogenes can be classified into several different categories according to their function. This classification includes genes that code for proteins involved in signal transduction such as: growth factors (i.e., sis, int-2); receptor and non-receptor protein-tyrosine kinases (i.e., erbB, src, bcr-abl, met, trk); membrane-associated G proteins (i.e., ras); cytoplasmic protein kinases (i.e., mitogen-activated protein kinase -MAPK- family, raf, mos, pak), or nuclear transcription factors (i.e., myc, myb, fos, jun, rel) (for review see Hunter T, 1991 Cell 64:249; Fanger GR et al., 1997 Curr.Op.Genet.Dev.7:67-74; Weiss FU et al., ibid. 80-86).

The second group of genes which are frequently mutated in cancer, called tumor suppressor genes, are genes whose products inhibit cell growth. Mutant versions in cancer cells have lost their normal function, and act in the cell in a recessive way in that both copies of the gene must be inactivated in order to change the cell phenotype. Most importantly, the tumor phenotype can be rescued by the wild type allele, as shown by cell fusion experiments first described by Harris and colleagues (Harris H et al., 1969, Nature 223:363-368). Germline mutations of tumor suppressor genes is transmitted and thus studied in both constitutional and tumor DNA from familial or sporadic cases. The current family of tumor suppressors includes DNA-binding transcription factors (i.e., p53, WT1), transcription regulators (i.e., RB, APC, probably BRCA1), protein kinase inhibitors (i.e., p16), among others (for review, see Haber D & Harlow E, 1997, Nature Genet. 16:320-322).

The third group of genes which are frequently mutated in cancer, called mutator genes, are responsible for maintaining genome integrity and/or low mutation rates. Loss of function of both alleles increase cell mutation rates, and as consequence, proto-oncogenes and tumor suppressor genes is mutated. Mutator genes can also be classified as tumor suppressor genes, except for the fact that tumorigenesis caused by this class of genes cannot be suppressed simply by restoration of a wild-type allele, as described above. Genes whose inactivation may lead to a mutator phenotype include mismatch repair genes (i.e., MLH1, MSH2), DNA helicases (i.e., BLM, WRN) or other genes involved in DNA repair and genomic stability (i.e., p53, possibly BRCA1 and BRCA2) (For review see Haber D & Harlow E, 1997, Nature Genet. 16:320-322; Fishel R & Wilson T. 1997, Curr.Op.Genet.Dev.7: 105-113; Ellis NA, 1997 ibid.354-363).

The recent development of sophisticated techniques for genetic mapping has resulted in an ever expanding list of genes associated with particular types of human cancers. The human haploid genome contains an estimated 80,000 to 100,000 genes scattered on a  $3 \times 10^9$  base-long double-stranded DNA. Each human being is diploid, i.e., possesses two haploid genomes, one from paternal origin, the other from maternal origin. The sequence of a given genetic locus may vary between individuals in a population or between the two copies of the locus on the chromosomes of a single



individual. Genetic mapping techniques often exploit these differences, which are called polymorphisms, to map the location of genes associated with human phenotypes.

One mapping technique, called the loss of heterozygosity (LOH) technique, is often employed to detect genes in which a loss of function results in a cancer, such as the tumor suppressor genes described above. Tumor suppressor genes often produce cancer via a two hit mechanism in which a first mutation, such as a point mutation (or a small deletion or insertion) inactivates one allele of the tumor suppressor gene. Often, this first mutation is inherited from generation to generation.

A second mutation, often a spontaneous somatic mutation such as a deletion which deletes all or part of the chromosome carrying the other copy of the tumor suppressor gene, results in a cell in which both copies of the tumor suppressor gene are inactive.

As a consequence of the deletion in the tumor suppressor gene, one allele is lost for any genetic marker located close to the tumor suppressor gene. Thus, if the patient is heterozygous for a marker, the tumor tissue loses heterozygosity, becoming homozygous or hemizygous. This loss of heterozygosity generally provides strong evidence for the existence of a tumor suppressor gene in the lost region.

By genotyping pairs of blood and tumor samples from affected individuals with a set of highly polymorphic genetic markers, such as microsatellites, covering the whole genome, one can discover candidate locations for tumor suppressor genes. Due to the presence of contaminant non-tumor tissue in most pathological tumor samples, a decreased relative intensity rather than total loss of heterozygosity of informative microsatellites is observed in the tumor samples. Therefore, classic LOH analysis generally requires quantitative PCR analysis, often limiting the power of detection of this technique. Another limitation of LOH studies resides on the fact that they only allow the definition of rather large candidate regions, typically spanning over several megabases. Refinement of such candidate regions requires the definition of the minimally overlapping portion of LOH regions identified in tumor tissues from several hundreds of affected patients.

Another approach to genetic mapping, called linkage analysis, is based upon establishing a correlation between the transmission of genetic markers and that of a specific trait throughout generations within a family. In this approach, all members of a series of affected families are genotyped with a few hundred markers, typically microsatellite markers, which are distributed at an average density of one every 10 Mb. By comparing genotypes in all family members, one can attribute sets of alleles to parental haploid genomes (haplotyping or phase determination). The origin of recombined fragments is then determined in the offspring of all families. Those that co-segregate with the trait are tracked. After pooling data from all families, statistical methods are used to determine the likelihood that the marker and the trait are segregating independently in all families. As a result of the statistical analysis, one or several regions are selected as candidates, based on their high probability to carry a trait causing allele. The result of linkage analysis is considered as significant

when the chance of independent segregation is lower than 1 in 1000 (expressed as a LOD score  $> 3$ ). Identification of recombinant individuals using additional markers allows further delineation of the candidate linked region, which most usually ranges from 2 to 20 Mb.

Linkage analysis studies have generally relied on the use of microsatellite markers (also  
5 called simple tandem repeat polymorphisms, or simple sequence length polymorphisms). These include small arrays of tandem repeats of simple sequences (di- tri- tetra- nucleotide repeats), which exhibit a high degree of length polymorphism, and thus a high level of informativeness. To date, only just more than 5,000 microsatellites have been ordered along the human genome (Dib et al., Nature 1996, 380: 152), thus limiting the maximum attainable resolution of linkage analysis to ca. 600 kb on  
10 average.

Linkage analysis has been successfully applied to map simple genetic traits that show clear Mendelian inheritance patterns. About 100 pathological trait-causing genes were discovered by linkage analysis over the last 10 years.

However, linkage analysis approaches have proven difficult for complex genetic traits, those  
15 probably due to the combined action of multiple genes and/or environmental factors. In such cases, too large an effort and cost are needed to recruit the adequate number of affected families required for applying linkage analysis to these situations, as recently discussed by Risch, N. and Merikangas, K. (Science 1996, 273: 1516-1517). Finally, linkage analysis cannot be applied to the study of traits for which no available large informative families are available. Typically, this will be the case in any  
20 attempt to identify trait-causing alleles involved in sporadic cases.

The incidence of prostate cancer has dramatically increased over the last decades. It averages 30-50/100,000 males both in Western European countries as well as within the US White male population. In these countries, it has recently become the most commonly diagnosed malignancy, being one of every four cancers diagnosed in American males. Prostate cancer's incidence is very  
25 much population specific, since it varies from 2/100,000 in China, to over 80/100,000 among African-American males.

In France, the incidence of prostate cancer is 35/100,000 males and it is increasing by 10/100,000 per decade. Mortality due to prostate cancer is also growing accordingly. It is the second cause of cancer death among French males, and the first one among French males aged over 70. This  
30 makes prostate cancer a serious burden in terms of public health, especially in view of the aging of populations.

An average 40% reduction in life expectancy affects males with prostate cancer. If completely localized, prostate cancer can be cured by surgery, with however an average success rate of only ca. 50%. If diagnosed after metastasis from the prostate, prostate cancer is a fatal disease for  
35 which there is no curative treatment.

Early-stage diagnosis relies on Prostate Specific Antigen (PSA) dosage, and would allow the

detection of prostate cancer seven years before clinical symptoms become apparent. The effectiveness of PSA dosage diagnosis is however limited, due to its inability to discriminate between malignant and non-malignant affections of the organ.

Therefore, there is a strong need for both a reliable diagnostic procedure which would enable  
5 early-stage prostate cancer prognosis, and for preventive and curative treatments of the disease. The present invention relates to the PG1 gene, a gene associated with prostate cancer, as well as diagnostic methods and reagents for detecting alleles of the gene which may cause prostate cancer, and therapies for treating prostate cancer.

#### 10 Summary of the Invention

The present invention relates to the identification of a gene associated with prostate cancer, identified as the PG1 gene, and reagents, diagnostics, and therapies related thereto. The present invention is also based on the discovery of a novel set of PG1-related biallelic markers. See the definition of PG1-related biallelic markers in the Detailed Description Section. These markers are  
15 located in the coding regions as well as non-coding regions adjacent to the PG1 gene. The position of these markers and knowledge of the surrounding sequence has been used to design polynucleotide compositions which are useful in determining the identity of nucleotides at the marker position, as well as more complex association and haplotyping studies which are useful in determining the genetic basis for diseases including cancer and prostate cancer. In addition, the compositions and methods of  
20 the invention find use in the identification of the targets for the development of pharmaceutical agents and diagnostic methods, as well as the characterization of the differential efficacious responses to and side effects from pharmaceutical agents acting on diseases including cancer and prostate cancer.

A first embodiment of the invention is a recombinant, purified or isolated polynucleotide comprising, or consisting of a mammalian genomic sequence, gene, or fragments thereof. In one  
25 aspect the sequence is derived from a human, mouse or other mammal. In a preferred aspect, the genomic sequence is the human genomic sequence of SEQ ID NO: 179 or the complement thereto. In a second preferred aspect, the genomic sequence is selected from one of the two mouse genomic fragments of SEQ ID NO: 182 and 183. In yet another aspect of this embodiment, the nucleic acid comprises nucleotides 1629 through 1870 of the sequence of SEQ ID NO: 179. Optionally, said  
30 polynucleotide consists of, consists essentially of, or comprises a contiguous span of nucleotides of a mammalian genomic sequence, preferably a sequence selected the following SEQ ID NOs: 179, 182, and 183, wherein said contiguous span is at least 6, 8, 10, 12, 15, 20, 25, 30, 50, 100, 200, or 500 nucleotides in length.

A second embodiment of the present invention is a recombinant, purified or isolated  
35 polynucleotide comprising, or consisting of a mammalian cDNA sequence, or fragments thereof. In one aspect the sequence is derived from a human, mouse or other mammal. In a preferred aspect, the

cDNA sequence is selected from the human cDNA sequences of SEQ ID NO: 3, 69, 112-124 or the complement thereto. In a second preferred aspect, the cDNA sequence is the mouse cDNA sequence of SEQ ID NO: 184. Optionally, said polynucleotide consists of, consists essentially of, or comprises a contiguous span of nucleotides of a mammalian genomic sequence, preferably a sequence selected  
5 the following SEQ ID NOs: 3, 69, 112-124 and 184, wherein said contiguous span is at least 6, 8, 10, 12, 15, 20, 25, 30, 50, 100, 200, or 500 nucleotides in length.

A third embodiment of the present invention is a recombinant, purified or isolated polynucleotide, or the complement thereof, encoding a mammalian PG1 protein, or a fragment thereof. In one aspect the PG1 protein sequence is from a human, mouse or other mammal. In a  
10 preferred aspect, the PG1 protein sequence is selected from the human PG1 protein sequences of SEQ ID NO: 4, 5, 70, and 125-136. In a second preferred aspect, the PG1 protein sequence is the mouse PG1 protein sequences of SEQ ID NO: 74. Optionally, said fragment of PG1 polypeptide consists of, consists essentially of, or comprises a contiguous stretch of at least 8, 10, 12, 15, 20, 25, 30, 50, 100 or 200 amino acids from SEQ ID NOs: 4, 5, 70, 74, and 125-136, as well as any other human, mouse  
15 or mammalian PG1 polypeptide.

A fourth embodiment of the invention are the polynucleotide primers and probes disclosed herein

A fifth embodiment of the present invention is a recombinant, purified or isolated polypeptide comprising or consisting of a mammalian PG1 protein, or a fragment thereof. In one aspect the PG1  
20 protein sequence is from a human, mouse or other mammal. In a preferred aspect, the PG1 protein sequence is selected from the human PG1 protein sequences of SEQ ID NO: 4, 5, 70, and 125-136. In a second preferred aspect, the PG1 protein sequence is the mouse PG1 protein sequences of SEQ ID NO: 74. Optionally, said fragment of PG1 polypeptide consists of, consists essentially of, or comprises a contiguous stretch of at least 8, 10, 12, 15, 20, 25, 30, 50, 100 or 200 amino acids from  
25 SEQ ID NOs: 4, 5, 70, 74, and 125-136, as well as any other human, mouse or mammalian PG1 polypeptide.

A sixth embodiment of the present invention is an antibody composition capable of specifically binding to a polypeptide of the invention. Optionally, said antibody is polyclonal or monoclonal. Optionally, said polypeptide is an epitope-containing fragment of at least 8, 10, 12, 15,  
30 20, 25, or 30 amino acids of a human, mouse, or mammalian PG1 protein, preferably a sequence selected from SEQ ID NOs: 4, 5, 70, 74, or 125-136.

A seventh embodiment of the present invention is a vector comprising any polynucleotide of the invention. Optionally, said vector is an expression vector, gene therapy vector, amplification vector, gene targeting vector, or knock-out vector.

35 An eighth embodiment of the present invention is a host cell comprising any vector of the invention.

A ninth embodiment of the present invention is a mammalian host cell comprising a PG1 gene disrupted by homologous recombination with a knock out vector.

A tenth embodiment of the present invention is a nonhuman host mammal or animal comprising a vector of the invention.

5 A further embodiment of the present invention is a nonhuman host mammal comprising a PG1 gene disrupted by homologous recombination with a knock out vector.

Another embodiment of the present invention is a method of determining whether an individual is at risk of developing cancer or prostate cancer at a later date or whether the individual suffers from cancer or prostate cancer as a result of a mutation in the PG1 gene comprising obtaining  
10 a nucleic acid sample from the individual; and determining whether the nucleotides present at one or more of the PG1-related biallelic markers of the invention are indicative of a risk of developing prostate cancer at a later date or indicative of prostate cancer resulting from a mutation in the PG1 gene. Optionally, said PG1-related biallelic is a PG1-related biallelic markers positioned in SEQ ID NO: 179; a PG1-related biallelic marker selected from the group consisting of 99-1485/251, 99-  
15 622/95, 99-619/141, 4-76/222, 4-77/151, 4-71/233, 4-72/127, 4-73/134, 99-610/250, 99-609/225, 4-90/283, 99-602/258, 99-600/492, 99-598/130, 99-217/277, 99-576/421, 4-61/269, 4-66/145, and 4-67/40; or a PG1-related biallelic marker selected from the group consisting of 99-622, 4-77, 4-71, 4-73, 99-598, 99-576, and 4-66.

Another embodiment of the present invention is a method of determining whether an  
20 individual is at risk of developing prostate cancer at a later date or whether the individual suffers from prostate cancer as a result of a mutation in the PG1 gene comprising obtaining a nucleic acid sample from the individual and determining whether the nucleotides present at one or more of the polymorphic bases in a PG1-related biallelic marker. Optionally, said PG1-related biallelic is a PG1-related biallelic markers positioned in SEQ ID NO: 179; a PG1-related biallelic marker selected from  
25 the group consisting of 99-1485/251, 99-622/95, 99-619/141, 4-76/222, 4-77/151, 4-71/233, 4-72/127, 4-73/134, 99-610/250, 99-609/225, 4-90/283, 99-602/258, 99-600/492, 99-598/130, 99-217/277, 99-576/421, 4-61/269, 4-66/145, and 4-67/40; or a PG1-related biallelic marker selected from the group consisting of 99-622, 4-77, 4-71, 4-73, 99-598, 99-576, and 4-66.

Another embodiment of the present invention is a method of obtaining an allele of the PG1  
30 gene which is associated with a detectable phenotype comprising obtaining a nucleic acid sample from an individual expressing the detectable phenotype, contacting the nucleic acid sample with an agent capable of specifically detecting a nucleic acid encoding the PG1 protein, and isolating the nucleic acid encoding the PG1 protein. In one aspect of this method, the contacting step comprises contacting the nucleic acid sample with at least one nucleic acid probe capable of specifically  
35 hybridizing to said nucleic acid encoding the PG1 protein. In another aspect of this embodiment, the contacting step comprises contacting the nucleic acid sample with an antibody capable of specifically

binding to the PG1 protein. In another aspect of this embodiment, the step of obtaining a nucleic acid sample from an individual expressing a detectable phenotype comprises obtaining a nucleic acid sample from an individual suffering from prostate cancer.

Another embodiment of the present invention is a method of obtaining an allele of the PG1 gene which is associated with a detectable phenotype comprising obtaining a nucleic acid sample from an individual expressing the detectable phenotype, contacting the nucleic acid sample with an agent capable of specifically detecting a sequence within the 8p23 region of the human genome, identifying a nucleic acid encoding the PG1 protein in the nucleic acid sample, and isolating the nucleic acid encoding the PG1 protein. In one aspect of this embodiment, the nucleic acid sample is obtained from an individual suffering from cancer or prostate cancer.

Another embodiment of the present invention is a method of categorizing the risk of prostate cancer in an individual comprising the step of assaying a sample taken from the individual to determine whether the individual carries an allelic variant of PG1 associated with an increased risk of prostate cancer. In one aspect of this embodiment, the sample is a nucleic acid sample. In another aspect a nucleic acid sample is assayed by determining the frequency of the PG1 transcripts present. In another aspect of this embodiment, the sample is a protein sample. In another aspect of this embodiment, the method further comprises determining whether the PG1 protein in the sample binds an antibody specific for a PG1 isoform associated with prostate cancer.

Another embodiment of the present invention is a method of categorizing the risk of prostate cancer in an individual comprising the step of determining whether the identities of the polymorphic bases of one or more biallelic markers which are in linkage disequilibrium with the PG1 gene are indicative of an increased risk of prostate cancer.

Another embodiment of the present invention comprises a method of identifying molecules which specifically bind to a PG1 protein, preferably the protein of SEQ ID NO:4 or a portion thereof: comprising the steps of introducing a nucleic acid encoding the protein of SEQ ID NO:4 or a portion thereof into a cell such that the protein of SEQ ID NO:4 or a portion thereof contacts proteins expressed in the cell and identifying those proteins expressed in the cell which specifically interact with the protein of SEQ ID NO:4 or a portion thereof.

Another embodiment of the present invention is a method of identifying molecules which specifically bind to the protein of SEQ ID NO: 4 or a portion thereof. One step of the method comprises linking a first nucleic acid encoding the protein of SEQ ID NO:4 or a portion thereof to a first indicator nucleic acid encoding a first indicator polypeptide to generate a first chimeric nucleic acid encoding a first fusion protein. The first fusion protein comprises the protein of SEQ ID NO:4 or a portion thereof and the first indicator polypeptide. Another step of the method comprises linking a second nucleic acid encoding a test polypeptide to a second indicator nucleic acid encoding a second indicator polypeptide to generate a second chimeric nucleic acid encoding a second

fusion protein. The second fusion protein comprises the test polypeptide and the second indicator polypeptide. Association between the first indicator protein and the second indicator protein produces a detectable result. Another step of the method comprises introducing the first chimeric nucleic acid and the second chimeric nucleic acid into a cell. Another step comprises detecting the detectable  
5 result.

A further embodiment of the invention is a purified or isolated mammalian PG1 gene or cDNA sequence.

Further embodiments of the present invention include the nucleic acid and amino acid sequences of mutant or low frequency PG1 alleles derived from prostate cancer patients, tissues or  
10 cell lines. The present invention also encompasses methods which utilize detection of these mutant PG1 sequences in an individual or tissue sample to diagnosis prostate cancer, assess the risk of developing prostate cancer or assess the likely severity of a particular prostate tumor.

Another embodiment of the invention encompasses any polynucleotide of the invention attached to a solid support. In addition, the polynucleotides of the invention which are attached to a  
15 solid support encompass polynucleotides with any further limitation described in this disclosure, or those following: Optionally, said polynucleotides is specified as attached individually or in groups of at least 2, 5, 8, 10, 12, 15, 20, or 25 distinct polynucleotides of the inventions to a single solid support. Optionally, polynucleotides other than those of the invention may attached to the same solid support as polynucleotides of the invention. Optionally, when multiple polynucleotides are attached to a solid  
20 support they are attached at random locations, or in an ordered array. Optionally, said ordered array is addressable.

An additional embodiment of the invention encompasses the use of any polynucleotide for, or any polynucleotide for use in, determining the identity of an allele at a PG1-related biallelic marker. In addition, the polynucleotides of the invention for use in determining the identity of an allele at a  
25 PG1-related biallelic marker encompass polynucleotides with any further limitation described in this disclosure, or those following: Optionally, said PG1-related biallelic marker is a PG1-related biallelic markers positioned in SEQ ID NO: 179; a PG1-related biallelic marker selected from the group consisting of 99-1485/251, 99-622/95, 99-619/141, 4-76/222, 4-77/151, 4-71/233, 4-72/127, 4-73/134, 99-610/250, 99-609/225, 4-90/283, 99-602/258, 99-600/492, 99-598/130, 99-217/277, 99-576/421, 4-  
30 61/269, 4-66/145, and 4-67/40; or a PG1-related biallelic marker selected from the group consisting of 99-622, 4-77, 4-71, 4-73, 99-598, 99-576, and 4-66. Optionally, said polynucleotide may comprise a sequence disclosed in the present specification. Optionally, said polynucleotide may consist of, or consist essentially of any polynucleotide described in the present specification. Optionally, said determining is performed in a hybridization assay, sequencing assay, microsequencing assay, or allele-specific amplification assay. Optionally, said polynucleotide is  
35 attached to a solid support, array, or addressable array. Optionally, said polynucleotide is labeled.

Another embodiment of the invention encompasses the use of any polynucleotide for, or any polynucleotide for use in, amplifying a segment of nucleotides comprising an PG1-related biallelic marker. In addition, the polynucleotides of the invention for use in amplifying a segment of nucleotides comprising a PG1-related biallelic marker encompass polynucleotides with any further  
5 limitation described in this disclosure, or those following: Optionally, said PG1-related biallelic marker is a PG1-related biallelic markers positioned in SEQ ID NO: 179; a PG1-related biallelic marker selected from the group consisting of 99-1485/251, 99-622/95, 99-619/141, 4-76/222, 4-77/151, 4-71/233, 4-72/127, 4-73/134, 99-610/250, 99-609/225, 4-90/283, 99-602/258, 99-600/492, 99-598/130, 99-217/277, 99-576/421, 4-61/269, 4-66/145, and 4-67/40; or a PG1-related biallelic  
10 marker selected from the group consisting of 99-622, 4-77, 4-71, 4-73, 99-598, 99-576, and 4-66. Optionally, said polynucleotide may comprise a sequence disclosed in the present specification. Optionally, said polynucleotide may consist of, or consist essentially of any polynucleotide described in the present specification. Optionally, said amplifying is performed by a PCR or LCR. Optionally, said polynucleotide is attached to a solid support, array, or addressable array. Optionally, said  
15 polynucleotide is labeled.

A further embodiment of the invention encompasses methods of genotyping a biological sample comprising determining the identity of an allele at an PG1-related biallelic marker. In addition, the genotyping methods of the invention encompass methods with any further limitation described in this disclosure, or those following: Optionally, said PG1-related biallelic marker is a PG1-related  
20 biallelic markers positioned in SEQ ID NO: 179; a PG1-related biallelic marker selected from the group consisting of 99-1485/251, 99-622/95, 99-619/141, 4-76/222, 4-77/151, 4-71/233, 4-72/127, 4-73/134, 99-610/250, 99-609/225, 4-90/283, 99-602/258, 99-600/492, 99-598/130, 99-217/277, 99-576/421, 4-61/269, 4-66/145, and 4-67/40; or a PG1-related biallelic marker selected from the group consisting of 99-622, 4-77, 4-71, 4-73, 99-598, 99-576, and 4-66. Optionally, said method further  
25 comprises determining the identity of a second allele at said biallelic marker, wherein said first allele and second allele are not base paired (by Watson & Crick base pairing) to one another. Optionally, said biological sample is derived from a single individual or subject. Optionally, said method is performed in vitro. Optionally, said biallelic marker is determined for both copies of said biallelic marker present in said individual's genome. Optionally, said biological sample is derived from  
30 multiple subjects or individuals. Optionally, said method further comprises amplifying a portion of said sequence comprising the biallelic marker prior to said determining step. Optionally, wherein said amplifying is performed by PCR, LCR, or replication of a recombinant vector comprising an origin of replication and said portion in a host cell. Optionally, wherein said determining is performed by a hybridization assay, sequencing assay, microsequencing assay, or allele-specific amplification assay.

35 An additional embodiment of the invention comprises methods of estimating the frequency of an allele in a population comprising determining the proportional representation of an allele at a PG1-



related biallelic marker in said population. In addition, the methods of estimating the frequency of an allele in a population of the invention encompass methods with any further limitation described in this disclosure, or those following: Optionally, said PG1-related biallelic marker is a PG1-related biallelic markers positioned in SEQ ID NO: 179; a PG1-related biallelic marker selected from the group consisting of 99-1485/251, 99-622/95, 99-619/141, 4-76/222, 4-77/151, 4-71/233, 4-72/127, 4-73/134, 99-610/250, 99-609/225, 4-90/283, 99-602/258, 99-600/492, 99-598/130, 99-217/277, 99-576/421, 4-61/269, 4-66/145, and 4-67/40; or a PG1-related biallelic marker selected from the group consisting of 99-622, 4-77, 4-71, 4-73, 99-598, 99-576, and 4-66. Optionally, determining the proportional representation of an allele at a PG1-related biallelic marker is accomplished by determining the identity of the alleles for both copies of said biallelic marker present in the genome of each individual in said population and calculating the proportional representation of said allele at said PG1-related biallelic marker for the population. Optionally, determining the proportional representation is accomplished by performing a genotyping method of the invention on a pooled biological sample derived from a representative number of individuals, or each individual, in said population, and calculating the proportional amount of said nucleotide compared with the total.

A further embodiment of the invention comprises methods of detecting an association between a genotype and a phenotype, comprising the steps of a) genotyping at least one PG1-related biallelic marker in a trait positive population according to a genotyping method of the invention; b) genotyping said PG1-related biallelic marker in a control population according to a genotyping method of the invention; and c) determining whether a statistically significant association exists between said genotype and said phenotype. In addition, the methods of detecting an association between a genotype and a phenotype of the invention encompass methods with any further limitation described in this disclosure, or those following: Optionally, said PG1-related biallelic marker is a PG1-related biallelic markers positioned in SEQ ID NO: 179; a PG1-related biallelic marker selected from the group consisting of 99-1485/251, 99-622/95, 99-619/141, 4-76/222, 4-77/151, 4-71/233, 4-72/127, 4-73/134, 99-610/250, 99-609/225, 4-90/283, 99-602/258, 99-600/492, 99-598/130, 99-217/277, 99-576/421, 4-61/269, 4-66/145, and 4-67/40; or a PG1-related biallelic marker selected from the group consisting of 99-622, 4-77, 4-71, 4-73, 99-598, 99-576, and 4-66. Optionally, said control population is a trait negative population, or a random population. Optionally, each of said genotyping steps a) and b) is performed on a single pooled biological sample derived from each of said populations. Optionally, each of said genotyping of steps a) and b) is performed separately on biological samples derived from each individual in said population or a subsample thereof. Optionally, said phenotype is a disease, cancer or prostate cancer; a response to an anti-cancer agent or an anti-prostate cancer agent; or a side effect to an anti-cancer or anti-prostate cancer agent. Optionally, said method comprises the additional steps of determining the phenotype in said trait positive and said control populations prior to step c).

An additional embodiment of the present invention encompasses methods of estimating the frequency of a haplotype for a set of biallelic markers in a population, comprising the steps of: a) genotyping at least one PG1-related biallelic marker for both copies of said set of biallelic marker present in the genome of each individual in said population or a subsample thereof, according to a genotyping method of the invention; b) genotyping a second biallelic marker by determining the identity of the allele at said second biallelic marker for both copies of said second biallelic marker present in the genome of each individual in said population or said subsample, according to a genotyping method of the invention; and c) applying a haplotype determination method to the identities of the nucleotides determined in steps a) and b) to obtain an estimate of said frequency. In addition, the methods of estimating the frequency of a haplotype of the invention encompass methods with any further limitation described in this disclosure, or those following: Optionally, said PG1-related biallelic marker is a PG1-related biallelic markers positioned in SEQ ID NO: 179; a PG1-related biallelic marker selected from the group consisting of 99-1485/251, 99-622/95, 99-619/141, 4-76/222, 4-77/151, 4-71/233, 4-72/127, 4-73/134, 99-610/250, 99-609/225, 4-90/283, 99-602/258, 99-600/492, 99-598/130, 99-217/277, 99-576/421, 4-61/269, 4-66/145, and 4-67/40; or a PG1-related biallelic marker selected from the group consisting of 99-622, 4-77, 4-71, 4-73, 99-598, 99-576, and 4-66. Optionally, said second biallelic marker is a PG1-related biallelic marker; a PG1-related biallelic markers positioned in SEQ ID NO: 179; a PG1-related biallelic marker selected from the group consisting of 99-1485/251, 99-622/95, 99-619/141, 4-76/222, 4-77/151, 4-71/233, 4-72/127, 4-73/134, 99-610/250, 99-609/225, 4-90/283, 99-602/258, 99-600/492, 99-598/130, 99-217/277, 99-576/421, 4-61/269, 4-66/145, and 4-67/40; or a PG1-related biallelic marker selected from the group consisting of 99-622, 4-77, 4-71, 4-73, 99-598, 99-576, and 4-66. Optionally, said PG1-related biallelic marker and said second biallelic marker are 4-77/151 and 4-66/145. Optionally, said haplotype determination method is an expectation-maximization algorithm.

An additional embodiment of the present invention encompasses methods of detecting an association between a haplotype and a phenotype, comprising the steps of: a) estimating the frequency of at least one haplotype in a trait positive population, according to a method of the invention for estimating the frequency of a haplotype; b) estimating the frequency of said haplotype in a control population, according to a method of the invention for estimating the frequency of a haplotype; and c) determining whether a statistically significant association exists between said haplotype and said phenotype. In addition, the methods of detecting an association between a haplotype and a phenotype of the invention encompass methods with any further limitation described in this disclosure, or those following: Optionally, said PG1-related biallelic is a PG1-related biallelic markers positioned in SEQ ID NO: 179; a PG1-related biallelic marker selected from the group consisting of 99-1485/251, 99-622/95, 99-619/141, 4-76/222, 4-77/151, 4-71/233, 4-72/127, 4-73/134, 99-610/250, 99-609/225, 4-90/283, 99-602/258, 99-600/492, 99-598/130, 99-217/277, 99-576/421, 4-

61/269, 4-66/145, and 4-67/40; or a PG1-related biallelic marker selected from the group consisting of 99-622, 4-77, 4-71, 4-73, 99-598, 99-576, and 4-66. Optionally, said PG1-related biallelic marker and said second biallelic marker are 4-77/151 and 4-66/145. Optionally, said haplotype exhibits a p-value of  $< 1 \times 10^{-3}$  in an association with a trait positive population with cancer, preferably prostate cancer. Optionally, said control population is a trait negative population, or a random population. Optionally, said phenotype is a disease, cancer or prostate cancer; a response to an anti-cancer agent or an anti-prostate cancer agent, or a side effects to an anti-cancer or anti-prostate cancer agent. Optionally, said method comprises the additional steps of determining the phenotype in said trait positive and said control populations prior to step c).

10 Additional embodiments and aspects of the present invention are set forth in the Detailed Description of the Invention and the Examples.

#### Brief Description of the Drawings

Figure 1 is a diagram showing the BAC contig containing the PG1 gene and the positions of biallelic markers along the contig.

Figure 2 is a graph showing the results of the first screening of a prostate cancer association study and the significance of various biallelic markers as measured by their chi squared and p-values for a low density set of markers.

Figure 3 is a graph showing the results of the first screening of a prostate cancer association study and the significance of various biallelic markers as measured by their chi squared and p-values for a higher density set of markers.

Figure 4 is a table demonstrating the results of an haplotype analysis. Among all the theoretical potential different haplotypes based on 2 to 9 markers, 11 haplotypes showing a strong association with prostate cancer were selected, and their haplotype analysis results are shown here.

Figure 5 is a bar graph demonstrating the results of an experiment evaluating the significance (p-values) of the haplotype analysis shown in Figure 4.

Figure 6A is a table listing the biallelic markers used in the haplotype analysis of Figure 4. Figure 6B is a table listing additional biallelic markers in linkage disequilibrium with the PG1 gene.

Figure 7 is a table listing the positions of exons, splice sites, a stop codon, and a poly A site in the PG1 gene.

Figure 8A is a diagram showing the genomic structure of PG1 in comparison with its most abundant mRNA transcript. Figure 8B is a more detailed diagram showing the genomic structure of PG1, including exons and introns.

Figure 9 is a table listing some of the homologies between the PG1 protein and known proteins.

Figure 10 is a half-tone reproduction of a fluorescence micrograph of the perinuclear/nuclear

expression of PG1 in tumoral (PC3) and normal prostatic cell lines (PNT2). Vector "PG1": includes all the coding exons from exon 1 to 8. For PC3 (upper panel) and PNT2 (lower panel), the nucleus was labelled with Propidium iodide (IP, left panel). Note that EGFP fluorescence was detected in and around the nucleus (GFP, middle panel), as shown when the two pictures were overlapped (right panel).

Figure 11 is a half-tone reproduction of a fluorescence micrograph of the perinuclear/nuclear expression of PG1/1-4 in tumoral (PC3) and normal prostatic cell lines (PNT2). Vector "PG1/1-4" corresponds to an alternative messenger which is due to an alternative splicing, joining exon 1 to exon 4, and resulting in the absence of exons 2 and 3. For PC3 (upper panel) and PNT2 (lower panel), the nucleus was labelled with Propidium iodide (IP, left panel). Note that EGFP fluorescence was detected in and around the nucleus (GFP, middle panel), as shown when the two pictures were overlapped (right panel).

Figure 12 is a half-tone reproduction of a fluorescence micrograph of the perinuclear/nuclear expression of PG1/1-5 in tumoral prostatic cell line (PC3) and cytoplasmic expression of PG1/1-5 in normal prostatic cell line (PNT2). Vector "PG1/1-5" corresponds to an alternative messenger which is due to an alternative splicing, joining exon 1 to exon 5, and resulting in the absence of exons 2, 3 and 4. For PC3 (upper panel) and PNT2 (lower panels), the nucleus was labelled with Propidium iodide (IP). Note that in PC3 cells, EGFP fluorescence was detected in and around the nucleus (GFP, upper middle panel), as shown when the two picture were overlapped (upper right panel). In PNT2A cells, EGFP fluorescence was detected in the cytoplasm (GFP, lower left panel), as shown when the two pictures were overlapped (lower right panel).

Figure 13 is a half-tone reproduction of a fluorescence micrograph of the perinuclear/nuclear expression of a mutated form PG1 (PG1mut229) in normal prostatic cell line (PNT2). Vector "PG1/1-7" includes exons 1 to 6, and corresponds to the mutated form identified in genomic DNA of the prostatic tumoural cell line LNCaP. The nucleus was labelled with Propidium iodide (IP, left panel). EGFP fluorescence was detected in the cytoplasm (GFP, middle panel), as shown when the two pictures were overlapped (lower right panel).

Figure 14 is a diagram of the structure of the 14 alternative splice species found for human PG1 by the exons present. An \* indicates that there is a stop codon in frame at that location. An arrow to the right at the right-hand side of a splice species indicates that the open-reading frame continues off of the chart. a space between exons indicates that the exon(s) is missing from that particular alternative splice species. An up arrow indicates that either exon 1bis, 3bis, or 5bis has been inserted depending upon which is indicated. A bracket notation in exon 6, over an exon 6bis notation indicates that the first 60 bases is missing from exon 6, and exon 6bis is therefor present as a truncated form of exon 6.

Figure 15 is a table listing the results of a series of RT-PCR experiments that were performed

on RNA of normal prostate, normal prostatic cell lines (PNT1A, PNT1B and PNT2), and tumoral prostatic cell lines (LnCaPFCG, LnCaPIMB, CaHPV, Du145, PC3, and prostate tumors (ECP5 to ECP24) using all the possible combinations of primers (SEQ ID NOs: 137-178) specific to all of the possible splice junctions or exon borders in human PG1. An NT indicates that the experiment was not performed. An [+] indicates the use of an alternative splice species with exons 1, 3, 4, 7, and 8.

Figure 16 is a graph showing the results of association studies using markers spanning the 650 kb region of the 8p23 locus around PG1, using both single point analysis and haplotyping studies.

Figure 17 is a graph showing an enlarged view of the single point association results within a 160 kb region comprising the PG1 gene.

Figure 18A is a graph showing an enlarged view of the single point association results of 40 kb within the PG1 gene. Figure 18B is a table listing the location of markers within PG1 gene, the two possible alleles at each site. For each marker, the disease-associated allele is indicated first; its frequencies in cases and controls as well as the difference between both are shown; the odd-ratio and the p-value of each individual marker association are also shown.

Figure 19A is a table showing the results of a haplotype analysis study using 4 markers (marker Nos. 4-14, 99-217, 4-66 and 99-221) within the 160 kb region shown in Figure 17. Figure 19B is a table showing the segmented haplotyping results according to the subject's age, and whether the prostate cancer cases were sporadic or familial, using the same markers 4 markers and the same individuals as were used to generate the results in Figure 19A.

Figure 20 is a table listing the haplotyping results and odd ratios for combinations of the 7 markers (99-622 ; 4-77 ; 4-71 ; 4-73 ; 99-598 ; 99-576 ; 4-66) within PG1 gene that were shown in Figure 18 to have p-values more significant than  $1.10^{-2}$ . All of the 2-, 3-, 4-, 5-, 6- and 7-marker haplotypes were tested.

Figure 21 is a graph showing the distribution of statistical significance, as measured by Chi-square values, for each series of possible x-marker haplotypes, (x =2, 3 or 4) using all of the 19 markers listed in Figure 18B.

#### Detailed Description of the Preferred Embodiment

The practice of the present invention encompasses conventional techniques of chemistry, immunology, molecular biology, biochemistry, protein chemistry, and recombinant DNA technology, which are within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Oligonucleotide Synthesis (M. Gait ed. 1984); Nucleic Acid Hybridization (B. Hames & S. Higgins, eds., 1984); Sambrook, Fritsch & Maniatis, Molecular Cloning: A Laboratory Manual, Second Edition (1989); PCR Technology (H.A. Erlich ed., Stockton Press); R. Scope, Protein Purification Principles and Practice (Springer-Verlag); and the series Methods in Enzymology (S. Colowick and N. Kaplan eds., Academic Press, Inc.).

### Definitions

As used interchangeably herein, the terms "nucleic acid" "oligonucleotide", and "polynucleotides" include RNA, DNA, or RNA/DNA hybrid sequences of more than one nucleotide in either single chain or duplex form. The term "nucleotide" as used herein as an adjective to describe molecules comprising RNA, DNA, or RNA/DNA hybrid sequences of any length in single-stranded or duplex form. The term "nucleotide" is also used herein as a noun to refer to individual nucleotides or varieties of nucleotides, meaning a molecule, or individual unit in a larger nucleic acid molecule, comprising a purine or pyrimidine, a ribose or deoxyribose sugar moiety, and a phosphate group, or phosphodiester linkage in the case of nucleotides within an oligonucleotide or polynucleotide. Although the term "nucleotide" is also used herein to encompass "modified nucleotides" which comprise at least one modifications (a) an alternative linking group, (b) an analogous form of purine, (c) an analogous form of pyrimidine, or (d) an analogous sugar, for examples of analogous linking groups, purine, pyrimidines, and sugars see for example PCT publication No. WO 95/04064. However, the polynucleotides of the invention are preferably comprised of greater than 50% conventional deoxyribose nucleotides, and most preferably greater than 90% conventional deoxyribose nucleotides. The polynucleotide sequences of the invention is prepared by any known method, including synthetic, recombinant, ex vivo generation, or a combination thereof, as well as utilizing any purification methods known in the art.

As used herein, the term "purified" does not require absolute purity; rather, it is intended as a relative definition Purification of starting material or natural material to at least one order of magnitude, preferably two or three orders, and more preferably four or five orders of magnitude is expressly contemplated.

The term "purified" is used herein to describe a polynucleotide or polynucleotide vector of the invention which has been separated from other compounds including, but not limited to other nucleic acids, carbohydrates, lipids and proteins (such as the enzymes used in the synthesis of the polynucleotide), or the separation of covalently closed polynucleotides from linear polynucleotides. A polynucleotide is substantially pure when at least about 50 %, preferably 60 to 75% of a sample exhibits a single polynucleotide sequence and conformation (linear versus covalently close). A substantially pure polynucleotide typically comprises about 50 %, preferably 60 to 90% weight/weight of a nucleic acid sample, more usually about 95%, and preferably is over about 99% pure. Polynucleotide purity or homogeneity is indicated by a number of means well known in the art, such as agarose or polyacrylamide gel electrophoresis of a sample, followed by visualizing a single polynucleotide band upon staining the gel. For certain purposes higher resolution can be provided by using HPLC or other means well known in the art.

The term "polypeptide" refers to a polymer of amino without regard to the length of the polymer; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide.

This term also does not specify or exclude post-expression modifications of polypeptides, for example, polypeptides which include the covalent attachment of glycosyl groups, acetyl groups, phosphate groups, lipid groups and the like are expressly encompassed by the term polypeptide. Also included within the definition are polypeptides which contain one or more analogs of an amino acid  
5 (including, for example, non-naturally occurring amino acids, amino acids which only occur naturally in an unrelated biological system, modified amino acids from mammalian systems etc.), polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring.

As used herein, the term "isolated" requires that the material be removed from its original  
10 environment (e.g., the natural environment if it is naturally occurring).

The term "purified" is used herein to describe a polypeptide of the invention which has been separated from other compounds including, but not limited to nucleic acids, lipids, carbohydrates and other proteins. A polypeptide is substantially pure when at least about 50 %, preferably 60 to 75% of a sample exhibits a single polypeptide sequence. A substantially pure polypeptide typically comprises  
15 about 50 %, preferably 60 to 90% weight/weight of a protein sample, more usually about 95%, and preferably is over about 99% pure. Polypeptide purity or homogeneity is indicated by a number of means well known in the art, such as agarose or polyacrylamide gel electrophoresis of a sample, followed by visualizing a single polypeptide band upon staining the gel. For certain purposes higher resolution can be provided by using HPLC or other means well known in the art.

As used herein, the term "non-human animal" refers to any non-human vertebrate, birds and more usually mammals, preferably primates, farm animals such as swine, goats, sheep, donkeys, and horses, rabbits or rodents, more preferably rats or mice. As used herein, the term "animal" is used to refer to any vertebrate, preferably a mammal. Both the terms "animal" and "mammal" expressly embrace human subjects unless preceded with the term "non-human".  
20

As used herein, the term "antibody" refers to a polypeptide or group of polypeptides which are comprised of at least one binding domain, where an antibody binding domain is formed from the folding of variable domains of an antibody molecule to form three-dimensional binding spaces with an internal surface shape and charge distribution complementary to the features of an antigenic determinant of an antigen, which allows an immunological reaction with the antigen. Antibodies  
25 include recombinant proteins comprising the binding domains, as well as fragments, including Fab, Fab', F(ab)<sub>2</sub>, and F(ab')<sub>2</sub> fragments.  
30

As used herein, an "antigenic determinant" is the portion of an antigen molecule, in this case an PG1 polypeptide, that determines the specificity of the antigen-antibody reaction. An "epitope" refers to an antigenic determinant of a polypeptide. An epitope can comprise as few as 3 amino acids  
35 in a spatial conformation which is unique to the epitope. Generally an epitope consists of at least 6 such amino acids, and more usually at least 8-10 such amino acids. Methods for determining the

amino acids which make up an epitope include x-ray crystallography, 2-dimensional nuclear magnetic resonance, and epitope mapping e.g. the Pepscan method described by H. Mario Geysen et al. 1984. Proc. Natl. Acad. Sci. U.S.A. 81:3998-4002; PCT Publication No. WO 84/03564; and PCT Publication No. WO 84/03506.

5       The term "DNA construct" and "vector" are used herein to mean a purified or isolated polynucleotide that has been artificially designed and which comprises at least two nucleotide sequences that are not found as contiguous nucleotide sequences in their natural environment.

      The terms "trait" and "phenotype" are used interchangeably herein and refer to any visible, detectable or otherwise measurable property of an organism such as symptoms of, or susceptibility to  
10 a disease for example. Typically the terms "trait" or "phenotype" are used herein to refer to symptoms of, or susceptibility to cancer or prostate cancer; or to refer to an individual's response to an anti-cancer agent or an anti-prostate cancer agent; or to refer to symptoms of, or susceptibility to side effects to an anticancer agent or an anti-prostate cancer agent.

      The term "allele" is used herein to refer to variants of a nucleotide sequence. A biallelic  
15 polymorphism has two forms. Typically the first identified allele is designated as the original allele whereas other alleles are designated as alternative alleles. Diploid organisms is homozygous or heterozygous for an allelic form.

      The term "heterozygosity rate" is used herein to refer to the incidence of individuals in a population, which are heterozygous at a particular allele. In a biallelic system the heterozygosity rate  
20 is on average equal to  $2P_a(1-P_a)$ , where  $P_a$  is the frequency of the least common allele. In order to be useful in genetic studies a genetic marker should have an adequate level of heterozygosity to allow a reasonable probability that a randomly selected person will be heterozygous.

      The term "genotype" as used herein refers the identity of the alleles present in an individual or a sample. In the context of the present invention a genotype preferably refers to the description of the  
25 biallelic marker alleles present in an individual or a sample. The term "genotyping" a sample or an individual for a biallelic marker consists of determining the specific allele or the specific nucleotide carried by an individual at a biallelic marker.

      The term "mutation" as used herein refers to a difference in DNA sequence between or among different genomes or individuals which has a frequency below 1%.

30       The term "haplotype" refers to a combination of alleles present in an individual or a sample. In the context of the present invention a haplotype preferably refers to a combination of biallelic marker alleles found in a given individual and which is associated with a phenotype.

      The term "polymorphism" as used herein refers to the occurrence of two or more alternative genomic sequences or alleles between or among different genomes or individuals. "Polymorphic"  
35 refers to the condition in which two or more variants of a specific genomic sequence can be found in a population. A "polymorphic site" is the locus at which the variation occurs. A single nucleotide



polymorphism is a single base pair change. Typically a single nucleotide polymorphism is the replacement of one nucleotide by another nucleotide at the polymorphic site. Deletion of a single nucleotide or insertion of a single nucleotide, also give rise to single nucleotide polymorphisms. In the context of the present invention "single nucleotide polymorphism" preferably refers to a single nucleotide substitution. Typically, between different genomes or between different individuals, the polymorphic site is occupied by two different nucleotides.

The terms "biallelic polymorphism" and "biallelic marker" are used interchangeably herein to refer to a nucleotide polymorphism having two alleles at a fairly high frequency in the population. A "biallelic marker allele" refers to the nucleotide variants present at a biallelic marker site. Usually a biallelic marker is a single nucleotide polymorphism. However, less commonly there are also insertions and deletions of up to 5 nucleotides which constitute biallelic markers for the purposes of the present invention. Typically the frequency of the less common allele of the biallelic markers of the present invention has been validated to be greater than 1%, preferably the frequency is greater than 10%, more preferably the frequency is at least 20% (i.e. heterozygosity rate of at least 0.32), even more preferably the frequency is at least 30% (i.e. heterozygosity rate of at least 0.42). A biallelic marker wherein the frequency of the less common allele is 30% or more is termed a "high quality biallelic marker."

The location of nucleotides in a polynucleotide with respect to the center of the polynucleotide are described herein in the following manner. When a polynucleotide has an odd number of nucleotides, the nucleotide at an equal distance from the 3' and 5' ends of the polynucleotide is considered to be "at the center" of the polynucleotide, and any nucleotide immediately adjacent to the nucleotide at the center, or the nucleotide at the center itself is considered to be "within 1 nucleotide of the center." With an odd number of nucleotides in a polynucleotide any of the five nucleotides positions in the middle of the polynucleotide would be considered to be within 2 nucleotides of the center, and so on. When a polynucleotide has an even number of nucleotides, there would be a bond and not a nucleotide at the center of the polynucleotide. Thus, either of the two central nucleotides would be considered to be "within 1 nucleotide of the center" and any of the four nucleotides in the middle of the polynucleotide would be considered to be "within 2 nucleotides of the center", and so on.

The term "upstream" is used herein to refer to a location which is toward the 5' end of the polynucleotide from a specific reference point.

The terms "base paired" and "Watson & Crick base paired" are used interchangeably herein to refer to nucleotides which can be hydrogen bonded to one another by virtue of their sequence identities in a manner like that found in double-helical DNA with thymine or uracil residues linked to adenine residues by two hydrogen bonds and cytosine and guanine residues linked by three hydrogen bonds (See Stryer, L., *Biochemistry*, 4<sup>th</sup> edition, 1995).

The terms "complementary" or "complement thereof" are used herein to refer to the sequences of polynucleotides which is capable of forming Watson & Crick base pairing with another specified polynucleotide throughout the entirety of the complementary region. This term is applied to pairs of polynucleotides based solely upon their sequences and not any particular set of conditions under which the two polynucleotides would actually bind.

As used herein the term "PG1-related biallelic marker" relates to a set of biallelic markers in linkage disequilibrium with PG1. The term PG1-related biallelic marker includes all of the biallelic markers used in the initial association studies shown below in Section I.D., including those biallelic markers contained in SEQ ID NOs: 21-38 and 57-62. The term PG1-related biallelic marker encompasses all of the following polymorphisms positioned in SEQ ID 179, and listed by internal reference number, including:

- 5-63-169 G or C in position 2159;
- 5-63-453 C or T in position 2443; 99-622-95 T or C in position 4452;
- 99-621-215 T or C in position 5733; 99-619-141 G or A in position 8438;
- 4-76-222 deletion of GT in position 11843; 4-76-361 C or T in position 11983;
- 4-77-151 G or C in position 12080; 4-77-294 A or G in position 12221;
- 4-71-33 G or T in position 12947; 4-71-233 A or G in position 13147;
- 4-71-280 G or A in position 13194; 4-71-396 G or C in position 13310;
- 4-72-127 A or G in position 13342; 4-72-152 A or G in position 13367;
- 4-72-380 deletion of A in position 13594; 4-73-134 G or C in position 13680;
- 4-73-356 G or C in position 13902; 99-610-250 T or C in position 16231;
- 99-610-93 A or T in position 16388; 99-609-225 A or T in position 17608;
- 4-90-27 A or C in position 18034; 4-90-283 A or C in position 18290;
- 99-607-397 T or C in position 18786; 99-602-295 deletion of A in position 22835;
- 99-602-258 T or C in position 22872;
- 99-600-492 deletion of TATTG in position 25183;
- 99-600-483 T or G in position 25192; 5-23-288 A or G in position 25614;
- 99-598-130 T or C in position 26911; 99-592-139 A or T in position 32703;
- 99-217-277 C or T in position 34491; 5-47-284 A or G in position 34756;
- 99-589-267 T or G in position 34934; 99-589-41 G or C in position 35160;
- 99-12899-307 C or T in position 39897; 4-12-68 A or G in position 40598;
- 99-582-263 T or C in position 40816; 99-582-132 T or C in position 40947;
- 99-576-421 G or C in position 45783; 4-13-51 C or T in position 47929;
- 4-13-328 A or T in position 48206; 4-13-329 G or C in position 48207;
- 99-12903-381 C or T in position 49282; 5-56-208 A or G in position 50037;
- 5-56-225 A or G in position 50054; 5-56-272 A or G in position 50101;
- 5-56-391 G or T in position 50220; 4-61-269 A or G in position 50440;

- 4-61-391 A or G in position 50562; 4-63-99 A or G in position 50653;  
 4-62-120 A or G in position 50660; 4-62-205 A or G in position 50745;  
 4-64-113 A or T in position 50885; 4-65-104 A or G in position 51249;  
 5-28-300 A or G in position 51333; 5-50-269 C or T in position 51435;  
 5 4-65-324 C or T in position 51468; 5-71-129 G or C in position 51515;  
 5-50-391 G or C in position 51557; 5-71-180 A or G in position 51566;  
 4-67-40 C or T in position 51632; 5-71-280 A or C in position 51666;  
 5-58-167 A or G in position 52016; 5-30-325 C or T in position 52096;  
 5-58-302 A or T in position 52151; 5-31-178 A or G in position 52282;  
 10 5-31-244 A or G in position 52348; 5-31-306 deletion of A in position 52410;  
 5-32-190 C or T in position 52524; 5-32-246 C or T in position 52580;  
 5-32-378 deletion of A in position 52712; 5-53-266 G or C in position 52772;  
 5-60-158 C or T in position 52860; 5-60-390 A or G in position 53092;  
 5-68-272 G or C in position 53272; 5-68-385 A or T in position 53389;  
 15 5-66-53 deletion of GA in position 53511; 5-66-142 G or C in position 53600;  
 5-66-207 A or G in position 53665; 5-37-294 A or G in position 53815;  
 5-62-163 insertion of A in position 54365; 5-62-340 A or T in position 54541; and the compliments thereof. The term PG1-related biallelic marker also includes all of the following biallelic markers listed by internal reference number, and two SEQ ID NOs each of which contains a 47-mers with one  
 20 of the two alternative bases at position 24:  
 4-14-107 of SEQ ID NOs 185 and 262; 4-14-317 of SEQ ID NOs 186 and 263; 4-14-35 of  
 SEQ ID NOs 187 and 264; 4-20-149 of SEQ ID NOs 188 and 265;  
 4-20-77 of SEQ ID NOs 189 and 266; 4-22-174 of SEQ ID NOs 190 and 267;  
 4-22-176 of SEQ ID NOs 191 and 268; 4-26-60 of SEQ ID NOs 192 and 269;  
 25 4-26-72 of SEQ ID NOs 193 and 270; 4-3-130 of SEQ ID NOs 194 and 271;  
 4-38-63 of SEQ ID NOs 195 and 272;  
 4-38-83 of SEQ ID NOs 196 and 273; 4-4-152 of SEQ ID NOs 197 and 274;  
 4-4-187 of SEQ ID NOs 198 and 275; 4-4-288 of SEQ ID NOs 199 and 276;  
 4-42-304 of SEQ ID NOs 200 and 277; 4-42-401 of SEQ ID NOs 201 and 278;  
 30 4-43-328 of SEQ ID NOs 202 and 279; 4-43-70 of SEQ ID NOs 203 and 280;  
 4-50-209 of SEQ ID NOs 204 and 281; 4-50-293 of SEQ ID NOs 205 and 282;  
 4-50-323 of SEQ ID NOs 206 and 283; 4-50-329 of SEQ ID NOs 207 and 284;  
 4-50-330 of SEQ ID NOs 208 and 285; 4-52-163 of SEQ ID NOs 209 and 286;  
 4-52-88 of SEQ ID NOs 210 and 287; 4-53-258 of SEQ ID NOs 211 and 288;  
 35 4-54-283 of SEQ ID NOs 212 and 289; 4-54-388 of SEQ ID NOs 213 and 290;  
 4-55-70 of SEQ ID NOs 214 and 291; 4-55-95 of SEQ ID NOs 215 and 292;

- 4-56-159 of SEQ ID NOs 216 and 293; 4-56-213 of SEQ ID NOs 217 and 294;  
 4-58-289 of SEQ ID NOs 218 and 295; 4-58-318 of SEQ ID NOs 219 and 296;  
 4-60-266 of SEQ ID NOs 220 and 297; 4-60-293 of SEQ ID NOs 221 and 298;  
 4-84-241 of SEQ ID NOs 222 and 299; 4-84-262 of SEQ ID NOs 223 and 300;  
 5 4-86-206 of SEQ ID NOs 224 and 301; 4-86-309 of SEQ ID NOs 225 and 302;  
 4-88-349 of SEQ ID NOs 226 and 303; 4-89-87 of SEQ ID NOs 227 and 304;  
 99-123-184 of SEQ ID NOs 228 and 305; 99-128-202 of SEQ ID NOs 229 and 306;  
 99-128-275 of SEQ ID NOs 230 and 307; 99-128-313 of SEQ ID NOs 231 and 308; 99-128-60 of  
 SEQ ID NOs 232 and 309; 99-12907-295 of SEQ ID NOs 233 and 310; 99-130-58 of SEQ ID NOs  
 10 234 and 311; 99-134-362 of SEQ ID NOs 235 and 312;  
 99-140-130 of SEQ ID NOs 236 and 313; 99-1462-238 of SEQ ID NOs 237 and 314; 99-147-181 of  
 SEQ ID NOs 238 and 315; 99-1474-156 of SEQ ID NOs 239 and 316; 99-1474-359 of SEQ ID  
 NOs 240 and 317; 99-1479-158 of SEQ ID NOs 241 and 318; 99-1479-379 of SEQ ID NOs 242 and  
 319; 99-148-129 of SEQ ID NOs 243 and 320; 99-148-132 of SEQ ID NOs 244 and 321; 99-148-139  
 15 of SEQ ID NOs 245 and 322;  
 99-148-140 of SEQ ID NOs 246 and 323; 99-148-182 of SEQ ID NOs 247 and 324;  
 99-148-366 of SEQ ID NOs 248 and 325; 99-148-76 of SEQ ID NOs 249 and 326;  
 99-1480-290 of SEQ ID NOs 250 and 327; 99-1481-285 of SEQ ID NOs 251 and 328; 99-1484-101 of  
 SEQ ID NOs 252 and 329; 99-1484-328 of SEQ ID NOs 253 and 330; 99-1485-251 of SEQ ID NOs  
 20 254 and 331; 99-1490-381 of SEQ ID NOs 255 and 332; 99-1493-280 of SEQ ID NOs 256 and  
 333; 99-151-94 of SEQ ID NOs 257 and 334;  
 99-211-291 of SEQ ID NOs 258 and 335; 99-213-37 of SEQ ID NOs 259 and 336;  
 99-221-442 of SEQ ID NOs 260 and 337; 99-222-109 of SEQ ID NOs 261 and 338; and the  
 compliments thereof.
- 25 The term "non-genic" is used herein to describe PG1-related biallelic markers, as well as  
 polynucleotides and primers which do not occur in the human PG1 genomic sequence of SEQ ID NO:  
 179. The term "genic" is used herein to describe PG1-related biallelic markers as well as  
 polynucleotides and primers which do occur in the human PG1 genomic sequence of SEQ ID NO:  
 179.
- 30 The terms "an anti-cancer agent" refers to a drug or a compound that is capable of reducing  
 the growth rate, rate of metastasis, or viability of tumor cells in a mammal, is capable of reducing the  
 size or eliminating tumors in a mammal, or is capable of increasing the average life span of a mammal  
 or human with cancer. Anti-cancer agents also include compounds which are able to reduce the risk  
 of cancer developing in a population, particularly a high risk population. The terms "an anti-prostate  
 35 cancer agent" is an anti-cancer agent that has these effects on cells or tumors that are derived from  
 prostate cancer cells.

The terms "response to an anti-cancer agent" and "response to an anti-prostate cancer agent" refer to drug efficacy, including but not limited to ability to metabolize a compound, to the ability to convert a pro-drug to an active drug, and to the pharmacokinetics (absorption, distribution, elimination) and the pharmacodynamics (receptor-related) of a drug in an individual.

5       The terms "side effects to an anti-cancer agent" and "side effects to an anti-prostate cancer agent" refer to adverse effects of therapy resulting from extensions of the principal pharmacological action of the drug or to idiosyncratic adverse reactions resulting from an interaction of the drug with unique host factors. These side effects include, but are not limited to, adverse reactions such as dermatological, hematological or hepatological toxicities and further includes gastric and intestinal  
10 ulceration, disturbance in platelet function, renal injury, nephritis, vasomotor rhinitis with profuse watery secretions, angioneurotic edema, generalized urticaria, and bronchial asthma to laryngeal edema and bronchoconstriction, hypotension, sexual dysfunction, and shock.

As used herein the term "homology" refers to comparisons between protein and/or nucleic acid sequences and is evaluated using any of the variety of sequence comparison algorithms and  
15 programs known in the art. Such algorithms and programs include, but are by no means limited to, TBLASTN, BLASTP, FASTA, TFASTA, and CLUSTALW (Pearson and Lipman, 1988, Proc. Natl. Acad. Sci. USA 85(8):2444-2448; Altschul et al., 1990, J. Mol. Biol. 215(3):403-410; Thompson et al., 1994, Nucleic Acids Res. 22(2):4673-4680; Higgins et al., 1996, Methods Enzymol. 266:383-402; Altschul et al., 1990, J. Mol. Biol. 215(3):403-410; Altschul et al., 1993, Nature Genetics 3:266-272).  
20 In a particularly preferred embodiment, protein and nucleic acid sequence homologies are evaluated using the Basic Local Alignment Search Tool ("BLAST") which is well known in the art (see, e.g., Karlin and Altschul, 1990, Proc. Natl. Acad. Sci. USA 87:2267-2268; Altschul et al., 1990, J. Mol. Biol. 215:403-410; Altschul et al., 1993, Nature Genetics 3:266-272; Altschul et al., 1997, Nuc. Acids Res. 25:3389-3402). In particular, five specific BLAST programs are used to perform the following  
25 task:

- (1) BLASTP and BLAST3 compare an amino acid query sequence against a protein sequence database;
- (2) BLASTN compares a nucleotide query sequence against a nucleotide sequence database;
- 30 (3) BLASTX compares the six-frame conceptual translation products of a query nucleotide sequence (both strands) against a protein sequence database;
- (4) TBLASTN compares a query protein sequence against a nucleotide sequence database translated in all six reading frames (both strands); and

- (5) TBLASTX compares the six-frame translations of a nucleotide query sequence against the six-frame translations of a nucleotide sequence database.

The BLAST programs identify homologous sequences by identifying similar segments, which are referred to herein as "high-scoring segment pairs," between a query amino or nucleic acid sequence and a test sequence which is preferably obtained from a protein or nucleic acid sequence database. High-scoring segment pairs are preferably identified (i.e., aligned) by means of a scoring matrix, many of which are known in the art. Preferably, the scoring matrix used is the BLOSUM62 matrix (Gonnet et al., 1992, Science 256:1443-1445; Henikoff and Henikoff, 1993, Proteins 17:49-61). Less preferably, the PAM or PAM250 matrices may also be used (see, e.g., Schwartz and Dayhoff, eds., 1978, Matrices for Detecting Distance Relationships: Atlas of Protein Sequence and Structure, Washington: National Biomedical Research Foundation). The BLAST programs evaluate the statistical significance of all high-scoring segment pairs identified, and preferably selects those segments which satisfy a user-specified threshold of significance, such as a user-specified percent homology. Preferably, the statistical significance of a high-scoring segment pair is evaluated using the statistical significance formula of Karlin (see, e.g., Karlin and Altschul, 1990, Proc. Natl. Acad. Sci. USA 87:2267-2268).

## **I. ISOLATION AND CHARACTERIZATION OF THE PG1 GENE AND PROTEINS**

### **I.A. The 8p23 Region- LOH Studies: Implications of 8p23 Region in Distinct Cancer Types**

Substantial amounts of LOH data support the hypothesis that genes associated with distinct cancer types are located within 8p23 region of the human genome. Emi et al., demonstrated the implication of 8p23.1-8p21.3 region in cases of hepatocellular carcinoma, colorectal cancer, and non-small cell lung cancer. (Emi M, Fujiwara Y, Nakajima T, Tsuchiya E, Tsuda H, Hirohashi S, Maeda Y, Tsuruta K, Miyaki M, Nakamura Y, Cancer Res. 1992 Oct 1; 52(19): 5368-5372) Yaremko, et al., showed the existence of two major regions of LOH for chromosome 8 markers in a sample of 87 colorectal carcinomas. The most prominent loss was found for 8p23.1-pter, where 45% of informative cases demonstrated loss of alleles. (Yaremko ML, Wasylyshyn ML, Paulus KL, Michelassi F, Westbrook CA, Genes Chromosomes Cancer 1994 May;10(1):1-6). Scholnick et al. demonstrated the existence of three distinct regions of LOH for the markers of chromosome 8 in cases of squamous cell carcinoma of the supraglottic larynx. They showed that the allelic loss of 8p23 marker D8S264 serves as a statistically significant, independent predictor of poor prognosis for patients with supraglottic squamous cell carcinoma. (Scholnick SB, Haughey BH, Sunwoo JB, el-Mofty SK, Baty JD, Piccirillo JF, Zequeira MR, J. Natl. Cancer Inst. 1996 Nov 20; 88(22): 1676-1682 and Sunwoo JB, Holt MS, Radford DM, Deeker C, Scholnick S, Genes Chromosomes Cancer 1996 Jul; 16(3):164-169).

In other studies, Nagai et al. demonstrated the highest loss of heterozygosity in the specific region of 8p23 by genome wide scanning of LOH in 120 cases of hepatocellular carcinoma (HCC). (Nagai H, Pineau P, Tiollais P, Buendia MA, Dejean A, *Oncogene* 1997 Jun 19; 14(24): 2927-2933). Gronwald et al. demonstrated 8p23-pter loss in renal clear cell carcinomas. (Gronwald J, Storkel S, Holtgreve-Grez H, Hadaczek P, Brinkschmidt C, Jauch A, Lubinski J, Cremer, *Cancer Res.* 1997 Feb 1; 57(3): 481-487).

The same region is involved in specific cases of prostate cancer. Matsuyama et al. showed the specific deletion of the 8p23 band in prostate cancer cases, as monitored by FISH with D8S7 probe. (Matsuyama H, Pan Y, Skoog L, Tribukait B, Naito K, Ekman P, Lichter P, Bergerheim US *Oncogene* 1994 Oct; 9(10): 3071-3076). They were able to document a substantial number of cases with deletions of 8p23 but retention of the 8p22 marker LPL. Moreover, Ichikawa et al. deduced the existence of a prostate cancer metastasis suppressor gene and localized it to 8p23-q12 by studies of metastasis suppression in highly metastatic rat prostate cells after transfer of human chromosomes. (Ichikawa T, Nihei N, Kuramochi H, Kawana Y, Killary AM, Rinker-Schaeffer CW, Barrett JC, Isaacs JT, Kugoh H, Oshimura M, Shimazaki J, *Prostate Suppl.* 1996; 6: 31-35).

Recently Washburn et al. were able to find substantial numbers of tumors with the allelic loss specific to 8p23 by LOH studies of 31 cases of human prostate cancer. (Washburn J, Woino K, and Macoska J, *Proceedings of American Association for Cancer Research*, March 1997; 38). In these samples they were able to define the minimal overlapping region with deletions covering genetic interval D8S262-D8S277.

#### Linkage Analysis Studies: Search for Prostate Cancer

##### Linked Regions on Chromosome 8

Microsatellite markers mapping to chromosome 8 were used by the inventors to perform linkage analysis studies on 194 individuals issued from 47 families affected with prostate cancer. While multiple point analysis led to weak linkage results, two point lod score analysis led to non significant results, as shown below.

##### *Two point lod (parametric analysis)*

MARKER	Distance (cM)	Z(lod) scores
D8S1742	0.8	-0.13
D8S561		-0.07

# of families analyzed	47
Total # of individuals genotyped	194
Total # of affected individuals genotyped	122

In view of the non-significant results obtained with linkage analysis, a new mapping approach based on linkage disequilibrium of biallelic markers was utilised to identify genes responsible for sporadic cases of prostate cancer.

### **I.B. Linkage Disequilibrium Using Biallelic Markers To Identify Candidate Loci Responsible**

#### **For Disease**

##### Linkage Disequilibrium

Once a chromosomal region has been identified as potentially harboring a candidate gene associated with a sporadic trait, an excellent approach to refine the candidate gene's location within the identified region is to look for statistical associations between the trait and some marker genotype when comparing an affected (trait<sup>+</sup>) and a control (trait<sup>-</sup>) population.

Association studies have most usually relied on the use of biallelic markers. Biallelic markers are genome-derived polynucleotides that exhibit biallelic polymorphism at one single base position. By definition, the lowest allele frequency of a biallelic polymorphism is 1%; sequence variants that show allele frequencies below 1% are called rare mutations. There are potentially more than 10<sup>7</sup> biallelic markers lying along the human genome.

Association studies seek to establish correlations between traits and genetic markers and are based on the phenomenon of linkage disequilibrium (LD). LD is defined as the trend for alleles at nearby loci on haploid genomes to correlate in the population. If two genetic loci lie on the same chromosome, then sets of alleles on the same chromosomal segment (i.e., haplotypes) tend to be transmitted as a block from generation to generation. When not broken up by recombination, haplotypes can be tracked not only through pedigrees but also through populations. The resulting phenomenon at the population level is that the occurrence of pairs of specific alleles at different loci on the same chromosome is not random, and the deviation from random is called linkage disequilibrium.

Since results generated by association studies are essentially based on the quantitative calculation of allele frequencies, they best apply to the analysis of germline mutations. This is mainly due to the fact that allelic frequencies are difficult to quantify within tumor tissue samples because of the usual presence of normal cells within the studied tumor samples. Association studies applied to cancer genetics will therefore be best suited to the identification of tumor suppressor genes.

##### Trait Localization by Linkage Disequilibrium Mapping

Any gene responsible or partly responsible for a given trait will be in LD with some flanking markers. To map such a gene, specific alleles of these flanking markers which are associated with the gene or genes responsible for the trait are identified. Although the following discussion of techniques for finding the gene or genes associated with a particular trait using linkage disequilibrium mapping, refers to locating a single gene which is responsible for the trait, it will be appreciated that the same techniques may also be used to identify genes which are partially responsible for the trait.



Association studies is conducted within the general population (as opposed to the linkage analysis techniques discussed above which are limited to studies performed on related individuals in one or several affected families).

Association between a biallelic marker A and a trait T may primarily occur as a result of three possible relationships between the biallelic marker and the trait. First, allele *a* of biallelic marker A is directly responsible for trait T (e.g., Apo E e4 allele and Alzheimer's disease). However, since the majority of the biallelic markers used in genetic mapping studies are selected randomly, they mainly map outside of genes. Thus, the likelihood of allele *a* being a functional mutation directly related to trait T is therefore very low.

10 An association between a biallelic marker A and a trait T may also occur when the biallelic marker is very closely linked to the trait locus. In other words, an association occurs when allele *a* is in linkage disequilibrium with the trait-causing allele. When the biallelic marker is in close proximity to a gene responsible for the trait, more extensive genetic mapping will ultimately allow a gene to be discovered near the marker locus which carries mutations in people with trait T (i.e. the gene responsible for the trait or one of the genes responsible for the trait). As will be further exemplified below using a group of biallelic markers which are in close proximity to the gene responsible for the trait, the location of the causal gene can be deduced from the profile of the association curve between the biallelic markers and the trait. The causal gene will be found in the vicinity of the marker showing the highest association with the trait.

20 Finally, an association between a biallelic marker and a trait may occur when people with the trait and people without the trait correspond to genetically different subsets of the population who, coincidentally, also differ in the frequency of allele *a* (population stratification). This phenomenon is avoided by using large heterogeneous samples.

Association studies are particularly suited to the efficient identification of susceptibility genes that present common polymorphisms, and are involved in multifactorial traits whose frequency is relatively higher than that of diseases with monofactorial inheritance.

#### Application of Linkage Disequilibrium Mapping to Candidate Gene Identification

The general strategy of association studies using a set of biallelic markers, is to scan two pools of individuals (affected individuals and unaffected controls) characterized by a well defined phenotype in order to measure the allele frequencies for a number of the chosen markers in each of these pools. If a positive association with a trait is identified using an array of biallelic markers having a high enough density, the causal gene will be physically located in the vicinity of the associated markers, since the markers showing positive association to the trait are in linkage disequilibrium with the trait locus. Regions harboring a gene responsible for a particular trait which

are identified through association studies using high density sets of biallelic markers will, on average, be 20 - 40 times shorter in length than those identified by linkage analysis.

Once a positive association is confirmed as described above, BACs (bacterial artificial chromosomes) obtained from human genomic libraries, constructed as described below, harboring the  
5 markers identified in the association analysis are completely sequenced.

Once a candidate region has been sequenced and analyzed, the functional sequences within the candidate region (exons and promoters, and other potential regulatory regions) are scanned for mutations which are responsible for the trait by comparing the sequences of a selected number of controls and affected individuals using appropriate software. Candidate mutations are further  
10 confirmed by screening a larger number of affected individuals and controls using the microsequencing techniques described below.

Candidate mutations are identified as follows. A pair of oligonucleotide primers is designed in order to amplify the sequences of every predicted functional region. PCR amplification of each predicted functional sequence is carried out on genomic DNA samples from affected patients and  
15 unaffected controls. Amplification products from genomic PCR are subjected to automated dideoxy terminator sequencing reactions and electrophoresed on ABI 377 sequencers. Following gel image analysis and DNA sequence extraction, the sequence data are automatically analyzed to detect the presence of sequence variations among affected cases and unaffected controls. Sequences are systematically verified by comparing the sequences of both DNA strands of each individual.

20 Polymorphisms are then verified by screening a larger population of affected individuals and controls using the microsequencing technique described below in an individual test format. Polymorphisms are considered as candidate mutations when present in affected individuals and controls at frequencies compatible with the expected association results.

#### Association Studies: Statistical Analysis and Haplotyping

25 As mentioned above, linkage analysis typically localizes a disease gene to a chromosomal region of several megabases. Further refinement in location requires the analysis of additional families in order to increase the number of recombinants. However, this approach becomes unfeasible because recombination is rarely observed even within large pedigrees (Boehnke, M, 1994, Am. J. Hum. Genet. 55: 379-390).

30 Linkage disequilibrium, the nonrandom association of alleles at linked loci, may offer an alternative method of obtaining additional recombinants. When a chromosome carrying a mutant allele of a gene responsible for a given trait is first introduced into a population as a result of either mutation or migration, the mutant allele necessarily resides on a chromosome having a unique set of linked markers (haplotype). Consequently, there is complete disequilibrium between these markers  
35 and the disease mutation: the disease mutation is present only linked to a specific set of marker alleles. Through subsequent generations, recombinations occur between the disease mutation and

these marker polymorphisms, resulting in a gradual disappearance of disequilibrium. The degree of disequilibrium dissipation depends on the recombination frequency, so the markers closest to the disease gene will tend to show higher levels of disequilibrium than those that are farther away (Jorde LB, 1995, Am. J. Hum. Genet. 56: 11-14). Because linkage disequilibrium patterns in a present-day  
5 population reflect the action of recombination through many past generations, disequilibrium analysis effectively increases the sample of recombinants. Thus the mapping resolution achieved through the analysis of linkage disequilibrium patterns is much higher than that of linkage analysis.

In practice, in order to define the regions bearing a candidate gene, the affected and control populations are genotyped using an appropriate number of biallelic markers (at a density of 1 marker  
10 every 50-150 kilobases). Then, a marker/trait association study is performed that compares the genotype frequency of each biallelic marker in the affected and control populations by means of a chi square statistical test (one degree of freedom).

After the first screening, additional markers within the region showing positive association are genotyped in the affected and control populations. Two types of complementary analysis are then  
15 performed. First, a marker/trait association study (as described above) is performed to refine the location of the gene responsible for the trait. In addition, a haplotype association analysis is performed to define the frequency and the type of the ancestral/preferential carrier haplotype. Haplotype analysis, by combining the informativeness of a set of biallelic markers increases the power of the association analysis, allowing false positive and/or negative data that may result from the  
20 single marker studies to be eliminated.

The haplotype analysis is performed by estimating the frequencies of all possible haplotypes for a given set of biallelic markers in the case and control populations, and comparing these frequencies by means of a chi square statistical test (one degree of freedom). Haplotype estimations are performed by applying the Expectation-Maximization (EM) algorithm (Excoffier L & Slatkin M,  
25 1995, Mol. Biol. Evol. 12: 921-927), using the EM-HAPLO program (Hawley ME, Pakstis AJ & Kidd KK, 1994, Am. J. Phys. Anthropol. 18: 104). The EM algorithm is used to estimate haplotype frequencies in the case when only genotype data from unrelated individuals are available. The EM algorithm is a generalized iterative maximum likelihood approach to estimation that is useful when data are ambiguous and/or incomplete.

30 The application of biallelic marker based linkage disequilibrium analysis to the 8p23 region to identify a gene associated with prostate cancer is described below.

#### **I.C. Application of Linkage Disequilibrium Mapping to the 8p23 Region**

##### **YAC Contig Construction in 8p23 Region**

First, a YAC contig which contains the 8p23 region was constructed as follows. The CEPH-  
35 Genethon YAC map for the entire human genome (Chumakov I.M. et al. A YAC contig map of the human genome, Nature, 377 Supp.: 175-297, 1995) was used for detailed contig building in the region

around D8S262 and D8S277 genetic markers. Screening data available for regional genetic markers D8S1706, D8S277, D8S1742, D8S518, D8S262, D8S1798, D8S1140, D8S561 and D8S1819 were used to select the following set of CEPH YACs, localized within this region: 832\_g\_12, 787\_c\_11, 920\_h\_7, 807\_a\_1, 842\_b\_1, 745\_a\_3, 910\_d\_3, 879\_f\_11, 918\_c\_6, 764\_c\_7, 910\_f\_12, 967\_c\_11, 856\_d\_8, 792\_a\_6, 812\_h\_4, 873\_c\_8, 930\_a\_2, 807\_a\_1, 852\_d\_10. This set of YACs was tested by PCR with the above mentioned genetic markers as well as with other publicly available markers supposedly located within the 8p23 region. As a result of these studies, a YAC STS contig map was generated around genetic markers D8S262 and D8S277. The two CEPH YACs, 920\_h\_7 (1170 kb insert size) and 910\_f\_12 (1480 kb insert size) constitute a minimal tiling path in this region, with an estimated size of ca. 2 Megabases.

During this mapping effort, the following publicly known STS markers were precisely located within the contig: WI-14718, WI-3831, D8S1413E, WI-8327, WI-3823, ND4.

#### BAC Contig Construction Covering D8S262-D8S277

#### Fragment Within 8p23 Region of the Human Genome

Following construction of the YAC contig, a BAC contig was constructed as follows. BAC libraries were obtained as described in Woo et al. Nucleic Acids Res., 1994, 22, 4922-4931. Briefly, two different whole human genome libraries were produced by cloning BamHI or HindIII partially digested DNA from a lymphoblastoid cell line (derived from individual N°8445, CEPH families) into the pBeloBAC11 vector (Kim et al. Genomics, 1996, 34, 213-218). The library produced with the BamHI partial digestion contained 110,000 clones with an average insert size of 150 kb, which corresponds to 5 human haploid genome equivalents. The library prepared with the HindIII partial digestion corresponds to 3 human genome equivalents with an average insert size of 150 kb.

#### BAC Screening

The human genomic BAC libraries obtained as described above were screened with all of the above mentioned STSs. DNA from the clones in both libraries was isolated and pooled in a three dimensional format ready for PCR screening with the above mentioned STSs using high throughput PCR methods (Chumakov et al., Nature 1995, 377: 175-298). Briefly, three dimensional pooling consists in rearranging the samples to be tested in a manner which allows the number of PCR reactions required to screen the clones with STSs to be reduced by at least 100 fold, as compared to screening each clone individually. PCR amplification products were detected by conventional agarose gel electrophoresis combined with automated image capturing and processing.

In a final step, STS-positive clones were checked individually. Subchromosomal localization of BACs was systematically verified by fluorescence in situ hybridization (FISH), performed on metaphasic chromosomes as described by Cherif et al. Proc. Natl. Acad. Sci. USA 1990, 87: 6639-6643.

BAC insert size was determined by Pulsed Field Gel Electrophoresis after digestion with restriction enzyme NotI.

#### BAC Contig Analysis

The ordered BACs selected by STS screening and verified by FISH, were assembled into  
5 contigs and new markers were generated by partial sequencing of insert ends from some of them. These markers were used to fill the gaps in the contig of BAC clones covering the chromosomal region around D8S277, having an estimated size of 2 megabases. Selected BAC clones from the contig were subcloned and sequenced.

#### BAC Subcloning

10 Each BAC human DNA was first extracted using the alkaline lysis procedure and then sheared by sonication. The obtained DNA fragments were end-repaired and electrophoresed on preparative agarose gels. The fragments in the desired size range were isolated from the gel, purified and ligated to a linearized, dephosphorylated, blunt-ended plasmid cloning vector (pBluescript II Sk (+)). Example 1 describes the BAC subcloning procedure.

15

#### Example 1

The cells obtained from three liters overnight culture of each BAC clone were treated by alkaline lysis using conventional techniques to obtain the BAC DNA containing the genomic DNA inserts. After centrifugation of the BAC DNA in a cesium chloride gradient, ca. 50µg of BAC DNA was purified. 5-10µg of BAC DNA was sonicated using three distinct conditions, to obtain fragments  
20 of the desired size. The fragments were treated in a 50 µl volume with two units of Vent polymerase for 20 min at 70°C, in the presence of the four deoxytriphosphates (100µM). The resulting blunt-ended fragments were separated by electrophoresis on low-melting point 1% agarose gels (60 Volts for 3 hours). The fragments were excised from the gel and treated with agarase. After chloroform extraction and dialysis on Microcon 100 columns, DNA in solution was adjusted to a 100 ng/µl  
25 concentration. A ligation was performed overnight by adding 100 ng of BAC fragmented DNA to 20 ng of pBluescript II Sk (+) vector DNA linearized by enzymatic digestion, and treated by alkaline phosphatase. The ligation reaction was performed in a 10 µl final volume in the presence of 40 units/µl T4 DNA ligase (Epicentre). The ligated products were electroporated into the appropriate cells (ElectroMAX E.coli DH10B cells). IPTG and X-gal were added to the cell mixture, which was  
30 then spread on the surface of an ampicillin-containing agar plate. After overnight incubation at 37°C, recombinant (white) colonies were randomly picked and arrayed in 96 well microplates for storage and sequencing.

#### Partial Sequencing of BACs

At least 30 of the obtained BAC clones were sequenced by the end pair-wise method (500 bp  
35 sequence from each end) using a dye-primer cycle sequencing procedure. Pair-wise sequencing was performed until a map allowing the relative positioning of selected markers along the corresponding

DNA region was established. Example 2 describes the sequencing and ordering of the BAC inserts.

#### Example 2

The subclone inserts were amplified by PCR on overnight bacterial cultures, using vector primers flanking the insertions. The insert extremity sequences (on average 500 bases at each end) were determined by fluorescent automated sequencing on ABI 377 sequencers, with a ABI Prism DNA Sequencing Analysis software (2.1.2 version).

The sequence fragments from BAC subclones were assembled using Gap4 software from R. Staden (Bonfield et al. 1995). This software allows the reconstruction of a single sequence from sequence fragments. The sequence deduced from the alignment of different fragments is called the consensus sequence. We used directed sequencing techniques (primer walking) to complete sequences and link contigs.

Figure 1 shows the overlapping BAC subclones (labeled BAC) which make up the assembled contig and the positions of the publicly known STS markers along the contig.

#### Identification of Biallelic Markers Lying Along the BAC Contig

Following assembly of the BAC contig, biallelic markers lying along the contig were then identified. Given that the assessed distribution of informative biallelic markers in the human genome (biallelic polymorphisms with a heterozygosity rate higher than 42%) is one in 2.5 to 3 kb, six 500 bp genomic fragments have to be screened in order to identify 1 biallelic marker. Six pairs of primers per potential marker, each one defining a ca. 500 bp amplification fragment, were derived from the above mentioned BAC partial sequences. All primers contained a common upstream oligonucleotide tail enabling the easy systematic sequencing of the resulting amplification fragments. Amplification of each BAC-derived sequence was carried out on pools of DNA from ca. 100 individuals. The conditions used for the polymerase chain reaction were optimized so as to obtain more than 95% of PCR products giving 500bp-sequence reads.

The amplification products from genomic PCR using the oligonucleotides derived from the BAC subclones were subjected to automated dideoxy terminator sequencing reactions using a dye-primer cycle sequencing protocol. Following gel image analysis and DNA sequence extraction, sequence data were automatically processed with appropriate software to assess sequence quality and to detect the presence of biallelic sites among the pooled amplified fragments. Biallelic sites were systematically verified by comparing the sequences of both strands of each pool.

The detection limit for the frequency of biallelic polymorphisms detected by sequencing pools of 100 individuals is 0.3 +/- 0.05 for the minor allele, as verified by sequencing pools of known allelic frequencies. Thus, the biallelic markers selected by this method will be "informative biallelic markers" since they have a frequency of 0.3 to 0.5 for the minor allele and 0.5 to 0.7 for the major allele, therefore an average heterozygosity rate higher than 42%.

Example 3 describes the preparation of genomic DNA samples from the individuals screened

to identify biallelic markers.

### Example 3

The population used in order to generate biallelic markers in the region of interest consisted of ca. 100 unrelated individuals corresponding to a French heterogeneous population.

5 DNA was extracted from peripheral venous blood of each donor as follows.

30 ml of blood were taken in the presence of EDTA. Cells (pellet) were collected after centrifugation for 10 minutes at 2000 rpm. Red cells were lysed by a lysis solution (50 ml final volume : 10 mM Tris pH7.6; 5 mM MgCl<sub>2</sub>; 10 mM NaCl). The solution was centrifuged (10 minutes, 2000 rpm) as many times as necessary to eliminate the residual red cells present in the supernatant, 10 after resuspension of the pellet in the lysis solution.

The pellet of white cells was lysed overnight at 42°C with 3.7 ml of lysis solution composed of:

- 3 ml TE 10-2 (Tris-HCl 10 mM, EDTA 2 mM) / NaCl 0.4 M
- 200 µl SDS 10%
- 15 - 500 µl K-proteinase (2 mg K-proteinase in TE 10-2 / NaCl 0.4 M).

For the extraction of proteins, 1 ml saturated NaCl (6M) (1/3.5 v/v) was added. After vigorous agitation, the solution was centrifuged for 20 minutes at 10000 rpm.

For the precipitation of DNA, 2 to 3 volumes of 100% ethanol were added to the previous supernatant, and the solution was centrifuged for 30 minutes at 2000 rpm. The DNA solution was 20 rinsed three times with 70% ethanol to eliminate salts, and centrifuged for 20 minutes at 2000 rpm. The pellet was dried at 37°C, and resuspended in 1 ml TE 10-1 or 1 ml water. The DNA concentration was evaluated by measuring the OD at 260 nm (1 unit OD = 50 µg/ml DNA).

To determine the presence of proteins in the DNA solution, the OD 260 / OD 280 ratio was determined. Only DNA preparations having a OD 260 / OD 280 ratio between 1.8 and 2 were used in 25 the subsequent steps described below.

### DNA Amplification

Once each BAC was isolated, pairs of primers, each one defining a 500 bp-amplification fragment, were designed. Each of the primers contained a common oligonucleotide tail upstream of the specific bases targeted for amplification, allowing the amplification products from each set of 30 primers to be sequenced using the common sequence as a sequencing primer. The primers used for the genomic amplification of sequences derived from BACs were defined with the OSP software (Hillier L. and Green P. Methods Appl., 1991, 1: 124-8). The synthesis of primers was performed following the phosphoramidite method, on a GENSET UFPS 24.1 synthesizer.

Example 4 provides the procedures used in the amplification reactions.

The amplification of each sequence was performed by PCR (Polymerase Chain Reaction) as follows:

	- final volume	50 $\mu$ l
	- genomic DNA	100 ng
5	- MgCl <sub>2</sub>	2 mM
	- dNTP (each)	200 $\mu$ M
	- primer (each)	7.5 pmoles
	- Ampli Taq Gold DNA polymerase (Perkin)	1 unit
	- PCR buffer (10X=0.1 M Tris HCl pH 8.3, 0.5 M KCl) 1X.	

10 The amplification was performed on a Perkin Elmer 9600 Thermocycler or MJ Research PTC200 with heating lid. After heating at 94°C for 10 minutes, 35 cycles were performed. Each cycle comprised: 30 sec at 94°C, 1 minute at 55°C, and 30 sec at 72°C. For final elongation, 7 minutes at 72°C ended the amplification.

The obtained quantity of amplification products was determined on 96-well microtiter plates, using a fluorimeter and Picogreen as intercalating agent (Molecular Probes).

The sequences of the amplification products were determined for each of the approximately 100 individuals from whom genomic DNA was obtained. Those amplification products which contained biallelic markers were identified.

Figure 1 shows the locations of the biallelic markers along the 8p23 BAC contig. This first set of markers corresponds to a medium density map of the candidate locus, with an inter-marker distance averaging 50kb-150kb.

A second set of biallelic markers was then generated as described above in order to provide a very high-density map of the region identified using the first set of markers which can be used to conduct association studies, as explained below. The high density map has markers spaced on average every 2-50kb.

The biallelic markers were then used in association studies as described below.

#### Collection of DNA samples from affected and non-affected individuals

Prostate cancer patients were recruited according to clinical inclusion criteria based on pathological or radical prostatectomy records. Control cases included in this study were both ethnically- and age-matched to the affected cases; they were checked for both the absence of all clinical and biological criteria defining the presence or the risk of prostate cancer, and for the absence of related familial prostate cancer cases. Both affected and control individuals corresponded to unrelated cases.

The two following pools of independent individuals were used in the association studies. The first pool, comprising individuals suffering from prostate cancer, contained 185 individuals. Of these



185 cases of prostate cancer, 45 cases were sporadic and 140 cases were familial. The second pool, the control pool, contained 104 non-diseased individuals.

Haplotype analysis was conducted using additional diseased (total samples: 281) and control samples (total samples: 130), from individuals recruited according to similar criteria.

5                                   Genotyping Affected and Control Individuals

The general strategy to perform the association studies was to individually scan the DNA samples from all individuals in each of the two populations described above in order to establish the allele frequencies of the above described biallelic markers in each of these populations.

10                   Allelic frequencies of the above-described biallelic markers in each population were determined by performing microsequencing reactions on amplified fragments obtained by genomic PCR performed on the DNA samples from each individual.

DNA samples and amplification products from genomic PCR were obtained in similar conditions as those described above for the generation of biallelic markers, and subjected to automated microsequencing reactions using fluorescent ddNTPs (specific fluorescence for each 15 ddNTP) and the appropriate oligonucleotide microsequencing primers which hybridized just upstream of the polymorphic base. Once specifically extended at the 3' end by a DNA polymerase using the complementary fluorescent dideoxynucleotide analog (thermal cycling), the primer was precipitated to remove the unincorporated fluorescent ddNTPs. The reaction products were analyzed by electrophoresis on ABI 377 sequencing machines.

20                   Example 5 describes one microsequencing procedure.

Example 5

5 µl of PCR products in a microtiter plate were added to 5 µl purification mix {2U SAP (Amersham); 2U Exonuclease I (Amersham); 1 µl SAP10X buffer : 400mM Tris-HCl pH8, 100 mM MgCl<sub>2</sub>; H<sub>2</sub>O final volume 5 µl}. The reaction mixture was incubated 30 minutes at 37°C, and 25 denatured 10 minutes at 94°C. After 10 sec centrifugation, the microsequencing reaction was performed on line with the whole purified reaction mixture (10 µl) in the microplate using 10 pmol microsequencing oligonucleotide (23mers, GENSET, crude synthesis, 5 OD), 0.5 U Thermosequenase (Amersham), 1.25 µl Thermosequenase 16X buffer (Amersham), both of the fluorescent ddNTPs (Perkin Elmer) corresponding to the polymorphism {0.025 µl ddTTP and ddCTP, 0.05 µl ddATP and 30 ddGTP}, H<sub>2</sub>O to a final volume of 20 µl. A PCR program on a GeneAmp 9600 thermocycler was carried out as follows: 4 minutes at 94°C; 5 sec at 55°C / 10 sec at 94°C for 20 cycles. The reaction product was incubated at 4°C until precipitation. The microtiter plate was centrifuged 10 sec at 1500 rpm. 19 µl MgCl<sub>2</sub> 2mM and 55 µl 100 % ethanol were added in each well. After 15 minute incubation at room temperature, the microtiter plate was centrifuged at 3300 rpm 15 minutes at 4°C. 35 Supernatants were discarded by inverting the microtitre plate on a box folded to proper size and by centrifugation at 300 rpm 2 minutes at 4°C afterwards. The microplate was then dried 5 minutes in a

vacuum drier. The pellets were resuspended in 2.5 µl formamide EDTA loading buffer (0.7µl of 9 µg/µl dextran blue in 25 mM EDTA and 1.8 µl formamide). A 10% polyacrylamide gel / 12 cm / 64 wells was pre-run for 5 minutes on a 377 ABI 377 sequencer. After 5 minutes denaturation at 100°C, 0.8 µl of each microsequencing reaction product was loaded in each well of the gel. After migration (2 h 30 for 2 microtiter plates of PCR products per gel), the fluorescent signals emitted by the incorporated ddNTPs were analyzed on the ABI 377 sequencer using the GENESCAN software (Perkin Elmer). Following gel analysis, data were automatically processed with a software that allowed the determination of the alleles of biallelic markers present in each amplified fragment.

#### **I.D. Initial Association Studies**

10 Association studies were run in two successive steps. In a first step, a rough localization of the candidate gene was achieved by determining the frequencies of the biallelic markers of Figure 1 in the affected and unaffected populations. The results of this rough localization are shown in Figure 2. This analysis indicated that a gene responsible for prostate cancer was located near the biallelic marker designated 4-67.

15 In a second phase of the analysis, the position of the gene responsible for prostate cancer was further refined using the very high density set of markers described above. The results of this localization are shown in Figure 3.

As shown in Figure 3, the second phase of the analysis confirmed that the gene responsible for prostate cancer was near the biallelic marker designated 4-67, most probably within a ca. 150kb region comprising the marker.

#### **Haplotype analysis**

The allelic frequencies of each of the alleles of biallelic markers 99-123, 4-26, 4-14, 4-77, 99-217, 4-67, 99-213, 99-221, and 99-135 (SEQ ID NOs: 21-38) were determined in the affected and unaffected populations. Table 1 lists the internal identification numbers of the markers used in the haplotype analysis (SEQ ID NOs: 21-38), the alleles of each marker, the most frequent allele in both unaffected individuals and individuals suffering from prostate cancer, the least frequent allele in both unaffected individuals and individuals suffering from prostate cancer, and the frequencies of these alleles in each population.

Among all the theoretical potential different haplotypes based on 2 to 9 markers, 11 haplotypes showing a strong association with prostate cancer were selected. The results of these haplotype analyses are shown in Figure 4.

Figures 2, 3, and 4 aggregate linkage analysis results with sequencing results which permitted the physical order and/or the distance between markers to be estimated.

The significance of the values obtained in Figure 4 are underscored by the following results of computer simulations. For the computer simulations, the data from the affected individuals and the unaffected controls were pooled and randomly allocated to two groups which contained the same

number of individuals as the affected and unaffected groups used to compile the data summarized in Figure 4. A haplotype analysis was run on these artificial groups for the six markers included in haplotype 5 of Figure 4. This experiment was reiterated 100 times and the results are shown in Figure 5. Among 100 iterations, only 5% of the obtained haplotypes are present with a p-value below  $1 \times 10^{-4}$  as compared to the p-value of  $9 \times 10^{-7}$  for haplotype 5 of Figure 4. Furthermore, for haplotype 5 of Figure 4, only 6% of the obtained haplotypes have a significance level below  $5 \times 10^{-3}$ , while none of them show a significance level below  $5 \times 10^{-5}$ .

Thus, using the data of Figure 4 and evaluating the associations for single maker alleles or for haplotypes will permit estimation of the risk a corresponding carrier has to develop prostate cancer. Significance thresholds of relative risks will be adapted to the reference sample population used.

The diagnostic techniques may employ a variety of methodologies to determine whether a test subject has a biallelic marker pattern associated with an increased risk of developing prostate cancer or suffers from prostate cancer resulting from a mutant PG1 allele. These include any method enabling the analysis of individual chromosomes for haplotyping, such as family studies, single sperm DNA analysis or somatic hybrids.

In each of these methods, a nucleic acid sample is obtained from the test subject and the biallelic marker pattern for one or more of the biallelic markers listed in Figures 4, 6A and 6B is determined. The biallelic markers listed in Figure 6A are those which were used in the haplotype analysis of Figure 4. The first column of Figure 6A lists the BAC clones in which the biallelic markers lie. The second column of Figure 6A lists the internal identification number of the marker. The third column of Figure 6A lists the sequence identification number for a first allele of the biallelic markers. The fourth column of Figure 6A lists the sequence identification number for a second allele of the biallelic markers. For example, the first allele of the biallelic marker 99-123 has the sequence of SEQ ID NO:21 and the second allele of the biallelic marker has the sequence of SEQ ID NO: 30.

The fifth column of Figure 6A lists the sequences of upstream primers which is used to generate amplification products containing the polymorphic bases of the biallelic markers. The sixth column of Figure 6A lists the sequence identification numbers for the upstream primers.

The seventh column of Figure 6A lists the sequences of downstream primers which is used to generate amplification products containing the polymorphic bases of the biallelic markers. The eighth column of Figure 6A lists the sequence identification numbers for the downstream primers.

The ninth column of Figure 6A lists the position of the polymorphic base in the amplification products generated using the upstream and downstream primers. The tenth column lists the identities of the polymorphic bases found at the polymorphic positions in the biallelic markers. The eleventh and twelfth columns list the locations of microsequencing primers in the biallelic markers which can be used to determine the identities of the polymorphic bases.

In addition to the biallelic markers of SEQ ID NOs: 21-38, other biallelic markers (designated 99-1482, 4-73, 4-65) have been identified which are closely linked to one or more of the biallelic markers of SEQ ID NOs: 21-38, SEQ ID NOs: 57-62, and the PG1 gene. These biallelic markers include the markers of SEQ ID NOs: 57-62, which are listed in Figure 6B. The columns in Figure 6B  
5 are identical to the corresponding columns in Figure 6A. SEQ ID NOs: 58, 59, 61, and 62 lie within the PG1 gene of SEQ ID NO:1 at the positions indicated in the accompanying Sequence Listing.

Genetic analysis of these additional biallelic markers is performed as follows. Nucleic acid samples are obtained from individuals suffering from prostate cancer and unaffected individuals. The frequencies at which each of the two alleles occur in the affected and unaffected populations is  
10 determined using the methodologies described above. Association values are calculated to determine the correlation between the presence of a particular allele or spectrum of alleles and prostate cancer. The markers of SEQ ID NOs: 21-38 may also be included in the analysis used to calculate the risk factors. The markers of SEQ ID NOs: 21-38 and SEQ ID NOs: 57-62 is used in diagnostic techniques, such as those described below, to determine whether an individual is at risk for  
15 developing prostate cancer or suffers from prostate cancer as a result of a mutation in the PG1 gene.

Example 6 describes methods for determining the biallelic marker pattern.

#### Example 6

A nucleic acid sample is obtained from an individual to be tested for susceptibility to prostate cancer or PG1 mediated prostate cancer. The nucleic acid sample is an RNA sample or a DNA  
20 sample.

A PCR amplification is conducted using primer pairs which generate amplification products containing the polymorphic nucleotides of one or more biallelic markers associated with prostate cancer-related forms of PG1, such as the biallelic markers of SEQ ID NOs: 21-38, SEQ ID NOs: 57-62, biallelic markers which are in linkage disequilibrium with the biallelic markers of SEQ ID NOs:  
25 21-38, SEQ ID NOs: 57-62, biallelic markers in linkage disequilibrium with the PG1 gene, or combinations thereof. In some embodiments, the PCR amplification is conducted using primer pairs which generate amplification products containing the polymorphic nucleotides of several biallelic markers. For example, in one embodiment, amplification products containing the polymorphic bases of several biallelic markers selected from the group consisting of SEQ ID NOs: 21-38, SEQ ID NOs:  
30 57-62, and biallelic markers which are in linkage disequilibrium with the biallelic markers of SEQ ID NOs: 21-38, SEQ ID NOs: 57-62 or with the PG1 gene is generated. In another embodiment, amplification products containing the polymorphic bases of two or more biallelic markers selected from the group consisting of SEQ ID NOs: 21-38, SEQ ID NOs: 57-62, and biallelic markers which are in linkage disequilibrium with the biallelic markers of SEQ ID NOs: 21-38, SEQ ID NOs: 57-62  
35 or with the PG1 gene is generated. In another embodiment, amplification products containing the polymorphic bases of five or more biallelic markers selected from the group consisting of SEQ ID

NOs: 21-38, SEQ ID NOs: 57-62, and biallelic markers which are in linkage disequilibrium with the biallelic markers of SEQ ID NOs: 21-38, SEQ ID NOs: 57-62 or with the PG1 gene is generated. In another embodiment, amplification products containing the polymorphic bases of more than five of the biallelic markers selected from the group consisting of SEQ ID NOs: 21-38, SEQ ID NOs: 57-62, and biallelic markers which are in linkage disequilibrium with the biallelic markers of SEQ ID NOs: 21-38, SEQ ID NOs: 57-62 or with the PG1 gene is generated.

For example, the primers used to generate the amplification products may comprise the primers listed in Figure 6A or 6B (SEQ ID NOs: 39-56 and SEQ ID NOs: 63-68). Figures 6A and Figure 6B provide exemplary primers which is used in the amplification reactions and the identities and locations of the polymorphic bases in the amplification products which are produced with the exemplary primers. The sequences of each of the alleles of the biallelic markers resulting from amplification using the primers in Figures 6A and 6B are listed in the accompanying Sequence Listing as SEQ ID NOs: 21-38 and 57-62.

The PCR primers is oligonucleotides of 10, 15, 20 or more bases in length which enable the amplification of the polymorphic site in the markers. In some embodiments, the amplification product produced using these primers is at least 100 bases in length (i.e. 50 nucleotides on each side of the polymorphic base). In other embodiments, the amplification product produced using these primers is at least 500 bases in length (i.e. 250 nucleotides on each side of the polymorphic base). In still further embodiments, the amplification product produced using these primers is at least 1000 bases in length (i.e. 500 nucleotides on each side of the polymorphic base).

It will be appreciated that the primers listed in Figure 6A and 6B are merely exemplary and that any other set of primers which produce amplification products containing the polymorphic nucleotides of one or more of the biallelic markers of SEQ ID NOs. 21-38 and SEQ ID NOs: 57-62 or biallelic markers in linkage disequilibrium with the sequences of SEQ ID NOs. 21-38 and SEQ ID NOs: 57-62 or with the PG1 gene, or a combination thereof is used in the diagnostic methods.

Following the PCR amplification, the identities of the polymorphic bases of one or more of the biallelic markers of SEQ ID NOs: 21-38 and SEQ ID NOs: 57-62, or biallelic markers in linkage disequilibrium with the sequences of SEQ ID NOs. 21-38 and SEQ ID NOs: 57-62 or with the PG1 gene, or a combination thereof, are determined. The identities of the polymorphic bases is determined using the microsequencing procedures described in Example 5 above and the microsequencing primers listed as features in the sequences of SEQ ID NOs: 21-38 and SEQ ID NOs: 57-62. It will be appreciated that the microsequencing primers listed as features in the sequences of SEQ ID NOs: 21-38 and SEQ ID NOs: 57-62 are merely exemplary and that any primer having a 3' end near the polymorphic nucleotide, and preferably immediately adjacent to the polymorphic nucleotide, is used. Alternatively, the microsequencing analysis is performed as described in Pastinen et al., Genome Research 7:606-614 (1997), which is described in more detail below.

Alternatively, the PCR product is completely sequenced to determine the identities of the polymorphic bases in the biallelic markers. In another method, the identities of the polymorphic bases in the biallelic markers is determined by hybridizing the amplification products to microarrays containing allele specific oligonucleotides specific for the polymorphic bases in the biallelic markers.

5 The use of microarrays comprising allele specific oligonucleotides is described in more detail below.

It will be appreciated that the identities of the polymorphic bases in the biallelic markers is determined using techniques other than those listed above, such as conventional dot blot analyses.

Nucleic acids used in the above diagnostic procedures may comprise at least 10 consecutive nucleotides in the biallelic markers of SEQ ID NOs: 21-38 and SEQ ID NOs: 57-62 or the sequences

10 complementary thereto. Alternatively, the nucleic acids used in the above diagnostic procedures may comprise at least 15 consecutive nucleotides in the biallelic markers of SEQ ID NOs: 21-38 and SEQ ID NOs: 57-62 or the sequences complementary thereto. In some embodiments, the nucleic acids used in the above diagnostic procedures may comprise at least 20 consecutive nucleotides in the biallelic markers of SEQ ID NOs: 21-38 and SEQ ID NOs: 57-62 or the sequences complementary thereto. In

15 still other embodiments, the nucleic acids used in the above diagnostic procedures may comprise at least 30 consecutive nucleotides in the biallelic markers of SEQ ID NOs: 21-38 and SEQ ID NOs: 57-62 or the sequences complementary thereto. In further embodiments, the nucleic acids used in the above diagnostic procedures may comprise more than 30 consecutive nucleotides in the biallelic markers of SEQ ID NOs: 21-38 and SEQ ID NOs: 57-62 or the sequences complementary thereto. In

20 still further embodiments, the nucleic acids used in the above diagnostic procedures may comprise the entire sequence of the biallelic markers of SEQ ID NOs: 21-38 and SEQ ID NOs: 57-62 or the sequences complementary thereto.

#### **I.E. Identification and Sequencing, of the PG1 Gene, and Localization of the PG1 Protein**

The above haplotype analysis indicated that 171kb of genomic DNA between biallelic

25 markers 4-14 and 99-221 totally or partially contains a gene responsible for prostate cancer. Therefore, the protein coding sequences lying within this region were characterized to locate the gene associated with prostate cancer. This analysis, described in further detail below, revealed a single protein coding sequence in the 171 kb, which was designated as the PG1 gene.

Template DNA for sequencing the PG1 gene was obtained as follows. BACs 189EO8 and

30 463FO1 were subcloned as previously described. Plasmid inserts were first amplified by PCR on PE 9600 thermocyclers (Perkin-Elmer), using appropriate primers, AmpliTaqGold (Perkin-Elmer), dNTPs (Boehringer), buffer and cycling conditions as recommended by the Perkin-Elmer Corporation.

PCR products were then sequenced using automatic ABI Prism 377 sequencers (Perkin Elmer, Applied Biosystems Division, Foster City, CA). Sequencing reactions were performed using PE 9600

35 thermocyclers (Perkin Elmer) with standard dye-primer chemistry and ThermoSequenase (Amersham Life Science). The primers were labeled with the JOE, FAM, ROX and TAMRA dyes. The dNTPs and

ddNTPs used in the sequencing reactions were purchased from Boehringer. Sequencing buffer, reagent concentrations and cycling conditions were as recommended by Amersham.

Following the sequencing reaction, the samples were precipitated with EtOH, resuspended in formamide loading buffer, and loaded on a standard 4% acrylamide gel. Electrophoresis was performed  
5 for 2.5 hours at 3000V on an ABI 377 sequencer, and the sequence data were collected and analyzed using the ABI Prism DNA Sequencing Analysis Software, version 2.1.2.

The sequence data obtained as described above were transferred to a proprietary database, where quality control and validation steps were performed. A proprietary base-caller ("Trace"), working using a Unix system automatically flagged suspect peaks, taking into account the shape of the peaks, the inter-  
10 peak resolution, and the noise level. The proprietary base-caller also performed an automatic trimming. Any stretch of 25 or fewer bases having more than 4 suspect peaks was considered unreliable and was discarded. Sequences corresponding to cloning vector oligonucleotides were automatically removed from the sequence. However, the resulting sequence may contain 1 to 5 bases belonging to the vector sequences at their 5' end. If needed, these can easily be removed on a case by case basis.

15 The genomic sequence of the PG1 gene is provided in the accompanying Sequence Listing and is designated as SEQ ID NO: 1.

Potential exons in BAC-derived human genomic sequences were located by homology searches on protein, nucleic acid and EST (Expressed Sequence Tags) public databases. Main public databases were locally reconstructed. The protein database, NRPU (Non-redundant Protein Unique) is formed by a  
20 non-redundant fusion of the Genpept (Benson D.A. et al., Nucleic Acids Res. 24: 1-5 (1996), Swissprot (Bairoch, A. and Apweiler, R, Nucleic Acids Res. 24: 21-25 (1996) and PIR/NBRF (George, D.G. et al., Nucleic Acids Res. 24:17-20 (1996) databases. Redundant data were eliminated by using the NRDB software (Benson et al., supra) and internal repeats were masked with the XNU software (Benson et al., supra). Homologies found using the NRPU database allowed the identification of sequences  
25 corresponding to potential coding exons related to known proteins.

The EST local database is composed by the gbest section (1-9) of GenBank (Benson et al., supra), and thus contains all publicly available transcript fragments. Homologies found with this database allowed the localization of potentially transcribed regions.

The local nucleic acid database contained all sections of GenBank and EMBL (Rodriguez-  
30 Tome, P. et al., Nucleic Acids Res. 24: 6-12 (1996) except the EST sections. Redundant data were eliminated as previously described.

Similarity searches in protein or nucleic acid databases were performed using the BLAS software (Altschul, S.F. et al., J. Mol. Biol. 215: 403-410 (1990). Alignments were refined using the Fasta software, and multiple alignments used Clustal W. Homology thresholds were adjusted for each  
35 analysis based on the length and the complexity of the tested region, as well as on the size of the reference database.

Potential exon sequences identified as above were used as probes to screen cDNA libraries. Extremities of positive clones were sequenced and the sequence stretches were positioned on the genomic sequence of SEQ ID NO:1. Primers were then designed using the results from these alignments in order to enable the PG1 cloning procedure described below.

5

#### Cloning PG1 cDNA

PG1 cDNA was obtained as follows. 4:1 of ethanol suspension containing 1mg of human prostate total RNA (Clontech laboratories, Inc., Palo Alto, USA; catalogue N. 64038-1, lot 7040869) was centrifuged, and the resulting pellet was air dried for 30 minutes at room temperature.

First strand cDNA synthesis was performed using the Advantage<sup>TM</sup> RT-for-PCR kit (Clontech laboratories, Inc., Palo Alto, USA; catalogue N. K1402-1). 1:1 of 20mM solution of primer PGRT32: TTTTTTTTTTTTTTTTTTTTGAAAT (SEQ ID NO:10) was added to 12.5 :l of RNA solution in water, heated at 74°C for two and a half minutes and rapidly quenched in an ice bath. 10:1 of 5xRT buffer (50mM Tris-HCl, pH 8.3, 75mM KCl, 3 mM MgCl<sub>2</sub>), 2.5 :l of dNTP mix (10mM each), 1.25:1 of human recombinant placental RNA inhibitor were mixed with 1 ml of MMLV reverse transcriptase (200 units). 6.5:1 of this solution were added to RNA-primer mix and incubated at 42°C for one hour. 80:1 of water were added and the solution was incubated at 94°C for 5 minutes.

5:1 of the resulting solution were used in a Long Range PCR reaction with hot start, in 50 :l final volume, using 2 units of rTHXL, 20 pmol/μl of each of GC1.5p.1: CTGTCCCTGGTGCTCCACACGTA CTC (SEQ ID NO:6) or GC1.5p2 TGGTGCTCCACACGTA CTC CATGCGC (SEQ ID NO: 7) and GC1.3p: CTTGCCTGCTGGAGACACAGAATTTTCGATAGCAC (SEQ ID NO:9) primers with 35 cycles of elongation for 6 minutes at 67°C in thermocycler.

The sequence of the PG1 cDNA obtained as described above (SEQ ID NO 3) is provided in the accompanying Sequence Listing. Results of Northern blot analysis of prostate mRNAs support the existence of a major PG1 cDNA having a 5-6kb length.

#### Characterization of the PG1 Gene

The intron/exon structure of the gene was deduced by aligning the mRNA sequence from the cDNA of SEQ ID NO:3 and the genomic DNA sequence of SEQ ID NO: 1.

The positions of the introns and exons in the PG1 genomic DNA are provided in Figures 7 and 8. Figure 7 lists positions of the start and end nucleotides defining each of the at least 8 exons (labeled Exons A-H) in the sequence of SEQ ID NO: 1, the locations and phases of the 5' and 3' splice sites in the sequence of SEQ ID NO: 1, the position of the stop codon in the sequence of SEQ ID NO: 1, and the position of the polyadenylation site in the sequence of SEQ ID NO: 1. Figure 8 shows the positions of the exons within the PG1 genomic DNA and the PG1 mRNA, the location of a tyrosine phosphatase retro-pseudogene in the PG1 genomic DNA, the positions of the coding region in the mRNA, and the locations of the polyadenylation signal and polyA stretch in the mRNA.



As indicated in Figures 7 and 8, the PG1 gene comprises at least 8 exons, and spans more than 52kb. The first intron contains a tyrosine phosphatase retropseudogene. A G/C rich putative promoter region lies between nucleotide 1629 and 1870 of SEQ ID NO: 1. A CCAAT box is present at nucleotide 1661 of SEQ ID NO: 1. The promoter region was identified as described in Prestridge,  
5 D.S., Predicting Pol II Promoter Sequences Using Transcription Factor Binding Sites, J. Mol. Biol. 249:923-932 (1995).

It is possible that the methionine listed as being the initiating methionine in the PG1 protein sequence of SEQ ID NO: 4 (based on the cDNA sequence of SEQ ID NO: 3) may actually be downstream but in phase with another methionine which acts as the initiating methionine. The  
10 genomic DNA sequence of SEQ ID NO: 1 contains a methionine upstream from the methionine at position number 1 of the protein sequence of SEQ ID NO: 4. If the upstream methionine is in fact the authentic initiation site, the sequence of the PG1 protein would be that of SEQ ID NO: 5. This possibility is investigated by determining the exact position of the 5N end of the PG1 mRNA as follows.

15 One way to determine the exact position of the 5N end of the PG1 mRNA is to perform a 5NRACE reaction using the Marathon-Ready human prostate cDNA kit from Clontech (Catalog. No. PT1156-1). For example, the RACE reaction may employ the PG1 primers PG15RACE196 CAATATCTGGACCCCGGTGTAATTCTC (SEQ ID NO: 8) as the first primer. The second primer in the RACE reaction is PG15RACE130n having the sequence  
20 GGTCGTCCAGCGCTTGGTAGAAG (SEQ ID NO: 2). The sequence analysis of the resulting PCR product, or the product obtained with other PG1 specific primers, will give the exact sequence of the initiation point of the PG1 transcript.

Alternatively, the 5Nsequence of the PG1 transcript can be determined by conducting a PCR amplification with a series of primers extending from the 5Nend of the presently identified coding  
25 region. In any event, the present invention contemplates use of PG1 nucleic acids and/or polypeptides coding for or corresponding to either SEQ ID NO:4 or SEQ ID NO:5 or fragments thereof.

It is also possible that alternative splicing of the PG1 gene may result in additional translation products not described above. It is also possible that there are sequences upstream or downstream of the genomic sequence of SEQ ID NO: 1 which contribute to the translation products of the gene.  
30 Finally, alternative promoters may result in PG1derived transcripts other than those described herein.

The promoter activity of the region between nucleotides 1629 and 1870 can be verified as described below. Alternatively, should this region lack promoter activity, the promoter responsible for driving expression of the PG1 gene is identified as described below.

Genomic sequences lying upstream of the PG1 gene are cloned into a suitable promoter reporter  
35 vector, such as the pSEAP-Basic, pSEAP-Enhancer, pβgal-Basic, pβgal-Enhancer, or pEGFP-1 Promoter Reporter vectors available from Clontech. Briefly, each of these promoter reporter vectors

include multiple cloning sites positioned upstream of a reporter gene encoding a readily assayable protein such as secreted alkaline phosphatase,  $\beta$  galactosidase, or green fluorescent protein. The sequences upstream of the PG1 coding region are inserted into the cloning sites upstream of the reporter gene in both orientations and introduced into an appropriate host cell. The level of reporter protein is  
5 assayed and compared to the level obtained from a vector which lacks an insert in the cloning site. The presence of an elevated expression level in the vector containing the insert with respect to the control vector indicates the presence of a promoter in the insert. If necessary, the upstream sequences can be cloned into vectors which contain an enhancer for augmenting transcription levels from weak promoter sequences. A significant level of expression above that observed with the vector lacking an insert  
10 indicates that a promoter sequence is present in the inserted upstream sequence.

Promoter sequences within the upstream genomic DNA is further defined by constructing nested deletions in the upstream DNA using conventional techniques such as Exonuclease III digestion. The resulting deletion fragments can be inserted into the promoter reporter vector to determine whether the deletion has reduced or obliterated promoter activity. In this way, the boundaries of the promoters is  
15 defined. If desired, potential individual regulatory sites within the promoter is identified using site directed mutagenesis or linker scanning to obliterate potential transcription factor binding sites within the promoter individually or in combination. The effects of these mutations on transcription levels is determined by inserting the mutations into the cloning sites in the promoter reporter vectors.

Sequences within the PG1 promoter region which are likely to bind transcription factors is  
20 identified by homology to known transcription factor binding sites or through conventional mutagenesis or deletion analyses of reporter plasmids containing the promoter sequence. For example, deletions is made in a reporter plasmid containing the promoter sequence of interest operably linked to an assayable reporter gene. The reporter plasmids carrying various deletions within the promoter region are transfected into an appropriate host cell and the effects of the deletions on expression levels is assessed.  
25 Transcription factor binding sites within the regions in which deletions reduce expression levels is further localized using site directed mutagenesis, linker scanning analysis, or other techniques familiar to those skilled in the art.

The promoters and other regulatory sequences located upstream of the PG1 gene is used to design expression vectors capable of directing the expression of an inserted gene in a desired spatial,  
30 temporal, developmental, or quantitative manner. For example, since the PG1 promoter is presumably active in the prostate, it can be used to construct expression vectors for directing gene expression in the prostate.

Preferably, in such expression vectors, the PG1 promoter is placed near multiple restriction sites to facilitate the cloning of an insert encoding a protein for which expression is desired downstream of the  
35 promoter, such that the promoter is able to drive expression of the inserted gene. The promoter is inserted in conventional nucleic acid backbones designed for extrachromosomal replication, integration

into the host chromosomes or transient expression. Suitable backbones for the present expression vectors include retroviral backbones, backbones from eukaryotic episomes such as SV40 or Bovine Papilloma Virus, backbones from bacterial episomes, or artificial chromosomes.

Preferably, the expression vectors also include a polyA signal downstream of the multiple  
5 restriction sites for directing the polyadenylation of mRNA transcribed from the gene inserted into the expression vector.

Nucleic acids encoding proteins which interact with sequences in the PG1 promoter is identified using one-hybrid systems such as those described in the manual accompanying the Matchmaker One-Hybrid System kit available from Clontech (Catalog No. K1603-1). Briefly, the Matchmaker One-  
10 hybrid system is used as follows. The target sequence for which it is desired to identify binding proteins is cloned upstream of a selectable reporter gene and integrated into the yeast genome. Preferably, multiple copies of the target sequences are inserted into the reporter plasmid in tandem.

A library comprised of fusions between cDNAs to be evaluated for the ability to bind to the promoter and the activation domain of a yeast transcription factor, such as GAL4, is transformed into the  
15 yeast strain containing the integrated reporter sequence. The yeast are plated on selective media to select cells expressing the selectable marker linked to the promoter sequence. The colonies which grow on the selective media contain genes encoding proteins which bind the target sequence. The inserts in the genes encoding the fusion proteins are further characterized by sequencing. In addition, the inserts is inserted into expression vectors or in vitro transcription vectors. Binding of the polypeptides encoded by the  
20 inserts to the promoter DNA is confirmed by techniques familiar to those skilled in the art, such as gel shift analysis or DNase protection analysis.

#### Analysis of PG1 Protein Sequence

The PG1 cDNA of SEQ ID NO: 3 encodes a 353 amino-acid protein (SEQ ID NO:4). As indicated in the accompanying Sequence Listing, a Prosite analysis indicated that the PG1 protein has  
25 a leucine zipper motif, a potential glycosylation site, 3 potential casein kinase II phosphorylation sites, a potential cAMP dependent protein kinase phosphorylation site, 2 potential tyrosine kinase phosphorylation sites, 4 potential protein kinase C phosphorylation sites, 5 potential N-myristoylation sites, 1 potential tyrosine sulfation site, and one potential amidation site.

A search for membrane associated domains was conducted according to the methods  
30 described in Argos, P. et al., Structural Prediction of Membrane-bound Proteins, *Elur. J. Biochem.* 128:565-575 (1982); Klein et al., *Biochimica & Biophysica Acta* 815:468-476 (1985); and Eisenberg et al., *J. Mol. Biol.* 179:125-142 (1984). The search revealed 5 potential transmembrane domains predicted to be integral membrane domains. These results suggest that the PG1 protein is likely to be membrane-associated and is an integral membrane protein.

35 A homology search was conducted to identify proteins homologous to the PG1 protein. Several proteins were identified which share homology with the PG1 protein. Figure 9 lists the

accession numbers of several proteins which share homology with the PG1 protein in three regions designated box1, box2 and box3.

It will be appreciated that each of the motifs described above is also present in the protein of SEQ ID NO: 5, which would be produced if by translation initiation translated from the potential  
5 upstream methionine in the nucleic acid of SEQ ID NO: 1.

As indicated in Figure 9, a distinctive pattern of homology to box 1, box 2 (SEQ ID NOs: 11-14) and box 3 (SEQ ID NOs: 15-20) is found amongst acyl glycerol transferases. For example, the plsC protein from E. coli ( Accession Number P26647) shares homology with the box1 and box2 sequences, but not the box 3 sequence, of the PG1 protein. The product of this gene transfers acyl  
10 from acyl-coenzymeA to the sn2 position of 1-Acyl-sn-glycerol-3-phosphate (lysophosphatidic acid, LPA)(Coleman J., Mol Gen Genet. 1992 Mar 1; 232(2): 295-303).

Box1 and box2 homologies, but not box 3 homologies, are also found in the SLCI gene product from baker's yeast (Accession Number P33333) and the mouse gene AB005623. Each of these genes are able to complement in vivo mutations in the bacterial plsC gene. (Nagiec MM, Wells  
15 GB, Lester RL, Dickson RC, J. Biol. Chem., 1993 Oct 15; 268(29): 22156-22163, A suppressor gene that enables Saccharomyces cerevisiae to grow without making sphingolipids encodes a protein that resembles an Escherichia coli fatty acyltransferase; and Kume K, Shimizu T, Biochem. Biophys. Res. Commun. 1997, Aug. 28; 237(3): 663-666, cDNA cloning and expression of murine 1-acyl-sn-glycerol-3-phosphate acyltransferase).

20 Recently two different human homologues of the mouse AB005623 gene, Accession Numbers U89336 and U56417 were cloned and found to be localized to human chromosomes 6 and 9 (Eberhardt. C., Gray, P.W. and Tjoelker, L.W., J. Biol. Chem. 1997; 272, 20299-20305, Human lysophosphatidic acid acyltransferase cDNA cloning, expression, and localization to chromosome 9q34.3; and West, J., Tompkins, C.K., Balantac, N., Nudelman, E., Meengs, B., White, T., Bursten,  
25 S., Coleman, J., Kumar, A., Singer, J.W. and Leung, D.W, DNA Cell Biol. 6, 691-701 (1997), Cloning and expression of two human lysophosphatidic acid acyltransferase cDNAs that enhance cytokine induced signaling responses in cells).

The enzymatic acylation of LPA results in 1,2-diacyl-sn-glycerol 3-phosphate, an intermediate to the biosynthesis of both glycerophospholipids and triacylglycerol. Several important signaling  
30 messengers participating in the transduction of mitogenic signals, induction of apoptosis, transmission of nerve impulses and other cellular responses mediated by membrane bound receptors belong to this metabolic pathway.

LPA itself is a potent regulator of mammalian cell proliferation. In fact, LPA is one of the major mitogens found in blood serum. (For a review: Durieux ME, Lynch KR, Trends Pharmacol.  
35 Sci. 1993 Jun; 14(6):249-254, Signaling properties of lysophosphatidic acid. LPA can act as a survival factor to inhibit apoptosis of primary cells; and Levine JS, Koh JS, Triaca V, Lieberthal W,

Am. J. Physiol. 1997 Oct; 273(4Pt2): F575-F585, Lysophosphatidic acid: a novel growth and survival factor for renal proximal tubular cells). This function of LPA is mediated by the lipid kinase phosphatidylinositol 3-kinase.

Phosphatidylinositol and its derivatives present another class of messengers emerging from the 1-acyl-sn-glycerol-3-phosphate acyltransferase pathway. (Toker A, Cantley LC, Nature 1997 Jun 12; 387(6634): 673-676, Signaling through the lipid products of phosphoinositide-3-OH kinase; Martin TF, Curr. Opin. Neurobiol. 1997 Jun; 7(3):331-338, Phosphoinositides as spatial regulators of membrane traffic; and Hsuan JJ, et al., Int. J. Biochem. Cell Biol. 1997 Mar 1<sup>st</sup>; 29(3): 415-435, Growth factor-dependent phosphoinositide signaling).

Cell growth, differentiation and apoptosis can be affected and modified by enzymes involved in this metabolic pathway. Consequently, alteration of this pathway could facilitate cancer cell progression. Modulation of the activity of enzymes in this pathway using agents such as enzymatic inhibitors could be a way to restore a normal phenotype to cancerous cells.

Ashagbley A, Samadder P, Bittman R, Erukulla RK, Byun HS, Arthur G have recently shown that ether-linked analogue of lysophosphatidic acid: 4-O-hexadecyl-3(S)-O-methoxybutanephosphonate can effectively inhibit the proliferation of several human cancerous cell lines, including DU145 line of prostate cancer origin. (Anticancer Res 1996 Jul; 16(4A): 1813-1818, Synthesis of ether-linked analogues of lysophosphatidate and their effect on the proliferation of human epithelial cancer cells in vitro).

Structural differences between the PG1 family of cellular proteins and the functionally confirmed 1-acyl-sn-glycerol-3-phosphate acyltransferase family, evidenced by the existence of a different pattern of homology to box3, could point to unique substrate specificity in the phospholipid metabolic pathway, to specific interaction with other cellular components or to both.

Further analysis of the function of the PG1 gene can be conducted, for example, by constructing knockout mutations in the yeast homologues of the PG1 gene in order to elucidate the potential function of this protein family, and to test potential substrate analogs in order to revert the malignant phenotype of human prostate cancer cells as described in Section VIII, below.

#### Example 7

##### Analysis of the Intracellular Localisation of the PG1 Isoforms

To study the intracellular localisation of PG1 protein, different isoforms of PG1 were cloned in the expression vector pEGFP-N1(Clontech), transfected and expressed in normal (PNT2A) or adenocarcinoma (PC3) prostatic cell line.

First, to generate cDNA inserts, 5' and 3' primers were synthesised allowing to amplify different regions of the PG1 open reading frame. Respectively, these primers were designed with an internal EcoRI or BamHI site which allowed the insertion of the amplified product into the EcoRI and BamHI sites of the expression vector. The restriction sites were introduced into the primer so that

after cloning into pEGFP-N1, the PG1 open reading frame would be fused in frame, to the EGFP open reading frame. The translated protein would be a fusion between PG1 and EGFP. EGFP being a variant form of the GFP protein (Green Fluorescent Protein), it is possible to detect the intracellular localisation of the different PG1 isoforms by examining the fluorescence emitted by the EGFP fused protein.

The different forms that were analysed correspond either to different messengers identified by RT-PCR performed on total normal human prostatic RNA or to a truncated form resulting from a non sense mutation identified in a tumoural prostatic cell line LnCaP. The different PG1 constructions were transfected using the lipofectine technique and EGFP expression was examined 20 hours post transfection.

Name and description of the different forms transfected are listed below :

- A) PG1 includes all the coding exons from exon 1 to 8.
- B) PG1/1-4 corresponds to an alternative messenger which is due to an alternative splicing, joining exon 1 to exon 4, and resulting in the absence of exons 2 and 3.
- 15 C) PG1/1-5 corresponds to an alternative messenger which is due to an alternative splicing, joining exon 1 to exon 5, and resulting in the absence of exons 2, 3 and 4.
- D) PG1/1-7 includes exons 1 to 6, and corresponds to the mutated form identified in genomic DNA of the prostatic tumoural cell line LNCaP.

Cloning of the PG1 cDNA inserts in the EGFP-N1 expression vector

20 cDNAs from human prostate were obtained by RT-PCR using the Advantage RT-for-PCR Kit (CLONTECH ref K1402-2). First, 1µl of oligodT-containing PG1 specific primer PGRT32 TTTTTTTTTTTTTTTTTTGGAAAT (20pmoles) and 11.5 µl of DEPC treated H<sub>2</sub>O were added to 1µl of total mRNA (1µg) extracted from human prostate (CLONTECH ref 64038-1). The mRNA was heat denaturated for 2.5 min at 74°C and then quickly chilled on ice. A mix containing 4µl of 5X buffer, 1µl of dNTPs (10mM each), 0.5µl of recombinant RNase inhibitor (20U) and 1µl of MoMuLV Reverse Transcriptase (200U) was added to the denaturated mRNA. Reverse transcription was performed for 60 min. at 42°C. Enzymes were heat denaturated for 5 min. at 94°C. Then, 80µl of DEPC treated H<sub>2</sub>O were added to the reaction mix and the cDNA mix was stored at -20°C. Primers PG15Eco3 (5' CCTGAATTCCGCCGAGCTGAGAAGATGC 3'), and PG13Bam2 (5' 25 CCTGGATCCGCTTTAATAGTAACCCACAGGCAG 3') were used for PCR amplification of the different PG1 cDNAs. A 50µl PCR reaction mix containing 5µl of the previously prepared prostate cDNA mix, 15µl of 3.3X PCR buffer, 4µl of dNTPs (2.5mM each), 20pmoles of primer PG15Eco3, 20pmoles of primer PG13Bam2, 1µl of RthXL enzyme, 2.2µl Mg(OAc)<sub>2</sub> (Hot Start) was set up and amplification was performed for 35 cycles of 30 sec at 94°C, 10 min. at 72°C, 4 min. at 67°C after an 35 initial denaturation step of 10 min. at 94°C. Size and integrity of the PCR product was assessed by migration on a 1% agarose gel. 2µg of the amplification product were digested with 2.4 units of

EcoRI (PROMEGA ref R601A) and 2.0 units of BamHI (PROMEGA ref R602A) in 50µl of 1X Multicore buffer for 2 hours at 37°C. Enzymes were then heat inactivated for 20 min, at 68°C, DNA was phenol/chloroform extracted and ethanol-precipitated and its concentration was estimated by migration on a 1% agarose gel.

5 To prepare the vector, 2µg of pEGFP-N1 vector (CLONTECH ref 6085-1) were digested with 2.4 units of EcoRI (PROMEGA ref R601A) and 2.0 units of BamHI (PROMEGA ref R602A) in 50µl of 1X multicore buffer for 2 hours at 37°C. Enzymes were then heat inactivated for 20 min, at 68°C, DNA was phenol/chloroform extracted and ethanol-precipitated and its concentration and integrity were estimated by migration on a 1% agarose gel. 20ng of the BamHI and EcoRI digested pEGFP-N1  
10 vector were added to 50ng of BamHI-EcoRI digested PG1 cDNAs. Ligation was performed over night at 13°C using 0.5units of T4 DNA ligase (BOEHRINGER ref 84333623) in a final volume of 20µl containing 1X ligase buffer. The ligation reaction mix was desalted by dialysis against water (MILLIPORE ref VSWP01300) for 30min. at room temperature. One fifth of the desalted ligation reaction was electroporated in 25µl of competent cells ElectroMAX DH10B (GIBCO BRL ref  
15 18290-015) using a resistance of 126 Ohms, capacitance of 50µF, and voltage of 2.5KV. Bacteria were then incubated in 500µl of SOB medium for 30min at 37°C. One fifth was plated on LB AGAR containing 40µg/µl KANAMYCINE (SIGMA ref K4000) and incubated over night at 37°C. Plasmid DNA was prepared from an overnight liquid culture of individual colonies and sequenced. Among the different forms identified 3 were used :

- 20 A) PG1 which includes all the coding exons from exon 1 to 8.  
B) PG1/1-4 which corresponds to an alternative messenger which is due to an alternative splicing, joining exon 1 to exon 4, and resulting in the absence of exons 2 and 3.  
C) PG1/1-5 which corresponds to an alternative messenger which is due to an alternative splicing, joining exon 1 to exon 5, and resulting in the absence of exons 2, 3 and 4.  
25 D) Vector PG1/1-7 : A cDNA insert encoding for a truncated protein was synthesized by PCR amplification, using primers PG15Eco3 and PG1mut29Bam (5' CCTGGATCCCCCTCCATCGTCTTTCCCTT 3') and vector PG1 as a template. The resulting PCR product was cloned following the same protocol as described above.

Transfection of the PG1 expression vectors in human prostate cell lines.

30 The DNA/lipofectin solution was prepared as followed: 1.5µl of lipofectin (GIBCO BRL ref 18292-011) was diluted in 100µl of OPTI-MEM medium (GIBCO BRL ref 31985-018), and incubated for 30min. at room temperature before being mixed to 0.5µg of vector diluted in 100µl of OPTI-MEM medium and incubated for 15 min. at room temperature. Cells were inoculated in RPMI1640 medium (Gibco BRL ref 61870-010) containing 5% fetal calf serum (Dutscher ref P30-  
35 3302) on slides (NUNC Lab-Tek ref 177402A) and grown at 37°C in 5%CO<sub>2</sub>. Cells reaching 40-60% confluency were rinsed with 300µl OPTI-MEM medium and incubated with the DNA/lipofectin

solution for 6 hours at 37°C. The medium containing DNA was replaced by medium supplemented in fetal calf serum and cells were incubated for at least 36 hours at 37°C. Slides were rinsed in PBS and cells were fixed in ethanol, treated with Propidium iodide, and examined with a fluorescence microscope using a double-pass filter set for FITC/PI.

5 After transfection of PG1 and PG1/1-4 in both the normal and tumoural prostatic cell line, green fluorescence was detected into and around the nucleus (Figures 10 and 11). This result shows that the PG1 protein is localised in the nucleus and/or the nuclear membrane. Furthermore, it suggests that exons 2 and 3 are dispensable for translocation of PG1 to the nucleus. In addition, no difference in the intracellular localisation of these two forms was detected between the tumoral and the normal  
10 prostatic cell line.

On the contrary, transfection experiments using PG1/1-5 show that this form is cytoplasmic in the normal prostatic cell line PNT2A. It suggests that exon 4 might be important for the regulation of the translocation to the nucleus. Interestingly, similar transfection experiments in the tumoral cell line PC3 show that PG1/1-5 remains nuclear and or perinuclear (Figure 12). This result shows that there is  
15 an abnormality in the regulation of the intracellular localization of the PG1 isoforms in this tumoral cell line. Furthermore, it indicates that the normal function of PG1 can be altered indirectly in prostatic tumors by an abnormality in the regulation of its intracellular location.

Finally, a non-sense mutation has been identified in the prostatic tumoural cell line LNCaP, in exon 6 of PG1 (SEQ ID NO: 69). This mutation is responsible for the production of a truncated  
20 protein (SEQ ID NO: 70). To determine the intracellular location of this truncated protein, PG1/1-7 and PG1 were transfected in the normal prostatic cell line PNT2A. Comparison of the fluorescence detected in both sets of experiments clearly showed that the truncated form was localised in the cytoplasm as the non-truncated protein was located in and/or around the nucleus (Figure 13). This result indicates that this mutated PG1 is translated in a truncated protein which is unable to reach the  
25 nucleus. It also suggests that exons 7 and 8 may play an important role in the regulation of the intracellular localisation of PG1. Furthermore, it supports the previous hypothesis that an altered regulation of PG1 intracellular localisation might be involved in prostate tumorigenesis.

	pEGFP N1	PG1	PG11-4	PG11-5	PG11-7
Transfection PNT2 06/17/98	NA	nuclear	nuclear	ND	ND

Transfection PNT2 06/30/98	cytoplasmic	nuclear	nuclear	ND	ND
----------------------------------	-------------	---------	---------	----	----



Transfection PNT2 07/16/98	cytoplasmic	NA	NA	cytoplasmic	ND
Transfection PC3 07/16/98	NA	nuclear	nuclear	nuclear	ND
Transfection PC3 07/16/98 bis	cytoplasmic	nuclear	NA	NA	ND

Transfection PC3 08/27/98	cytoplasmic	nuclear	nuclear	nuclear	NA
Transfection PNT2 08/28/98	cytoplasmic	nuclear	NA	cytoplasmic	cytoplasmic
		All exons	X2-3 Spliced out	X2-3-4Spliced out	mut aa229

NA : Not assessable

ND : Not done

- 5 Nuclear : localized in and around the nucleus (nuclear and perinuclear localization).

#### Alternative Splice Species

Alternative splicing is a common natural tool for the inhibition of function of full length gene products. Alternative splicing is known to result in enzyme isoforms, possessing different kinetic characteristics (pyruvate kinase: M1 and M2 Yamada K, Noguchi T, Biochem J. 1999 Jan1;337(Pt 1):1-11. Estrogen receptor (ER) gene is known to possess variant splicing yielding the deletions of exon 3, 5, or 7. The truncated ER protein induced from variant mRNA could mainly be exhibited as a repressor through dominant negative effects on normal ER protein (Iwase H, Omoto Y, Iwata H, Hara Y, Ando Y, Kobayashi S, Oncology 1998 Dec;55 Suppl S1:11-16 )'Yu et al ( Yu JJ, Mu C, Dabholkar M, Guo Y, Bostick-Bruton F, Reed E,Int J Mol Med 1998 Mar;1(3):617-620 ) demonstrated that there is an association between alternative splicing of ERCC1, and reduction in cellular capability to repair cisplatin-DNA adduct. Munoz-Sanjuan et al (Munoz-Sanjuan I, Simandl BK, Fallon JF, Nathans J, Development 1998 Dec 14;126(Pt 2):409-421) demonstrated existence of two differentially spliced isoforms of fibroblast growth factor(FGF) type two genes that are present in non-overlapping spatial

distributions in the neural tube and adjacent structures in developing chicken embryo. One of these forms is secreted and activates the expression of HoxD13, HoxD11, Fgf-4 and BMP-2 ectopically, consistent with cFHF-2 playing a role in anterior-posterior patterning of the limb.

The CD44 is a cell adhesion molecule that is present as numerous isoforms created by mRNA  
5 alternative splicing. Expression of variant isoforms of CD44 is associated with tumor growth and metastasis. (Shibuya Y, Okabayashi T, Oda K, Tanaka N, Jpn J Clin Oncol 1998 Oct;28(10):609-14) they showed that ratio of two particular isoforms is a useful indicator of prognosis in gastric and colorectal carcinoma. Zhang YF et al (Zhang YF, Jeffery S, Burchill SA, Berry PA, Kaski JC, Carter ND, Br J Cancer 1998 Nov;78(9):1141-6) showed that human endothelin receptor A is the subject to  
10 alternative splicing giving at least two isoforms. The truncated receptor was expressed in all tissues and cells examined, but the level of expression varied. In melanoma cell lines and melanoma tissues, the truncated receptor gene was the major species, whereas the wild-type ETA was predominant in other tissues. Zhang et al. conclude that the function and biological significance of this truncated ETA receptor is not clear, but it may have regulatory roles for cell responses to ETs.

15

#### Example 8

##### Identification of PG1 Alternative Splice Species

The PG1 cDNA was first cloned by screening of a human prostate cDNA library. Sequence analysis of about 400 cDNA clones showed that at least 14 isoforms were present in this cDNA library. Comparison of their sequences to the genomic sequence showed that these isoforms resulted  
20 from a complex set of different alternative splicing events between numerous exons (Figure 14).

To rule out the possibility of a cloning artefact generated during the cDNA library construction and to systematically identify all existing alternative splice junctions, RT-PCR experiments were performed on RNA of normal prostate as well as normal prostatic cell lines PNT1A, PNT1B and PNT2 using all the possible combinations of primers specific to the different  
25 exon borders. SEQ ID NOs: 137-178. The presence of multiple PCR bands in each reaction was assessed by migration in an agarose gel. Each band was analysed by sequencing, and the presence or absence of specific splicing events, as seen in the sequence by a specific splice junction, was scored as plus or minus in Figure 15.

Furthermore, to identify aberrant splicing event in prostate tumors, similar experiments were  
30 performed on RNA extracted from tumoral prostatic cell lines LnCaP (obtained from two different sources and named FCG and JMB), CaHPV, Du145 and PC3 as well as on RNA obtained from prostate tumors (ECP5 to ECP24).

As shown in the first five columns, all isoforms identified in the cDNA library were detected in RNA of normal prostate, normal prostatic cell lines or prostate tumors. In addition to the different  
35 splice junctions detected in the cDNA library, 19 other splice junctions were detected in normal prostate or in normal prostatic cell lines. Two types of exon junctions (exons 3-7, exons 3b-8) were

never detected in either normal prostate, normal prostatic cell lines, prostate tumors or prostatic tumoral cell lines. Comparison between normal and tumoral samples showed the presence of 2 additional exon junctions ( exons 3-8, exons 5-8) in the tumoral samples that were not detected previously in the normal samples. This result demonstrate that during tumorigenesis, the complex regulation of the PG1 splicing has been altered, resulting in an abnormal ratio of the different isoforms. It is of a specific interest since it has been shown in patients with a genetic predisposition to Wilms tumor, that an imbalance between different RNA isoforms might be involved in tumorigenesis (Bickmore et al., Science 1992, 257:325-7; Little et al, Hum Mol Genet 1995, 4:351-8).

Interestingly, comparison between normal and tumoral samples, also showed that some exon junctions are present in all normal samples, but are absent in numerous tumoral samples. It further indicates that the normal function of PG1 can be altered by an abnormality in the regulation of PG1 splicing and further support the previous hypothesis.

Furthermore, comparison between the different types of normal samples (Col.2 versus Col. 3, 4 and 5) also showed differences in the presence or absence of some exon junctions. It indicates that the transformation process necessary to the generation of these normal prostatic cell lines might result in similar alteration which further support the previous hypothesis.

#### Example 9

##### Determining the Tumor Suppressor Activity of the PG1 Gene Product, Mutants and Other PG1

##### Polypeptides

PG1 variants which results from either alternate splicing of the PG1 mRNA or from mutation of PG1 that introduce a stop codon (nucleotide of SEQ ID NO: 69 and protein of SEQ ID NO: 70) can no longer perform its role of tumor suppressor. It is possible and even likely that PG1 tumor suppressor role extends beyond prostate cancer to other form of malignancies. PG1 therefore represent a prime candidate for gene therapy of cancer by creating a targeting vector which knocks out the mutant and/or introduces a wild-type PG1 gene (e.g. SEQ ID NO 3 or 179) or a fragment thereof.

To validate this model, PG1 and its alternatively spliced or mutated variants are stably transfected in tumor cell line using methods described in Section VIII. The efficiency of transfection is determined by northern and western blotting; the latter is performed using antibodies prepared against PG1 synthetic peptides designed to distinguish the product of the most abundant PG1 mRNA from the alternatively spliced variants, the truncated variant, or other functional mutants. The production of synthetic peptides and of polyclonal antibodies is performed using the methods described herein in Sections III and VII.. After demonstrating that PG1 and its variant are efficiently expressed in various tumor cell line preferably derived from human prostate cancer, hepatocarcinoma, lung and colon carcinoma; we the effect of this gene on the rate of cell division, DNA synthesis, ability to grow in soft agar and ability to induce tumor progression and metastasis when injected in immunologically deficient nude mice are determined.

Alternatively the PG1 gene and its variant are inserted in adenoviruses that are used to obtain a high level of expression of these genes. This method is preferred to test the effect of PG1 expression in animal that are spontaneously developing tumor. The production of specific adenoviruses is obtained using methods familiar to those with normal skills in cell and molecular biology.

## **II. POLYNUCLEOTIDES:**

The present invention encompasses polynucleotides in the form of PG1 genomic or cDNA as well as polynucleotides for use as primers and probes in the methods of the invention. These polynucleotides may consist of, consist essentially of, or comprise a contiguous span of nucleotides of a sequence from any sequence in the Sequence Listing as well as sequences which are complementary thereto ("complements thereof"). Preferably said sequence is selected from SEQ ID NOs: 3, 112-125, 179, 182-184. The "contiguous span" is at least 6, 8, 10, 12, 15, 20, 25, 30, 50, 100, 200, or 500 nucleotides in length. It should be noted that the polynucleotides of the present invention are not limited to having the exact flanking sequences surrounding the polymorphic bases which are enumerated in Sequence Listing. Rather, it will be appreciated that the flanking sequences surrounding the biallelic markers, or any of the primers or probes of the invention which are more distant from a biallelic markers, is lengthened or shortened to any extent compatible with their intended use and the present invention specifically contemplates such sequences. It will be appreciated that the polynucleotides referred to in the Sequence Listing is of any length compatible with their intended use. Also the flanking regions outside of the contiguous span need not be homologous to native flanking sequences which actually occur in humans. The addition of any nucleotide sequence, which is compatible with the nucleotides intended use is specifically contemplated. The contiguous span may optionally include the PG1-related biallelic marker in said sequence. Optionally either allele of the biallelic markers described above in the definition of PG1-related biallelic marker is specified as being present at the PG1-related biallelic marker.

The invention also relates to polynucleotides that hybridize, under conditions of high or intermediate stringency, to a polynucleotide of a sequence from any sequence in the Sequence Listing as well as sequences, which are complementary thereto. Preferably said sequence is selected from SEQ ID NOs: 3, 112-125, 179, 182-184. Preferably such polynucleotides is at least 6, 8, 10, 12, 15, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 200, or 500 nucleotides in length. Preferred polynucleotides comprise an PG1-related biallelic marker. Optionally either allele of the biallelic markers described above in the definition of PG1-related biallelic marker is specified as being present at the biallelic marker site. Conditions of high and intermediate stringency are further described in Section X.C.4, below.

The invention embodies polynucleotides which encode an entire human, mouse or mammalian PG1 protein, or fragments thereof. Generally the polynucleotides of the invention

comprise the naturally occurring nucleotide sequence of the PG1. However, any naturally occurring silent codon variation or other silent codon variation can be employed to encode the PG1 amino acids sequence. As for those amino acids which are changed or added to the PG1 gene for any embodiment of the invention which requires the expression of a nucleotide sequence, the nucleic acid sequences

5 generally will be chosen to optimize expression in the specific human or non-human animal system in which the polynucleotide is intended to be used, making use of known codon preferences. The PG1 polynucleotides of the invention can be the native nucleotide sequence which encodes a human, mouse, or mammalian PG1 protein, preferably the PG1 polynucleotide sequence of SEQ ID NOs: 3, 112-125, 179, 182-184, and the compliments thereof. The polynucleotides of the invention include

10 those which encode PG1 polypeptides with a contiguous stretch of at least 8, 10, 12, 15, 20, 25, 30, 50, 100 or 200 amino acids from SEQ ID NOs: 4, 5, 70, 74, and 125-136, as well as any other human, mouse or mammalian PG1 polypeptide. In addition the present invention encompasses polynucleotides which comprise a contiguous stretch of at least 8, 10, 12, 15, 20, 25, 30, 50, 100, 200, 500 nucleotides of a human, mouse or mammalian PG1 genomic sequence as well as complete human,

15 mouse, or mammalian PG1 genes, preferably of SEQ ID NOs: 179, 182, 183, and the compliments thereof.

The present invention encompasses polynucleotides which consist of, consist essentially of, or comprise a contiguous stretch of at least 8, 10, 12, 15, 20, 25, 30, 50, 100, 200, or 500 nucleotides of a human, mouse or mammalian PG1 cDNA sequences as well as an entire human, mouse, or

20 mammalian PG1 cDNA. The cDNA species and polynucleotide fragments comprised by the polynucleotides of the invention include the predominant species derived from any human, mouse or mammal source, preferably SEQ ID NOs: 3, 184, and the compliments thereof. In addition, the polynucleotides of the invention comprise cDNA species, and fragments thereof, that result from the alternative splicing of PG1 transcripts in any human, mouse or other mammal, preferably the cDNA

25 species of SEQ ID NOs: 112-124, and compliments thereof. Moreover, the invention encompasses cDNA species and other polynucleotides which consist of or comprise the polynucleotides which span a splice junction, preferably including any one of SEQ ID NOs: 137 to 178, and the compliments thereof; more preferably any one of SEQ ID NOs: 137 to 149, 151 to 169, 171 to 178, and the compliments thereof. The polynucleotides of the invention also include cDNA and other

30 polynucleotides which comprise two covalently linked PG1 exons, derived from a single human, mouse or mammalian species, immediately adjacent to one another in the order shown, and selected from the following pairs of PG1 exons: 1:2, 1:3, 1:4, 1:5, 1:6, 1:7, 1:8, 2:3, 2:4, 2:5, 2:6, 2:7, 2:8, 3:4, 3:5, 3:6, 3:7, 3:8, 4:5, 4:6, 4:7, 4:8, 5:6, 5:7, 5:8, 6:7, 6:8, 7:8, 1:1bis, 1bis:2, 1bis:3, 1bis:4, 1bis:5, 1bis:6, 1bis:7, 1bis:8, 3:3bis, 3bis:4, 3bis:5, 3bis:6, 3bis:7, 3bis:8, 5:5bis, 5bis:6, 5bis:7, 5bis:8,

35 1:6bis, 2:6bis, 3:6bis, 4:6bis, 5:6bis, 6bis:7, 6bis:8, and the compliments thereof. In a preferred embodiment the sequences of the PG1 exons in each of the pairs of exons is selected as follows:

exon 1 – SEQ ID NO: 100; exon 2 – SEQ ID NO: 101; exon 3 – SEQ ID NO: 102;  
 exon 4 – SEQ ID NO: 103; exon 5 – SEQ ID NO: 104; exon 6 – SEQ ID NO: 105;  
 exon 7 – SEQ ID NO: 106; exon 8 – SEQ ID NO: 107; exon 1bis – SEQ ID NO: 108;  
 exon 3bis – SEQ ID NO: 109; exon 5bis – SEQ ID NO: 110; and

5 exon 6bis – SEQ ID NO: 111. Because of the 8 different polyadenylation sites in exon 8, any cDNA or polynucleotide of the invention comprising a human cDNA fragment encompassing exon 8 is truncated such that only the first 330 nucleotides, 699 nucleotides, 833 nucleotides, 1826 nucleotides, 2485 nucleotides, 2805 nucleotides, 4269 nucleotides or 4315 nucleotides of exon 8 shown in SEQ ID NO: 107 are present.

10 The primers of the present invention is designed from the disclosed sequences for any method known in the art. A preferred set of primers is fashioned such that the 3' end of the contiguous span of identity with the sequences of the Sequence Listing is present at the 3' end of the primer. Such a configuration allows the 3' end of the primer to hybridize to a selected nucleic acid sequence and dramatically increases the efficiency of the primer for amplification or sequencing reactions. Allele  
 15 specific primers is designed such that a biallelic marker is at the 3' end of the contiguous span and the contiguous span is present at the 3' end of the primer. Such allele specific primers tend to selectively prime an amplification or sequencing reaction so long as they are used with a nucleic acid sample that contains one of the two alleles present at a biallelic marker. The 3' end of primer of the invention is located within or at least 2, 4, 6, 8, 10, 12, 15, 18, 20, 25, 50, 100, 250, 500, or 1000 nucleotides  
 20 upstream of an PG1-related biallelic marker in said sequence or at any other location which is appropriate for their intended use in sequencing, amplification or the location of novel sequences or markers.

Preferred amplification primers include the polynucleotides disclosed in SEQ ID NOs: 39-56, and 63-68. Additional preferred amplification primers for particular non-genic PG1-related biallelic  
 25 markers are listed as follows by the internal reference number for the marker and the SEQ ID NOs for the PU and RP amplification primers respectively:

4-14-107 use SEQ ID NOs 339 and 382; 4-14-317 use SEQ ID NOs 339 and 382;  
 4-14-35 use SEQ ID NOs 339 and 382; 4-20-149 use SEQ ID NOs 340 and 383;  
 4-22-174 use SEQ ID NOs 341 and 384; 4-22-176 use SEQ ID NOs 341 and 384;  
 30 4-26-60 use SEQ ID NOs 342 and 385; 4-26-72 use SEQ ID NOs 342 and 385;  
 4-3-130 use SEQ ID NOs 343 and 386; 4-38-63 use SEQ ID NOs 344 and 387;  
 4-38-83 use SEQ ID NOs 344 and 387; 4-4-152 use SEQ ID NOs 345 and 388; 4-4-187 use  
 SEQ ID NOs 345 and 388; 4-4-288 use SEQ ID NOs 345 and 388;  
 4-42-304 use SEQ ID NOs 346 and 389; 4-42-401 use SEQ ID NOs 346 and 389;  
 35 4-43-328 use SEQ ID NOs 347 and 390; 4-43-70 use SEQ ID NOs 347 and 390;  
 4-50-209 use SEQ ID NOs 348 and 391; 4-50-293 use SEQ ID NOs 348 and 391;

- 4-50-323 use SEQ ID NOs 348 and 391; 4-50-329 use SEQ ID NOs 348 and 391;  
 4-50-330 use SEQ ID NOs 348 and 391; 4-52-163 use SEQ ID NOs 349 and 392;  
 4-52-88 use SEQ ID NOs 349 and 392; 4-53-258 use SEQ ID NOs 350 and 393;  
 4-54-283 use SEQ ID NOs 351 and 394; 4-54-388 use SEQ ID NOs 351 and 394;  
 5 4-55-70 use SEQ ID NOs 352 and 395; 4-55-95 use SEQ ID NOs 352 and 395;  
 4-56-159 use SEQ ID NOs 353 and 396; 4-56-213 use SEQ ID NOs 353 and 396;  
 4-58-289 use SEQ ID NOs 354 and 397; 4-58-318 use SEQ ID NOs 354 and 397;  
 4-60-266 use SEQ ID NOs 355 and 398; 4-60-293 use SEQ ID NOs 355 and 398;  
 4-84-241 use SEQ ID NOs 356 and 399; 4-84-262 use SEQ ID NOs 356 and 399;  
 10 4-86-206 use SEQ ID NOs 357 and 400; 4-86-309 use SEQ ID NOs 357 and 400;  
 4-88-349 use SEQ ID NOs 358 and 401; 4-89-87 use SEQ ID NOs 359 and 402;  
 99-123-184 use SEQ ID NOs 360 and 403; 99-128-202 use SEQ ID NOs 361 and 404; 99-128-  
 275 use SEQ ID NOs 361 and 404; 99-128-313 use SEQ ID NOs 361 and 404;  
 99-128-60 use SEQ ID NOs 361 and 404; 99-12907-295 use SEQ ID NOs 362 and 405; 99-  
 15 130-58 use SEQ ID NOs 363 and 406; 99-134-362 use SEQ ID NOs 364 and 407; 99-140-130  
 use SEQ ID NOs 365 and 408; 99-1462-238 use SEQ ID NOs 366 and 409; 99-147-181 use  
 SEQ ID NOs 367 and 410; 99-1474-156 use SEQ ID NOs 368 and 411; 99-1474-359 use SEQ  
 ID NOs 368 and 411;  
 99-1479-158 use SEQ ID NOs 369 and 412;  
 20 99-1479-379 use SEQ ID NOs 369 and 412; 99-148-129 use SEQ ID NOs 370 and 413;  
 99-148-132 use SEQ ID NOs 370 and 413; 99-148-139 use SEQ ID NOs 370 and 413;  
 99-148-140 use SEQ ID NOs 370 and 413; 99-148-182 use SEQ ID NOs 370 and 413;  
 99-148-366 use SEQ ID NOs 370 and 413; 99-148-76 use SEQ ID NOs 370 and 413; 99-1480-  
 290 use SEQ ID NOs 371 and 414;  
 25 99-1481-285 use SEQ ID NOs 372 and 415;  
 99-1484-101 use SEQ ID NOs 373 and 416;  
 99-1484-328 use SEQ ID NOs 373 and 416;  
 99-1485-251 use SEQ ID NOs 374 and 417;  
 99-1490-381 use SEQ ID NOs 375 and 418;  
 30 99-1493-280 use SEQ ID NOs 376 and 419; 99-151-94 use SEQ ID NOs 377 and 420;  
 99-211-291 use SEQ ID NOs 378 and 421; 99-213-37 use SEQ ID NOs 379 and 422;  
 99-221-442 use SEQ ID NOs 380 and 423; 99-222-109 use SEQ ID NOs 381 and 424; and the  
 compliments thereof.

- Primers with their 3' ends located 1 nucleotide upstream or downstream of a PG1-related  
 35 biallelic marker have a special utility in microsequencing assays. Preferred microsequencing primers  
 include the polynucleotides from position 1 to position 23 and from position 25 to position 47 of SEQ

- ID NOs: 21-38, and as well as the compliments thereof. Additional preferred microsequencing primers for particular non-genic PG1-related biallelic markers are listed as follows by the internal reference number for the marker and the SEQ ID NOs of the two preferred microsequencing primers:
- 4-14-107 of SEQ ID NOs 425 and 502\*; 4-14-317 of SEQ ID NOs 426 and 503\*;  
5 4-14-35 of SEQ ID NOs 427 and 504\*; 4-20-149 of SEQ ID NOs 428\* and 505;  
4-20-77 of SEQ ID NOs 429 and 506; 4-22-174 of SEQ ID NOs 430\* and 507;  
4-22-176 of SEQ ID NOs 431 and 508; 4-26-60 of SEQ ID NOs 432 and 509\*;  
4-26-72 of SEQ ID NOs 433 and 510; 4-3-130 of SEQ ID NOs 434 and 511\*;  
4-38-63 of SEQ ID NOs 435 and 512; 4-38-83 of SEQ ID NOs 436 and 513\*;  
10 4-4-152 of SEQ ID NOs 437 and 514; 4-4-187 of SEQ ID NOs 438\* and 515;  
4-4-288 of SEQ ID NOs 439 and 516; 4-42-304 of SEQ ID NOs 440 and 517;  
4-42-401 of SEQ ID NOs 441\* and 518; 4-43-328 of SEQ ID NOs 442 and 519;  
4-43-70 of SEQ ID NOs 443\* and 520; 4-50-209 of SEQ ID NOs 444\* and 521;  
4-50-293 of SEQ ID NOs 445\* and 522; 4-50-323 of SEQ ID NOs 446\* and 523;  
15 4-50-329 of SEQ ID NOs 447\* and 524; 4-50-330 of SEQ ID NOs 448 and 525;  
4-52-163 of SEQ ID NOs 449\* and 526; 4-52-88 of SEQ ID NOs 450\* and 527;  
4-53-258 of SEQ ID NOs 451 and 528\*; 4-54-283 of SEQ ID NOs 452\* and 529;  
4-54-388 of SEQ ID NOs 453 and 530; 4-55-70 of SEQ ID NOs 454 and 531\*;  
4-55-95 of SEQ ID NOs 455\* and 532; 4-56-159 of SEQ ID NOs 456\* and 533;  
20 4-56-213 of SEQ ID NOs 457 and 534; 4-58-289 of SEQ ID NOs 458\* and 535;  
4-58-318 of SEQ ID NOs 459\* and 536; 4-60-266 of SEQ ID NOs 460\* and 537;  
4-60-293 of SEQ ID NOs 461\* and 538; 4-84-241 of SEQ ID NOs 462 and 539\*;  
4-84-262 of SEQ ID NOs 463 and 540; 4-86-206 of SEQ ID NOs 464 and 541\*;  
4-86-309 of SEQ ID NOs 465 and 542; 4-88-349 of SEQ ID NOs 466 and 543.;  
25 4-89-87 of SEQ ID NOs 467\* and 544.; 99-123-184 of SEQ ID NOs 468 and 545;  
99-128-202 of SEQ ID NOs 469 and 546; 99-128-275 of SEQ ID NOs 470 and 547;  
99-128-313 of SEQ ID NOs 471 and 548; 99-128-60 of SEQ ID NOs 472\* and 549;  
99-12907-295 of SEQ ID NOs 473 and 550\*;  
99-130-58 of SEQ ID NOs 474\* and 551\*;  
30 99-134-362 of SEQ ID NOs 475 and 552\*; 99-140-130 of SEQ ID NOs 476\* and 553\*;  
99-1462-238 of SEQ ID NOs 477\* and 554; 99-147-181 of SEQ ID NOs 478 and 555\*;  
99-1474-156 of SEQ ID NOs 479 and 556\*; 99-1474-359 of SEQ ID NOs 480 and 557;  
99-1479-158 of SEQ ID NOs 481\* and 558; 99-1479-379 of SEQ ID NOs 482 and 559;  
99-148-129 of SEQ ID NOs 483 and 560; 99-148-132 of SEQ ID NOs 484 and 561;  
35 99-148-139 of SEQ ID NOs 485 and 562; 99-148-140 of SEQ ID NOs 486 and 563;  
99-148-182 of SEQ ID NOs 487 and 564\*; 99-148-366 of SEQ ID NOs 488 and 565;



- 99-148-76 of SEQ ID NOs 489 and 566; 99-1480-290 of SEQ ID NOs 490 and 567\*;  
 99-1481-285 of SEQ ID NOs 491 and 568\*; 99-1484-101 of SEQ ID NOs 492 and 569;  
 99-1484-328 of SEQ ID NOs 493\* and 570;  
 99-1485-251 of SEQ ID NOs 494 and 571\*;  
 5 99-1490-381 of SEQ ID NOs 495\* and 572;  
 99-1493-280 of SEQ ID NOs 496 and 573\*;  
 99-151-94 of SEQ ID NOs 497 and 574\*; 99-211-291 of SEQ ID NOs 498\* and 575;  
 99-213-37 of SEQ ID NOs 499 and 576; 99-221-442 of SEQ ID 500 and 577;  
 99-222-109 of SEQ ID NOs 501\* and 578; and compliments thereof.
- 10 Additional preferred microsequencing primers for particular genic PG1-related biallelic markers include a polynucleotide selected from the group consisting of the nucleotide sequences from position N-X to position N-1 of SEQ ID NO:179, nucleotide sequences from position N+1 to position N+X of SEQ ID NO:179, and the compliments thereof, wherein X is equal to 15, 18, 20, 25, 30, or a range of 15 to 30, and N is equal to one of the following values: 2159; 2443; 4452; 5733; 8438;  
 15 11843; 1983; 12080; 12221; 12947; 13147; 13194; 13310; 13342; 13367; 13594; 13680; 13902; 16231; 16388; 17608; 18034; 18290; 18786; 22835; 22872; 25183; 25192; 25614; 26911; 32703; 34491; 34756; 34934; 5160; 39897; 40598; 40816; 40947; 45783; 47929; 48206; 48207; 49282; 50037; 50054; 50101; 50220; 50440; 50562; 50653; 50660; 50745; 50885; 51249; 51333; 51435; 51468; 51515; 51557; 51566; 51632; 51666; 52016; 52096; 52151; 52282; 52348; 52410; 52580;  
 20 52712; 52772; 52860; 53092; 53272; 53389; 53511; 53600; 53665; 53815; 54365; and 54541.
- The probes of the present invention is designed from the disclosed sequences for any method known in the art, particularly methods which allow for testing if a particular sequence or marker disclosed herein is present. A preferred set of probes is designed for use in the hybridization assays of the invention in any manner known in the art such that they selectively bind to one allele of a biallelic  
 25 marker, but not the other under any particular set of assay conditions. Preferred hybridization probes may consists of, consist essentially of, or comprise a contiguous span which ranges in length from 8, 10, 12, 15, 18 or 20 to 25, 35, 40, 50, 60, 70, or 80 nucleotides, or be specified as being 12, 15, 18, 20, 25, 35, 40, or 50 nucleotides in length and including a PG1-related biallelic marker of said sequence. Optionally either of the two alleles specified in the definition of PG1-related biallelic marker is  
 30 specified as being present at the biallelic marker site. Optionally, said biallelic marker is within 6, 5, 4, 3, 2, or 1 nucleotides of the center of the hybridization probe or at the center of said probe. A preferred set of hybridization probes is disclosed in SEQ ID NOs: 21-38, 57-62, 185-338, and the compliments thereof. Another particularly preferred set of hybridization probes includes the polynucleotides from position X to position Y of any one of SEQ ID NOs: 21-38, 57-62, 185-338, or  
 35 the compliments thereof, wherein X is equal to 5, 8, 10, 12, 14, 16, 18 or a range of 5 to 18, and Y is equal to 30, 32, 34, 36, 38, 40, 43 or a range of 30 to 43; preferably X equals 12 and Y equals 36.

Additional preferred hybridization probes for particular genic PG1-related biallelic markers include a polynucleotide selected from the group consisting of the nucleotide sequences from position N-X to position N+Y of SEQ ID NO:179, and the compliments thereof, wherein X is equal to 8, 10, 12, 15, 20, 25, or a range of 8 to 30, Y is equal to 8, 10, 12, 15, 20, 25, or a range of 8 to 30, and N is equal to  
5 one of the following values: 2159; 2443; 4452; 5733; 8438; 11843; 1983; 12080; 12221; 12947; 13147; 13194; 13310; 13342; 13367; 13594; 13680; 13902; 16231; 16388; 17608; 18034; 18290; 18786; 22835; 22872; 25183; 25192; 25614; 26911; 32703; 34491; 34756; 34934; 5160; 39897; 40598; 40816; 40947; 45783; 47929; 48206; 48207; 49282; 50037; 50054; 50101; 50220; 50440; 50562; 50653; 50660; 50745; 50885; 51249; 51333; 51435; 51468; 51515; 51557; 51566; 51632;  
10 51666; 52016; 52096; 52151; 52282; 52348; 52410; 52580; 52712; 52772; 52860; 53092; 53272; 53389; 53511; 53600; 53665; 53815; 54365; and 54541; wherein the nucleotide at position N is selected from one of the two alleles specified in the definition of PG1-related biallelic marker at the biallelic marker site at position N.

Any of the polynucleotides of the present invention can be labeled, if desired, by  
15 incorporating a label detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. For example, useful labels include radioactive substances, fluorescent dyes or biotin. Preferably, polynucleotides are labeled at their 3' and 5' ends. A label can also be used to capture the primer, so as to facilitate the immobilization of either the primer or a primer extension product, such as amplified DNA, on a solid support. A capture label is attached to the primers or  
20 probes and can be a specific binding member which forms a binding pair with the solid's phase reagent's specific binding member (e.g. biotin and streptavidin). Therefore depending upon the type of label carried by a polynucleotide or a probe, it is employed to capture or to detect the target DNA. Further, it will be understood that the polynucleotides, primers or probes provided herein, may, themselves, serve as the capture label. For example, in the case where a solid phase reagent's binding  
25 member is a nucleic acid sequence, it is selected such that it binds a complementary portion of a primer or probe to thereby immobilize the primer or probe to the solid phase. In cases where a polynucleotide probe itself serves as the binding member, those skilled in the art will recognize that the probe will contain a sequence or "tail" that is not complementary to the target. In the case where a polynucleotide primer itself serves as the capture label, at least a portion of the primer will be free to  
30 hybridize with a nucleic acid on a solid phase. DNA Labeling techniques are well known to the skilled technician.

Any of the polynucleotides, primers and probes of the present invention can be conveniently immobilized on a solid support. Solid supports are known to those skilled in the art and include the walls of wells of a reaction tray, test tubes, polystyren beads, magnetic beads, nitrocellulose strips,  
35 membranes, microparticles such as latex particles, sheep (or other animal) red blood cells, duracytes® and others. The solid support is not critical and can be selected by one skilled in the art. Thus, latex

particles, microparticles, magnetic or non-magnetic beads, membranes, plastic tubes, walls of microtiter wells, glass or silicon chips, sheep (or other suitable animal's) red blood cells and duracytes are all suitable examples. Suitable methods for immobilizing nucleic acids on solid phases include ionic, hydrophobic, covalent interactions and the like. A solid support, as used herein, refers to any material which is insoluble, or can be made insoluble by a subsequent reaction. The solid support can be chosen for its intrinsic ability to attract and immobilize the capture reagent. Alternatively, the solid phase can retain an additional receptor which has the ability to attract and immobilize the capture reagent. The additional receptor can include a charged substance that is oppositely charged with respect to the capture reagent itself or to a charged substance conjugated to the capture reagent. As yet another alternative, the receptor molecule can be any specific binding member which is immobilized upon (attached to) the solid support and which has the ability to immobilize the capture reagent through a specific binding reaction. The receptor molecule enables the indirect binding of the capture reagent to a solid support material before the performance of the assay or during the performance of the assay. The solid phase thus can be a plastic, derivatized plastic, magnetic or non-magnetic metal, glass or silicon surface of a test tube, microtiter well, sheet, bead, microparticle, chip, sheep (or other suitable animal's) red blood cells, duracytes® and other configurations known to those of ordinary skill in the art. The polynucleotides of the invention can be attached to or immobilized on a solid support individually or in groups of at least 2, 5, 8, 10, 12, 15, 20, or 25 distinct polynucleotides of the inventions to a single solid support. In addition, polynucleotides other than those of the invention may be attached to the same solid support as one or more polynucleotides of the invention.

Any polynucleotide provided herein is attached in overlapping areas or at random locations on the solid support. Alternatively the polynucleotides of the invention is attached in an ordered array wherein each polynucleotide is attached to a distinct region of the solid support which does not overlap with the attachment site of any other polynucleotide. Preferably, such an ordered array of polynucleotides is designed to be "addressable" where the distinct locations are recorded and can be accessed as part of an assay procedure. Addressable polynucleotide arrays typically comprise a plurality of different oligonucleotide probes that are coupled to a surface of a substrate in different known locations. The knowledge of the precise location of each polynucleotide's location makes these "addressable" arrays particularly useful in hybridization assays. Any addressable array technology known in the art can be employed with the polynucleotides of the invention. One particular embodiment of these polynucleotide arrays is known as the Genechips™, and has been generally described in US Patent 5,143,854; PCT publications WO 90/15070 and 92/10092. These arrays may generally be produced using mechanical synthesis methods or light directed synthesis methods, which incorporate a combination of photolithographic methods and solid phase oligonucleotide synthesis (Fodor et al., Science, 251:767-777, 1991). The immobilization of arrays of oligonucleotides on solid

supports has been rendered possible by the development of a technology generally identified as "Very Large Scale Immobilized Polymer Synthesis" (VLSIPS™) in which, typically, probes are immobilized in a high density array on a solid surface of a chip. Examples of VLSIPS™ technologies are provided in US Patents 5,143,854 and 5,412,087 and in PCT Publications WO 90/15070, WO 92/10092 and WO 95/11995, which describe methods for forming oligonucleotide arrays through techniques such as light-directed synthesis techniques. In designing strategies aimed at providing arrays of nucleotides immobilized on solid supports, further presentation strategies were developed to order and display the oligonucleotide arrays on the chips in an attempt to maximize hybridization patterns and sequence information. Examples of such presentation strategies are disclosed in PCT Publications WO 94/12305, WO 94/11530, WO 97/29212 and WO 97/31256.

Oligonucleotide arrays may comprise at least one of the sequences selected from the group consisting of SEQ ID NOs: 3, 21-38, 57-62, 100-124, 179, 185-338, the preferred hybridization probes for genic PG1-related biallelic markers described above; and the sequences complementary thereto; or a fragment thereof of at least 15 consecutive nucleotides for determining whether a sample contains one or more alleles of the biallelic markers of the present invention. Oligonucleotide arrays may also comprise at least one of the sequences selected from the group consisting of SEQ ID NOs: 179, 339-424; and the sequences complementary thereto or a fragment thereof of at least 15 consecutive nucleotides for amplifying one or more alleles of the PG1-related biallelic markers. In other embodiments, arrays may also comprise at least one of the sequences selected from the group consisting of SEQ ID 425-578, the preferred microsequencing primers for genic PG1-related biallelic markers described above; and the sequences complementary thereto or a fragment thereof of at least 15 consecutive nucleotides for conducting microsequencing analyses to determine whether a sample contains one or more alleles of PG1-related biallelic marker.

The present invention further encompasses polynucleotide sequences that hybridize to any one of SEQ ID NOs: 3, 69, 100-112, or 179-184 under conditions of high or intermediate stringency as described below:

(i) By way of example and not limitation, procedures using conditions of high stringency are as follows: Prehybridization of filters containing DNA is carried out for 8 h to overnight at 65°C in buffer composed of 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 µg/ml denatured salmon sperm DNA. Filters are hybridized for 48 h at 65°C, the preferred hybridization temperature, in prehybridization mixture containing 100 µg/ml denatured salmon sperm DNA and 5-20 X 10<sup>6</sup> cpm of <sup>32</sup>P-labeled probe. Alternatively, the hybridization step can be performed at 65°C in the presence of SSC buffer, 1 x SSC corresponding to 0.15M NaCl and 0.05 M Na citrate. Subsequently, filter washes can be done at 37°C for 1 h in a solution containing 2X SSC, 0.01% PVP, 0.01% Ficoll, and 0.01% BSA, followed by a wash in 0.1X SSC at 50°C for 45 min. Alternatively, filter washes can be performed in a solution containing 2 x SSC and 0.1% SDS,

or 0.5 x SSC and 0.1% SDS, or 0.1 x SSC and 0.1% SDS at 68°C for 15 minute intervals. Following the wash steps, the hybridized probes are detectable by autoradiography. Other conditions of high stringency which is used are well known in the art and as cited in Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, Second Edition, Cold Spring Harbor Press, N.Y., pp. 9.47-9.57; and  
5 Ausubel et al., 1989, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y. Preferably, such sequences encode a homolog of a polypeptide encoded by one of ORF2 to ORF1297. In one embodiment, such sequences encode a mammalian PG1 polypeptide.

(ii) By way of example and not limitation, procedures using conditions of intermediate stringency are as follows: Filters containing DNA are prehybridized, and then hybridized at a  
10 temperature of 60°C in the presence of a 5 x SSC buffer and labeled probe. Subsequently, filters washes are performed in a solution containing 2x SSC at 50°C and the hybridized probes are detectable by autoradiography. Other conditions of intermediate stringency which is used are well known in the art and as cited in Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, Second Edition, Cold Spring Harbor Press, N.Y., pp. 9.47-9.57; and Ausubel et al., 1989, Current  
15 Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y. Preferably, such sequences encode a homolog of a polypeptide encoded by one of SEQ ID NOs: 3, 69, 100-112, or 179-184. In one embodiment, such sequences encode a mammalian PG1 polypeptide.

The present invention also encompasses diagnostic kits comprising one or more polynucleotides of the invention with a portion or all of the necessary reagents and instructions for  
20 genotyping a test subject by determining the identity of a nucleotide at a PG1-related biallelic marker. The polynucleotides of a kit may optionally be attached to a solid support, or be part of an array or addressable array of polynucleotides. The kit may provide for the determination of the identity of the nucleotide at a marker position by any method known in the art including, but not limited to, a sequencing assay method, a microsequencing assay method, a hybridization assay method, or an allele  
25 specific amplification method. Optionally such a kit may include instructions for scoring the results of the determination with respect to the test subjects' risk of contracting a cancer or prostate cancer, or likely response to an anti-cancer agent or anti-prostate cancer agent, or chances of suffering from side effects to an anti-cancer agent or anti-prostate cancer agent.

#### Use of PG1 Nucleic Acids as Reagents

30 The PG1 genomic DNA of SEQ ID NO: 179, the PG1 cDNA of SEQ ID NO: 3, 112-124 and PG1 alleles responsible for a detectable phenotype (such as those obtainable by the methods of Example 12, and SEQ ID NO:69) can be used to prepare PCR primers for use in diagnostic techniques or genetic engineering methods such as those described above. Example 10 describes the use of the PG1 genomic DNA of SEQ ID NO: 179, the PG1 cDNA of SEQ ID NO: 3, 112-124 and PG1 alleles responsible for a  
35 detectable phenotype (such as those obtainable by the methods of Example 12) in PCR amplification procedures.

### Example 10

The PG1 genomic DNA of SEQ ID NO: 179, the PG1 cDNA of SEQ ID NO: 3, and PG1 alleles responsible for a detectable phenotype (such as those obtainable by the methods of Example 12) is used to prepare PCR primers for a variety of applications, including isolation procedures for cloning nucleic acids capable of hybridizing to such sequences, diagnostic techniques and forensic techniques. The PCR primers comprise at least 10 consecutive bases of the PG1 genomic DNA of SEQ ID NO: 179, the PG1 cDNA of SEQ ID NO: 3, 112-124 and PG1 alleles responsible for a detectable phenotype (such as those obtainable by the methods of Example 12) or the sequences complementary thereto. Preferably, the PCR primers comprise at least 12, 15, or 17 consecutive bases of these sequences. More preferably, the PCR primers comprise at least 20-30 consecutive bases of the PG1 genomic DNA of SEQ ID NO: 179, the PG1 cDNA of SEQ ID NO: 3, 112-124 and PG1 alleles responsible for a detectable phenotype (such as those obtainable by the methods of Example 12) or the sequences complementary thereto. In some embodiments, the PCR primers may comprise more than 30 consecutive bases of the PG1 genomic DNA of SEQ ID NO: 179, the PG1 cDNA of SEQ ID NO: 3, 112-124 and PG1 alleles responsible for a detectable phenotype (such as those obtainable by the methods of Example 12) or the sequences complementary thereto. It is preferred that the primer pairs to be used together in a PCR amplification have approximately the same G/C ratio, so that melting temperatures are approximately the same.

A variety of PCR techniques are familiar to those skilled in the art. For a review of PCR technology, see Molecular Cloning to Genetic Engineering White, B.A. Ed. in Methods in Molecular Biology 67: Humana Press, Totowa 1997. In each of these PCR procedures, PCR primers on either side of the nucleic acid sequences to be amplified are added to a suitably prepared nucleic acid sample along with dNTPs and a thermostable polymerase such as Taq polymerase, Pfu polymerase, or Vent polymerase. The nucleic acid in the sample is denatured and the PCR primers are specifically hybridized to complementary nucleic acid sequences in the sample. The hybridized primers are extended. Thereafter, another cycle of denaturation, hybridization, and extension is initiated. The cycles are repeated multiple times to produce an amplified fragment containing the nucleic acid sequence between the primer sites.

The polynucleotides of the Invention also encompass vectors and DNA constructs as well as other forms of primers and probes. For a thorough description of these embodiments please see Sections VIII, X, and XI below.

## III. POLYPEPTIDES

### PG1 Proteins and Polypeptide Fragments

The term "PG1 polypeptides" is used herein to embrace all of the proteins and polypeptides of the present invention. Also forming part of the invention are polypeptides encoded by the polynucleotides of the invention, as well as fusion polypeptides comprising such polypeptides. The invention embodies PG1 proteins from human (SEQ ID NOs: 4, and 5), and mouse (SEQ ID NO: 74).

However, PG1 species from other varieties of mammals are expressly contemplated and is isolated using the antibodies of the present invention in conjunction with standard affinity chromatography methods as well as being expressed from the PG1 genes isolated from other mammalian sources using human and mouse PG1 nucleic acid sequences as primers and probes as well as the methods described  
5 herein.

The invention also embodies PG1 proteins translated from less common alternative splice species, including SEQ ID NOs: 125-136, and PG1 proteins which result from naturally occurring mutant, particularly functional mutants of PG1, including SEQ ID NO: 70, which is identified and obtained by the described herein. The present invention also embodies polypeptides comprising a  
10 contiguous stretch of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 50, or 100 amino acids of a PG1 protein. In a preferred embodiment the contiguous stretch of amino acids comprises the site of a mutation or functional mutation, including a deletion, addition, swap or truncation of the amino acids in the PG1 protein sequence. For instance, polypeptides that contain either the Arg and His residues at amino acid position 184, and polypeptides  
15 that contain either the Arg or Ile residue at amino acid position 293 of the SEQ ID NO: 4 in said contiguous stretch are particularly preferred embodiments of the invention and useful in the manufacture of antibodies to detect the presence and absence of these mutations. Similarly, polypeptides with a carboxy terminus at position 228 is a particularly preferred embodiment of the invention and useful in the manufacture of antibodies to detect the presence and absence of the  
20 mutation shown in SEQ ID NOs: 69 and 70.

Similarly, polypeptides that contain an peptide sequences of 8, 10, 12, 15, or 25 amino acids encoded over a naturally-occurring splice junction (the point at which two human PG1 exon (SEQ ID NOs: 100-111) are covalently linked) in said contiguous stretch are particularly preferred embodiments and useful in the manufacture of antibodies to detect the presence, localization, and  
25 quantity of the various protein products of the PG1 alternative splice species.

PG1 proteins are preferably isolated from human, mouse or mammalian tissue samples or expressed from human, mouse or mammalian genes.

The PG1 polypeptides of the invention can be made using routine expression methods known in the art, see, for instance, Example 11, below. The polynucleotide encoding the desired  
30 polypeptide, is ligated into an expression vector suitable for any convenient host. Both eukaryotic and prokaryotic host systems is used in forming recombinant polypeptides, and a summary of some of the more common systems are included in Sections II and VIII. The polypeptide is then isolated from lysed cells or from the culture medium and purified to the extent needed for its intended use. Purification is by any technique known in the art, for example, differential extraction, salt  
35 fractionation, chromatography, centrifugation, and the like. See, for example, *Methods in Enzymology* for a variety of methods for purifying proteins.

In addition, shorter protein fragments is produced by chemical synthesis. Alternatively the proteins of the invention is extracted from cells or tissues of humans or non-human animals. Methods for purifying proteins are known in the art, and include the use of detergents or chaotropic agents to disrupt particles followed by differential extraction and separation of the polypeptides by ion exchange chromatography, affinity chromatography, sedimentation according to density, and gel electrophoresis.

#### Expression of the PG1 Protein

Any PG1 cDNA, including SEQ ID NO: 3, 69, 112-124, or 184 or synthetic DNAs is use as described in Example 11 below to express PG1 proteins and polypeptides.

10

#### Example 11

The nucleic acid encoding the PG1 protein or polypeptide to be expressed is operably linked to a promoter in an expression vector using conventional cloning technology. The PG1 insert in the expression vector may comprise the full coding sequence for the PG1 protein or a portion thereof. For example, the PG1 derived insert may encode a polypeptide comprising at least 10 consecutive amino acids of the PG1 proteins of SEQ ID NO: 4.

15

The expression vector is any of the mammalian, yeast, insect or bacterial expression systems known in the art, see for example Section VIII. Commercially available vectors and expression systems are available from a variety of suppliers including Genetics Institute (Cambridge, MA), Stratagene (La Jolla, California), Promega (Madison, Wisconsin), and Invitrogen (San Diego, California). If desired, to enhance expression and facilitate proper protein folding, the codon context and codon pairing of the sequence is optimized for the particular expression organism in which the expression vector is introduced, as explained by Hatfield, et al., U.S. Patent No. 5,082,767.

20

The following is provided as one exemplary method to express the PG1 protein or a portion thereof. In one embodiment, the entire coding sequence of the PG1 cDNA through the poly A signal of the cDNA are operably linked to a promoter in the expression vector. Alternatively, if the nucleic acid encoding a portion of the PG1 protein lacks a methionine to serve as the initiation site, an initiating methionine can be introduced next to the first codon of the nucleic acid using conventional techniques. Similarly, if the insert from the PG1 cDNA lacks a poly A signal, this sequence can be added to the construct by, for example, splicing out the Poly A signal from pSG5 (Stratagene) using BglI and SalI restriction endonuclease enzymes and incorporating it into the mammalian expression vector pXT1 (Stratagene). pXT1 contains the LTRs and a portion of the gag gene from Moloney Murine Leukemia Virus. The position of the LTRs in the construct allow efficient stable transfection. The vector includes the Herpes Simplex Thymidine Kinase promoter and the selectable neomycin gene. The nucleic acid encoding the PG1 protein or a portion thereof is obtained by PCR from a bacterial vector containing the PG1 cDNA of SEQ ID NO: 3 using oligonucleotide primers complementary to the PG1 cDNA or portion thereof and containing restriction endonuclease sequences for Pst I incorporated into the 5' primer and

30

35



BglII at the 5' end of the corresponding cDNA 3' primer, taking care to ensure that the sequence encoding the PG1 protein or a portion thereof is positioned properly with respect to the poly A signal. The purified fragment obtained from the resulting PCR reaction is digested with PstI, blunt ended with an exonuclease, digested with Bgl II, purified and ligated to pXT1, now containing a poly A signal and digested with BglII.

The ligated product is transfected into mouse NIH 3T3 cells using Lipofectin (Life Technologies, Inc., Grand Island, New York) under conditions outlined in the product specification. Positive transfectants are selected after growing the transfected cells in 600ug/ml G418 (Sigma, St. Louis, Missouri).

Alternatively, the nucleic acids encoding the PG1 protein or a portion thereof is cloned into pED6dpc2 (Genetics Institute, Cambridge, MA). The resulting pED6dpc2 constructs is transfected into a suitable host cell, such as COS 1 cells. Methotrexate resistant cells are selected and expanded.

The above procedures may also be used to express a mutant PG1 protein responsible for a detectable phenotype or a portion thereof.

The expressed proteins is purified using conventional purification techniques such as ammonium sulfate precipitation or chromatographic separation based on size or charge. The protein encoded by the nucleic acid insert may also be purified using standard immunochromatography techniques. In such procedures, a solution containing the expressed PG1 protein or portion thereof, such as a cell extract, is applied to a column having antibodies against the PG1 protein or portion thereof is attached to the chromatography matrix. The expressed protein is allowed to bind the immunochromatography column. Thereafter, the column is washed to remove non-specifically bound proteins. The specifically bound expressed protein is then released from the column and recovered using standard techniques.

To confirm expression of the PG1 protein or a portion thereof, the proteins expressed from host cells containing an expression vector containing an insert encoding the PG1 protein or a portion thereof can be compared to the proteins expressed in host cells containing the expression vector without an insert. The presence of a band in samples from cells containing the expression vector with an insert which is absent in samples from cells containing the expression vector without an insert indicates that the PG1 protein or a portion thereof is being expressed. Generally, the band will have the mobility expected for the PG1 protein or portion thereof. However, the band may have a mobility different than that expected as a result of modifications such as glycosylation, ubiquitination, or enzymatic cleavage.

Antibodies capable of specifically recognizing the expressed PG1 protein or a portion thereof is generated as described below in Section VII.

If antibody production is not possible, the nucleic acids encoding the PG1 protein or a portion thereof is incorporated into expression vectors designed for use in purification schemes employing chimeric polypeptides. In such strategies the nucleic acid encoding the PG1 protein or a portion thereof is inserted in frame with the gene encoding the other half of the chimera. The other half of the chimera

is  $\beta$ -globin or a nickel binding polypeptide encoding sequence. A chromatography matrix having antibody to  $\beta$ -globin or nickel attached thereto is then used to purify the chimeric protein. Protease cleavage sites is engineered between the  $\beta$ -globin gene or the nickel binding polypeptide and the PG1 protein or portion thereof. Thus, the two polypeptides of the chimera is separated from one another by  
5 protease digestion.

One useful expression vector for generating  $\beta$ -globin chimerics is pSG5 (Stratagene), which encodes rabbit  $\beta$ -globin. Intron II of the rabbit  $\beta$ -globin gene facilitates splicing of the expressed transcript, and the polyadenylation signal incorporated into the construct increases the level of expression. These techniques are well known to those skilled in the art of molecular biology. Standard methods are  
10 published in methods texts such as Davis et al., (Basic Methods in Molecular Biology, L.G. Davis, M.D. DIBNER, and J.F. BATTEY, ed., Elsevier Press, NY, 1986) and many of the methods are available from Stratagene, Life Technologies, Inc., or Promega. Polypeptide may additionally be produced from the construct using in vitro translation systems such as the In vitro Express<sup>TM</sup> Translation Kit (Stratagene).

#### **IV. IDENTIFICATION OF MUTATIONS IN THE PG1 GENE WHICH ARE ASSOCIATED** 15 **WITH A DETECTABLE PHENOTYPE**

Mutations in the PG1 gene which are responsible for a detectable phenotype is identified by comparing the sequences of the PG1 genes from affected and unaffected individuals as described in Example 12, below. The detectable phenotype may comprise a variety of manifestations of altered PG1 function, including prostate cancer, hepatocellular carcinoma, colorectal cancer, non-small cell  
20 lung cancer, squamous cell carcinoma, or other conditions. The mutations may comprise point mutations, deletions, or insertions of the PG1 gene. The mutations may lie within the coding sequence for the PG1 protein or within regulatory regions in the PG1 gene.

##### Example 12

Oligonucleotide primers are designed to amplify the sequences of each of the exons or the  
25 promoter region of the PG1 gene. The oligonucleotide primers may comprise at least 10 consecutive nucleotides of the PG1 genomic DNA of SEQ ID NO:179 or the PG1 cDNA of SEQ ID NO: 3 or the sequences complementary thereto. Preferably, the oligonucleotides comprise at least 15 consecutive nucleotides of the PG1 genomic DNA of SEQ ID NO:179 or the PG1 cDNA of SEQ ID NO: 3 or the sequences complementary thereto. In some embodiments, the oligonucleotides may comprise at least  
30 20 consecutive nucleotides of the PG1 genomic DNA of SEQ ID NO: 179 or the PG1 cDNA of SEQ ID NO:3 or the sequences complementary thereto. In other embodiments, the oligonucleotides may comprise 25 or more consecutive nucleotides of the PG1 genomic DNA of SEQ ID NO: 179 or the PG1 cDNA of SEQ ID NO: 3 or the sequences complementary thereto.

Each primer pair is used to amplify the exon or promoter region from which it is derived.  
35 Amplification is carried out on genomic DNA samples from affected patients and unaffected controls using the PCR conditions described above. Amplification products from the genomic PCRs are

subjected to automated dideoxy terminator sequencing reactions and electrophoresed on ABI 377 sequencers. Following gel image analysis and DNA sequence extraction, ABI sequence data are automatically analyzed to detect the presence of sequence variations among affected and unaffected individuals. Sequences are verified by determining the sequences of both DNA strands for each individual. Preferably, these candidate mutations are detected by comparing individuals homozygous for haplotype 5 of Figure 4 and controls not carrying haplotype 5 or related haplotypes.

Candidate polymorphisms suspected of being responsible for the detectable phenotype, such as prostate cancer or other conditions, are then verified by screening a larger population of affected and unaffected individuals using the microsequencing technique described above. Polymorphisms which exhibit a statistically significant correlation with the detectable phenotype are deemed responsible for the detectable phenotype.

Other techniques may also be used to detect polymorphisms associated with a detectable phenotype such as prostate cancer or other conditions. For example, polymorphisms is detected using single stranded conformation analyses such as those described in Orita et al., Proc. Natl. Acad. Sci. U.S.A. 86: 2776-2770 (1989). In this approach, polymorphisms are detected through altered migration on SSCA gels.

Alternatively, polymorphisms is identified using clamped denaturing gel electrophoresis, heteroduplex analysis, chemical mismatch cleavage, and other conventional techniques as described in Sheffield, V.C. et al, Proc. Natl. Acad. Sci. U.S.A 49:699-706 (1991); White, M.B. et al., Genomics 12:301-306 (1992); Grompe, M. et al., Proc. Natl. Acad. Sci. U.S.A 86:5855-5892 (1989); and Grompe, M. Nature Genetics 5:111-117 (1993).

The PG1 genes from individuals carrying PG1 mutations responsible for the detectable phenotype, or cDNAs derived therefrom, is cloned as follows. Nucleic acid samples are obtained from individuals having a PG1 mutation associated with the detectable phenotype. The nucleic acid samples are contacted with a probe derived from the PG1 genomic DNA of SEQ ID NO: 179 or the PG1 cDNA of SEQ ID NO:3. Nucleic acids containing the mutant PG1 allele are identified using conventional techniques. For example, the mutant PG1 gene, or a cDNA derived therefrom, is obtained by conducting an amplification reaction using primers derived from the PG1 genomic DNA of SEQ ID NO: 179 or the PG1 cDNA of SEQ ID NO:3. Alternatively, the mutant PG1 gene, or a cDNA derived therefrom, is identified by hybridizing a genomic library or a cDNA library obtained from an individual having a mutant PG1 gene with a detectable probe derived from the PG1 genomic DNA of SEQ ID NO: 179 or the PG1 cDNA of SEQ ID NO: 3. Alternatively, the mutant PG1 allele is obtained by contacting an expression library from an individual carrying a PG1 mutation with a detectable antibody against the PG1 proteins of SEQ ID NO: 4 or SEQ ID NO: 5 which has been prepared as described below. Those skilled in the art will appreciate that the PG1 genomic DNA of

SEQ ID NO: 179, the PG1 cDNA of SEQ ID NO: 3 and the PG1 proteins of SEQ ID NOs: 4 and 5 is used in a variety of other conventional techniques to obtain the mutant PG1 gene.

In another embodiment the mutant PG1 allele which causes a detectable phenotype can be isolated by obtaining a nucleic acid sample such as a genomic library or a cDNA library from an individual expressing the detectable phenotype. The nucleic acid sample can be contacted with one or more probes lying in the 8p23 region of the human genome. Nucleic acids in the sample which contain the PG1 gene can be identified by conducting sequencing reactions on the nucleic acids which hybridize to the markers from the 8p23 region of the human genome.

The region of the PG1 gene containing the mutation responsible for the detectable phenotype may also be used in diagnostic techniques such as those described below. For example, oligonucleotides containing the mutation responsible for the detectable phenotype is used in amplification or hybridization based diagnostics, such as those described herein, for detecting individuals suffering from the detectable phenotype or individuals at risk of developing the detectable phenotype at a subsequent time. In addition, the PG1 allele responsible for the detectable phenotype is used in gene therapy as described herein. The PG1 allele responsible for the detectable phenotype may also be cloned into an expression vector to express the mutant PG1 protein as described herein.

During the search for biallelic markers associated with prostate cancer, a number of polymorphic bases were discovered which lie within the PG1 gene. The identities and positions of these polymorphic bases are listed as features in the accompanying Sequence Listing for the PG1 genomic DNA of SEQ ID NO: 179. The polymorphic bases is used in the above-described diagnostic techniques for determining whether an individual is at risk for developing prostate cancer at a subsequent date or suffers from prostate cancer as a result of a PG1 mutation. The identities of the nucleotides present at the polymorphic positions in a nucleic acid sample is determined using the techniques, such as microsequencing analysis, which are described above.

It is possible that one or more of these polymorphisms (or other polymorphic bases) is mutations which are associated with prostate cancer. To determine whether a polymorphism is responsible for prostate cancer, the frequency of each of the alleles in individuals suffering from prostate cancer and unaffected individuals is measured as described in the haplotype analysis above. Those mutations which occur at a statistically significant frequency in the affected population are deemed to be responsible for prostate cancer.

cDNAs containing the identified mutant PG1 gene is prepared as described above and cloned into expression vectors as described below. The proteins expressed from the expression vectors is used to generate antibodies specific for the mutant PG1 proteins as described below. In addition, allele specific probes containing the PG1 mutation responsible for prostate cancer is used in the diagnostic techniques described below.

Genes sharing homology to the PG1 gene is identified as follows.

Example 13

Alternatively, a cDNA library or genomic DNA library to be screened for genes sharing homology to the PG1 gene is obtained from a commercial source or made using techniques familiar to those skilled in the art. The cDNA library or genomic DNA library is hybridized to a detectable probe  
5 comprising at least 10 consecutive nucleotides from the PG1 cDNA of SEQ ID NO:3, the PG1 genomic DNA of SEQ ID NO: 179, or the sequences complementary thereto, using conventional techniques. Preferably, the probe comprises at least 12, 15, or 17 consecutive nucleotides from the PG1 cDNA of SEQ ID NO:3, the PG1 genomic DNA of SEQ ID NO: 179, or the sequences complementary thereto. More preferably, the probe comprises at least 20-30 consecutive nucleotides from the PG1 cDNA of  
10 SEQ ID NO:3, the PG1 genomic DNA of SEQ ID NO: 179, or the sequences complementary thereto. In some embodiments, the probe comprises more than 30 nucleotides from the PG1 cDNA of SEQ ID NO:3, the PG1 genomic DNA of SEQ ID NO: 179, or the sequences complementary thereto.

Techniques for identifying cDNA clones in a cDNA library which hybridize to a given probe sequence are disclosed in Sambrook et al., Molecular Cloning: A Laboratory Manual 2d Ed., Cold  
15 Spring Harbor Laboratory Press, 1989. The same techniques is used to isolate genomic DNAs sharing homology with the PG1 gene.

Briefly, cDNA or genomic DNA clones which hybridize to the detectable probe are identified and isolated for further manipulation as follows. A probe comprising at least 10 consecutive nucleotides from the PG1 cDNA of SEQ ID NO:3, the PG1 genomic DNA of SEQ ID NO: 179, or the sequences  
20 complementary thereto, is labeled with a detectable label such as a radioisotope or a fluorescent molecule. Preferably, the probe comprises at least 12, 15, or 17 consecutive nucleotides from the PG1 cDNA of SEQ ID NO:3, the PG1 genomic DNA of SEQ ID NO: 179, or the sequences complementary thereto. More preferably, the probe comprises 20-30 consecutive nucleotides from the PG1 cDNA of SEQ ID NO:3, the PG1 genomic DNA of SEQ ID NO: 179, or the sequences complementary thereto. In  
25 some embodiments, the probe comprises more than 30 nucleotides from the PG1 cDNA of SEQ ID NO:3, the PG1 genomic DNA of SEQ ID NO: 179, or the sequences complementary thereto.

Techniques for labeling the probe are well known and include phosphorylation with polynucleotide kinase, nick translation, in vitro transcription, and non-radioactive techniques. The cDNAs or genomic DNAs in the library are transferred to a nitrocellulose or nylon filter and denatured.  
30 After incubation of the filter with a blocking solution, the filter is contacted with the labeled probe and incubated for a sufficient amount of time for the probe to hybridize to cDNAs or genomic DNAs containing a sequence capable of hybridizing to the probe.

By varying the stringency of the hybridization conditions used to identify cDNAs or genomic DNAs which hybridize to the detectable probe, cDNAs or genomic DNAs having different levels of  
35 homology to the probe can be identified and isolated. To identify cDNAs or genomic DNAs having a

high degree of homology to the probe sequence, the melting temperature of the probe is calculated using the following formulas:

For probes between 14 and 70 nucleotides in length the melting temperature  $T_m$  is calculated using the formula:  $T_m = 81.5 + 16.6(\log(Na^+)) + 0.41(\text{fraction G+C}) - (600/N)$  where N is the length of the probe.

If the hybridization is carried out in a solution containing formamide, the melting temperature is calculated using the equation  $T_m = 81.5 + 16.6(\log(Na^+)) + 0.41(\text{fraction G+C}) - (0.63\% \text{ formamide}) - (600/N)$  where N is the length of the probe.

Prehybridization is carried out in 6X SSC, 5X Denhardt's reagent, 0.5% SDS, 100• g denatured fragmented salmon sperm DNA or 6X SSC, 5X Denhardt's reagent, 0.5% SDS, 100• g denatured fragmented salmon sperm DNA, 50% formamide. The formulas for SSC and Denhardt's solutions are listed in Sambrook et al., supra.

Hybridization is conducted by adding the detectable probe to the prehybridization solutions listed above. Where the probe comprises double stranded DNA, it is denatured before addition to the hybridization solution. The filter is contacted with the hybridization solution for a sufficient period of time to allow the probe to hybridize to cDNAs or genomic DNAs containing sequences complementary thereto or homologous thereto. For probes over 200 nucleotides in length, the hybridization is carried out at 15-25• C below the  $T_m$ . For shorter probes, such as oligonucleotide probes, the hybridization is conducted at 15-25• C below the  $T_m$ . Preferably, for hybridizations in 6X SSC, the hybridization is conducted at approximately 68• C. Preferably, for hybridizations in 50% formamide containing solutions, the hybridization is conducted at approximately 42• C.

All of the foregoing hybridizations would be considered to be under "stringent" conditions.

Following hybridization, the filter is washed in 2X SSC, 0.1% SDS at room temperature for 15 minutes. The filter is then washed with 0.1X SSC, 0.5% SDS at room temperature for 30 minutes to 1 hour. Thereafter, the solution is washed at the hybridization temperature in 0.1X SSC, 0.5% SDS. A final wash is conducted in 0.1X SSC at room temperature.

cDNAs or genomic DNAs homologous to the PG1 gene which have hybridized to the probe are identified by autoradiography or other conventional techniques.

The above procedure is modified to identify cDNAs or genomic DNAs having decreasing levels of homology to the probe sequence. For example, to obtain cDNAs or genomic DNAs of decreasing homology to the detectable probe, less stringent conditions is used. For example, the hybridization temperature is decreased in increments of 5• C from 68• C to 42• C in a hybridization buffer having a  $Na^+$  concentration of approximately 1M. Following hybridization, the filter is washed with 2X SSC, 0.5% SDS at the temperature of hybridization. These conditions are considered to be "moderate" conditions above 50• C and "low" conditions below 50• C.

Alternatively, the hybridization is carried out in buffers, such as 6X SSC, containing formamide at a temperature of 42° C. In this case, the concentration of formamide in the hybridization buffer is reduced in 5% increments from 50% to 0% to identify clones having decreasing levels of homology to the probe. Following hybridization, the filter is washed with 6X SSC, 0.5% SDS at 50° C. These conditions are considered to be "moderate" conditions above 25% formamide and "low" conditions below 25% formamide.

cDNAs or genomic DNAs which have hybridized to the probe are identified by autoradiography.

If it is desired to obtain nucleic acids homologous to the PG1 gene, such as allelic variants thereof or nucleic acids encoding proteins related to the PG1 protein, the level of homology between the hybridized nucleic acid and the PG1 gene may readily be determined. To determine the level of homology between the hybridized nucleic acid and the PG1 gene, the nucleotide sequences of the hybridized nucleic acid and the PG1 gene are compared. For example, using the above methods, nucleic acids having at least 95% nucleic acid homology to the PG1 gene is obtained and identified. Similarly, by using progressively less stringent hybridization conditions one can obtain and identify nucleic acids having at least 90%, at least 85%, at least 80% or at least 75% homology to the PG1 gene.

To determine whether a clone encodes a protein having a given amount of homology to the PG1 protein, the amino acid sequence of the PG1 protein is compared to the amino acid sequence encoded by the hybridizing nucleic acid. Homology is determined to exist when an amino acid sequence in the PG1 protein is closely related to an amino acid sequence in the hybridizing nucleic acid. A sequence is closely related when it is identical to that of the PG1 sequence or when it contains one or more amino acid substitutions therein in which amino acids having similar characteristics have been substituted for one another. Using the above methods, one can obtain nucleic acids encoding proteins having at least 95%, at least 90%, at least 85%, at least 80% or at least 75% homology to the proteins encoded by the PG1 probe.

#### Isolation and Use of Mutant or Low Frequency PG1 Alleles from Mammalian Prostate Tumor Tissues and Cell lines

A single mutant PG1 gene was isolated from a human prostate cancer cell line. The nucleic acid sequence and amino acid sequence of this mutant PG1 are disclosed in SEQ IN NOs: 69 and 70, respectively. This mutant was found to contain a stop codon at codon position number 229, and therefore results in a truncated gene product of only 228 amino acids. The present invention encompasses purified or isolated nucleic acids comprising at least 8, 10, 12, 15, 20, or 25 consecutive nucleotides of SEQ ID NO: 69, preferably containing the mutation in codon number 229. A preferred embodiment of the present invention encompasses purified or isolated nucleic acids comprising at least 8, 10, 12, 15, 20, or 25 consecutive nucleotides of SEQ ID NO: 71.

The present invention is also directed to methods of determining whether an individual is at risk of developing prostate cancer at a later date or whether said individual suffers from prostate

cancer as a result of a mutation in the PG1 gene comprising: obtaining a nucleic acid sample from said individual; and determining whether the nucleotides present at one or more of the polymorphic bases in the sequences selected from the group consisting of SEQ ID NOs: 69 and 71 are indicative of a risk of developing prostate cancer at a later date or indicative of prostate cancer resulting from a mutation in the PG1 gene. The present invention also includes purified or isolated nucleic acids encoding at least 4, 8, 10, 12, 15, or 20 consecutive amino acids of the polypeptide of SEQ ID NO: 70, preferably including the carboxy terminus of said polypeptide. The isolated or purified polypeptides of the invention include polypeptides comprising at least 4, 8, 10, 12, 15, or 20 consecutive amino acids of the polypeptide of SEQ ID NO: 70, preferably including the carboxy terminus of said polypeptide.

**V. DIAGNOSIS OF INDIVIDUALS AT RISK FOR DEVELOPING PROSTATE CANCER OR INDIVIDUALS SUFFERING FROM PROSTATE CANCER AS A RESULT OF A MUTATION IN THE PG1 GENE**

Individuals may then be screened for the presence of polymorphisms in the PG1 gene or protein which are associated with a detectable phenotype such as cancer, prostate cancer or other conditions as described in Example 13, below. The individuals is screened while they are asymptomatic to determine their risk of developing cancer, prostate cancer or other conditions at a subsequent time. Alternatively, individuals suffering from cancer, prostate cancer or other conditions is screened for the presence of polymorphisms in the PG1 gene or protein in order to determine whether therapies which target the PG1 gene or protein should be applied.

**Example 14**

Nucleic acid samples are obtained from a symptomatic or asymptomatic individual. The nucleic acid samples is obtained from blood cells as described above or is obtained from other tissues or organs. For individuals suffering from prostate cancer, the nucleic acid sample is obtained from the tumor. The nucleic acid sample may comprise DNA, RNA, or both. The nucleotides at positions in the PG1 gene where mutations lead to prostate cancer or other detectable phenotypes are determined for the nucleic acid sample.

In one embodiment, a PCR amplification is conducted on the nucleic acid sample as described above to amplify regions in which polymorphisms associated with prostate cancer or other detectable phenotypes have been identified. The amplification products are sequenced to determine whether the individual possesses one or more PG1 polymorphisms associated with prostate cancer or other detectable phenotypes.

Alternatively, the nucleic acid sample is subjected to microsequencing reactions as described above to determine whether the individual possesses one or more PG1 polymorphisms associated with prostate cancer or another detectable phenotype resulting from a mutation in the PG1 gene.



In another embodiment, the nucleic acid sample is contacted with one or more allele specific oligonucleotides which specifically hybridize to one or more PG1 alleles associated with prostate cancer or another detectable phenotype. The nucleic acid sample is also contacted with a second PG1 oligonucleotide capable of producing an amplification product when used with the allele specific  
5 oligonucleotide in an amplification reaction. The presence of an amplification product in the amplification reaction indicates that the individual possesses one or more PG1 alleles associated with prostate cancer or another detectable phenotype.

#### Determination of PG1 Expression Levels

As discussed above, PG1 polymorphisms associated with cancer, prostate cancer or other  
10 detectable phenotypes may exert their effects by increasing, decreasing, or eliminating PG1 expression, or in altering the frequency of various transcription species. Accordingly, PG1 expression levels in individuals suffering from cancer, prostate cancer or other detectable phenotypes is compared to those of unaffected individuals to determine whether over-expression, under-expression, loss of expression, or changes in the relative frequency of transcription species of PG1 causes cancer,  
15 prostate cancer or another detectable phenotype. Individuals is tested to determine whether they are at risk of developing cancer, or prostate cancer at a subsequent time or whether they suffer from prostate cancer resulting from a mutation in the PG1 gene by determining whether they exhibit a level of PG1 expression associated with prostate cancer. Similarly, individuals is tested to determine whether they suffer from another PG1 mediated detectable phenotype or whether they are at risk of suffering from  
20 such a condition at a subsequent time.

Expression levels in nucleic acid samples from affected and unaffected individuals is determined by performing Northern blots using detectable probes derived from the PG1 gene or the PG1 cDNA. A variety of conventional Northern blotting procedures is used to detect and quantitate PG1 expression and the frequencies of the various transcription species of PG1, including those  
25 disclosed in Current Protocols in Molecular Biology, John Wiley 503 Sons, Inc. 1997 and Sambrook et al. Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, 1989.

Alternatively, PG1 expression levels is determined as described in Example 15, below.

#### Example 15

30 Expression levels and patterns of PG1 is analyzed by solution hybridization with long probes as described in International Patent Application No. WO 97/05277. Briefly, the PG1 cDNA or the PG1 genomic DNA described above, or fragments thereof, is inserted at a cloning site immediately downstream of a bacteriophage (T3, T7 or SP6) RNA polymerase promoter to produce antisense RNA. Preferably, the PG1 insert comprises at least 100 or more consecutive nucleotides of the genomic DNA  
35 sequence of SEQ ID NO: 1 or the cDNA sequences of SEQ ID NO: 3. The plasmid is linearized and transcribed in the presence of ribonucleotides comprising modified ribonucleotides (i.e. biotin-UTP and

DIG-UTP). An excess of this doubly labeled RNA is hybridized in solution with mRNA isolated from cells or tissues of interest. The hybridizations are performed under standard stringent conditions (40-50° C for 16 hours in an 80% formamide, 0.4 M NaCl buffer, pH 7-8). The unhybridized probe is removed by digestion with ribonucleases specific for single-stranded RNA (i.e. RNases CL3, T1, Phy M, U2 or A). The presence of the biotin-UTP modification enables capture of the hybrid on a microtitration plate coated with streptavidin. The presence of the DIG modification enables the hybrid to be detected and quantified by ELISA using an anti-DIG antibody coupled to alkaline phosphatase.

Quantitative analysis of PG1 gene expression may also be performed using arrays as described in Sections II and X. As used here, the term array means an arrangement of a plurality of nucleic acids of sufficient length to permit specific detection of expression of PG1 mRNAs capable of hybridizing thereto. For example, the arrays may contain a plurality of nucleic acids derived from genes whose expression levels are to be assessed. The arrays may include the PG1 genomic DNA of SEQ ID NO:179, the PG1 cDNA of SEQ ID NO:3 or the sequences complementary thereto or fragments thereof. The array may contain some or all of the known alternative splice or transcription species of PG1, including the species in SEQ ID NOs: 3, and 112-124 to determine the relative frequency of particular transcription species. Alternatively, the array may contain polynucleotides which overlap all of the potential splice junctions, including, for example SEQ ID NOs: 137-178, so that the frequency of particular splice junctions can be determined and correlated with traits or used in diagnostics just as expressions levels are. Preferably, the fragments are at least 15 nucleotides in length. In other embodiments, the fragments are at least 25 nucleotides in length. In some embodiments, the fragments are at least 50 nucleotides in length. More preferably, the fragments are at least 100 nucleotides in length. In another preferred embodiment, the fragments are more than 100 nucleotides in length. In some embodiments the fragments is more than 500 nucleotides in length.

For example, quantitative analysis of PG1 gene expression is performed with a complementary DNA microarray as described by Schena et al. (Science 270:467-470, 1995; Proc. Natl. Acad. Sci. U.S.A. 93:10614-10619, 1996). Full length PG1 cDNAs or fragments thereof are amplified by PCR and arrayed from a 96-well microtiter plate onto silylated microscope slides using high-speed robotics. Printed arrays are incubated in a humid chamber to allow rehydration of the array elements and rinsed, once in 0.2% SDS for 1 min, twice in water for 1 min and once for 5 min in sodium borohydride solution. The arrays are submerged in water for 2 min at 95° C, transferred into 0.2% SDS for 1 min, rinsed twice with water, air dried and stored in the dark at 25° C.

Cell or tissue mRNA is isolated or commercially obtained and probes are prepared by a single round of reverse transcription. Probes are hybridized to 1 cm<sup>2</sup> microarrays under a 14 x 14 mm glass coverslip for 6-12 hours at 60° C. Arrays are washed for 5 min at 25° C in low stringency wash buffer (1 x SSC/0.2% SDS), then for 10 min at room temperature in high stringency wash buffer (0.1 x SSC/0.2% SDS). Arrays are scanned in 0.1 x SSC using a fluorescence laser scanning device fitted with

a custom filter set. Accurate differential expression measurements are obtained by taking the average of the ratios of two independent hybridizations.

Quantitative analysis of PG1 gene expression may also be performed with full length PG1 cDNAs or fragments thereof in complementary DNA arrays as described by Pietu et al. (Genome Research 6:492-503, 1996). The full length PG1 cDNA or fragments thereof is PCR amplified and spotted on membranes. Then, mRNAs originating from various tissues or cells are labeled with radioactive nucleotides. After hybridization and washing in controlled conditions, the hybridized mRNAs are detected by phospho-imaging or autoradiography. Duplicate experiments are performed and a quantitative analysis of differentially expressed mRNAs is then performed.

Alternatively, expression analysis using the PG1 genomic DNA, the PG1 cDNA, or fragments thereof can be done through high density nucleotide arrays as described by Lockhart et al. (Nature Biotechnology 14: 1675-1680, 1996) and Sosnowsky et al. (Proc. Natl. Acad. Sci. 94:1119-1123, 1997). Oligonucleotides of 15-50 nucleotides from the sequences of the PG1 genomic DNA of SEQ ID NO: 179, the PG1 cDNA of SEQ ID NO: 3, 112-124 or the sequences complementary thereto, are synthesized directly on the chip (Lockhart et al., supra) or synthesized and then addressed to the chip (Sosnowski et al., supra).

PG1 cDNA probes labeled with an appropriate compound, such as biotin, digoxigenin or fluorescent dye, are synthesized from the appropriate mRNA population and then randomly fragmented to an average size of 50 to 100 nucleotides. The said probes are then hybridized to the chip. After washing as described in Lockhart et al., supra and application of different electric fields (Sosnowsky et al., Proc. Natl. Acad. Sci. 94:1119-1123), the dyes or labeling compounds are detected and quantified. Duplicate hybridizations are performed. Comparative analysis of the intensity of the signal originating from cDNA probes on the same target oligonucleotide in different cDNA samples indicates a differential expression of PG1 mRNA.

The above methods may also be used to determine whether an individual exhibits a PG1 expression pattern associated with cancer, prostate cancer or other diseases. In such methods, nucleic acid samples from the individual are assayed for PG1 expression as described above. If a PG1 expression pattern associated with cancer, prostate cancer, or another disease is observed, an appropriate diagnosis is rendered and appropriate therapeutic techniques which target the PG1 gene or protein is applied.

The above methods may also be applied using allele specific probes to determine whether an individual possesses a PG1 allele associated with cancer, prostate cancer, or another disease. In such approaches, one or more allele specific oligonucleotides containing polymorphic nucleotides in the PG1 gene which are associated with prostate cancer are fixed to a microarray. The array is contacted with a nucleic acid sample from the individual being tested under conditions which permit allele specific hybridization of the sample nucleic acid to the allele specific PG1 probes. Hybridization of the sample

nucleic acid to one or more of the allele specific PG1 probes indicates that the individual suffers from prostate cancer caused by the PG1 gene or that the individual is at risk for developing prostate cancer at a subsequent time. Alternatively, any of the genotyping methods described in Section X is utilized..

Use of the Biallelic Markers Of The Invention In Diagnostics

5 The biallelic markers of the present invention can also be used to develop diagnostics tests capable of identifying individuals who express a detectable trait as the result of a specific genotype or individuals whose genotype places them at risk of developing a detectable trait at a subsequent time.

The diagnostic techniques of the present invention may employ a variety of methodologies to determine whether a test subject has a biallelic marker pattern associated with an increased risk of  
10 developing a detectable trait or whether the individual suffers from a detectable trait as a result of a particular mutation, including methods which enable the analysis of individual chromosomes for haplotyping, such as family studies, single sperm DNA analysis or somatic hybrids. The trait analyzed using the present diagnostics is any detectable trait, cancer, prostate cancer or another disease, a response to an anti-cancer, or anti-prostate cancer, or side effects to an anti-cancer or anti-  
15 prostate cancer agent. Diagnostics, which analyze and predict response to a drug or side effects to a drug, is used to determine whether an individual should be treated with a particular drug. For example, if the diagnostic indicates a likelihood that an individual will respond positively to treatment with a particular drug, the drug is administered to the individual. Conversely, if the diagnostic indicates that an individual is likely to respond negatively to treatment with a particular drug, an  
20 alternative course of treatment is prescribed. A negative response is defined as either the absence of an efficacious response or the presence of toxic side effects.

Clinical drug trials represent another application for the markers of the present invention. One or more markers indicative of response to an anti-cancer or anti-prostate cancer agent or to side effects to an anti-cancer or anti-prostate cancer agent is identified using the methods described in  
25 Section XI, below. Thereafter, potential participants in clinical trials of such an agent is screened to identify those individuals most likely to respond favorably to the drug and exclude those likely to experience side effects. In that way, the effectiveness of drug treatment is measured in individuals who respond positively to the drug, without lowering the measurement as a result of the inclusion of individuals who are unlikely to respond positively in the study and without risking undesirable safety  
30 problems. Preferably, in such diagnostic methods, a nucleic acid sample is obtained from the individual and this sample is genotyped using methods described in Section X.

Another aspect of the present invention relates to a method of determining whether an individual is at risk of developing a trait or whether an individual expresses a trait as a consequence of possessing a particular trait-causing allele. The present invention relates to a method of determining  
35 whether an individual is at risk of developing a plurality of traits or whether an individual expresses a plurality of traits as a result of possessing a particular trait-causing allele. These methods involve

obtaining a nucleic acid sample from the individual and determining whether the nucleic acid sample contains one or more alleles of one or more biallelic markers indicative of a risk of developing the trait or indicative that the individual expresses the trait as a result of possessing a particular trait-causing allele.

- 5 As described herein, the diagnostics is based on a single biallelic marker or a group of biallelic markers.

#### **VI. ASSAYING THE PG1 PROTEIN FOR INVOLVEMENT IN RECEPTOR/LIGAND INTERACTIONS**

- 10 The expressed PG1 protein or portion thereof is evaluated for involvement in receptor/ligand interactions as described in Example 16 below.

##### **Example 16**

- 15 The proteins encoded by the PG1 gene or a portion thereof may also be evaluated for their involvement in receptor/ligand interactions. Numerous assays for such involvement are familiar to those skilled in the art, including the assays disclosed in the following references: Chapter 7.28 (Measurement of Cellular Adhesion under Static Conditions 7.28.1-7.28.22) in Current Protocols in Immunology, J.E. Coligan et al. Eds. Greene Publishing Associates and Wiley-Interscience; Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 1987; Bierer et al., J. Exp. Med. 168:1145-1156, 1988; Rosenstein et al., J. Exp. Med. 169:149-160, 1989; Stoltenborg et al., J. Immunol. Methods 175:59-68, 1994; Stitt et al., Cell 80:661-670, 1995; Gyuris et al., Cell 75:791-803, 1993.

- 20 For example, the proteins of the present invention may demonstrate activity as receptors, receptor ligands or inhibitors or agonists of receptor/ligand interactions. Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their  
25 ligands) and receptor/ligand pairs involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses). Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction. A protein of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions.

- 30 The PG1 protein or portions thereof described above is used in drug screening procedures to identify molecules which are agonists, antagonists, or inhibitors of PG1 activity. The PG1 protein or portion thereof used in such analyses is free in solution or linked to a solid support. Alternatively, PG1 protein or portions thereof can be expressed on a cell surface. The cell may naturally express the PG1 protein or portion thereof or, alternatively, the cell may express the PG1 protein or portion thereof from  
35 an expression vector such as those described below.

In one method of drug screening, eukaryotic or prokaryotic host cells which are stably transformed with recombinant polynucleotides in order to express the PG1 protein or a portion thereof are used in conventional competitive binding assays or standard direct binding assays. For example, the formation of a complex between the PG1 protein or a portion thereof and the agent being tested is measured in direct binding assays. Alternatively, the ability of a test agent to prevent formation of a complex between the PG1 protein or a portion thereof and a known ligand is measured.

Alternatively, the high throughput screening techniques disclosed in the published PCT application WO 84/03564, is used. In such techniques, large numbers of small peptides to be tested for PG1 binding activity are synthesized on a surface and affixed thereto. The test peptides are contacted with the PG1 protein or a portion thereof, followed by a wash step. The amount of PG1 protein or portion thereof which binds to the test compound is quantitated using conventional techniques.

In some methods, PG1 protein or a portion thereof is fixed to a surface and contacted with a test compound. After a washing step, the amount of test compound which binds to the PG1 protein or portion thereof is measured.

In another approach, the three dimensional structure of the PG1 protein or a portion thereof may be determined and used for rational drug design.

Alternatively, the PG1 protein or a portion thereof is expressed in a host cell using expression vectors such as those described herein. The PG1 protein or portion thereof is an isotype which is associated with prostate cancer or an isotype which is not associated with prostate cancer. The cells expressing the PG1 protein or portion thereof are contacted with a series of test agents and the effects of the test agents on PG1 activity are measured. Test agents which modify PG1 activity is employed in therapeutic treatments.

The above procedures may also be applied to evaluate mutant PG1 proteins responsible for a detectable phenotype.

#### Identification of Proteins which Interact with the PG1 Protein

Proteins which interact with the PG1 protein is identified as described in Example 17, below.

##### Example 17

Proteins which interact with the PG1 protein or a portion thereof, is identified using two hybrid systems such as the Matchmaker Two Hybrid System 2 (Catalog No. K1604-1, Clontech). As described in the manual accompanying the Matchmaker Two Hybrid System 2 (Catalog No. K1604-1, Clontech), nucleic acids encoding the PG1 protein or a portion thereof, are inserted into an expression vector such that they are in frame with DNA encoding the DNA binding domain of the yeast transcriptional activator GAL4. cDNAs in a cDNA library which encode proteins which might interact with the polypeptides encoded by the nucleic acids encoding the PG1 protein or a portion thereof are inserted into a second expression vector such that they are in frame with DNA encoding the activation domain of GAL4. The two expression plasmids are transformed into yeast and the yeast are plated on selection medium which

selects for expression of selectable markers on each of the expression vectors as well as GAL4 dependent expression of the HIS3 gene. Transformants capable of growing on medium lacking histidine are screened for GAL4 dependent lacZ expression. Those cells which are positive in both the histidine selection and the lacZ assay contain plasmids encoding proteins which interact with the polypeptide  
5 encoded by the nucleic acid inserts.

Alternatively, the system described in Lustig et al., Methods in Enzymology 283: 83-99 (1997), is used for identifying molecules which interact with the PG1 protein or a portion thereof. In such systems, in vitro transcription reactions are performed on vectors containing an insert encoded the PG1 protein or a portion thereof cloned downstream of a promoter which drives in vitro transcription. The  
10 resulting mRNA is introduced into *Xenopus laevis* oocytes. The oocytes are then assayed for a desired activity.

Alternatively, the in vitro transcription products produced as described above is translated in vitro. The in vitro translation products can be assayed for a desired activity or for interaction with a known polypeptide.

15 The system described in U.S. Patent No. 5,654,150 may also be used to identify molecules which interact with the PG1 protein or a portion thereof. In this system, pools of cDNAs are transcribed and translated in vitro and the reaction products are assayed for interaction with a known polypeptide or antibody.

Proteins or other molecules interacting with the PG1 protein or portions thereof can be found  
20 by a variety of additional techniques. In one method, affinity columns containing the PG1 protein or a portion thereof can be constructed. In some versions of this method the affinity column contains chimeric proteins in which the PG1 protein or a portion thereof is fused to glutathione S-transferase. A mixture of cellular proteins or pool of expressed proteins as described above is applied to the affinity column. Proteins interacting with the polypeptide attached to the column can then be isolated  
25 and analyzed on 2-D electrophoresis gel as described in Ramunsen et al. Electrophoresis, 18, 588-598 (1997). Alternatively, the proteins retained on the affinity column can be purified by electrophoresis based methods and sequenced. The same method can be used to isolate antibodies, to screen phage display products, or to screen phage display human antibodies.

Proteins interacting with the PG1 protein or portions thereof can also be screened by using an  
30 Optical Biosensor as described in Edwards et Leatherbarrow, Analytical Biochemistry, 246, 1-6 (1997). The main advantage of the method is that it allows the determination of the association rate between the protein and other interacting molecules. Thus, it is possible to specifically select interacting molecules with a high or low association rate. Typically a target molecule is linked to the sensor surface (through a carboxymethyl dextran matrix) and a sample of test molecules is placed in  
35 contact with the target molecules. The binding of a test molecule to the target molecule causes a change in the refractive index and/ or thickness. This change is detected by the Biosensor provided it

occurs in the evanescent field (which extend a few hundred nanometers from the sensor surface). In these screening assays, the target molecule can be the PG1 protein or a portion thereof and the test sample can be a collection of proteins extracted from tissues or cells, a pool of expressed proteins, combinatorial peptide and/ or chemical libraries, or phage displayed peptides. The tissues or cells from which the test proteins are extracted can originate from any species.

In other methods, a target protein is immobilized and the test population is the PG1 protein or a portion thereof.

To study the interaction of the PG1 protein or a portion thereof with drugs, the microdialysis coupled to HPLC method described by Wang et al., *Chromatographia*, 44, 205-208(1997) or the affinity capillary electrophoresis method described by Busch et al., *J. Chromatogr.* 777:311-328 (1997).

The above procedures may also be applied to evaluate mutant PG1 proteins responsible for a detectable phenotype.

#### **VII. PRODUCTION OF ANTIBODIES AGAINST PG1 POLYPEPTIDES**

Any PG1 polypeptide or whole protein (SEQ ID NOs: 4, 5, 70, 74, 125-136) whether human, mouse or mammalian is used to generate antibodies capable of specifically binding to expressed PG1 protein or fragments thereof as described in Example 16, below. The antibodies is capable of binding the full length PG1 protein. PG1 proteins which result from naturally occurring mutant, particularly functional mutants of PG1, including SEQ ID NO: 70, which may used in the production of antibodies. The present invention also contemplates the use of polypeptides comprising a contiguous stretch of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 50, or 100 amino acids of any PG1 protein in the manufacture of antibodies. In a preferred embodiment the contiguous stretch of amino acids comprises the site of a mutation or functional mutation, including a deletion, addition, swap or truncation of the amino acids in the PG1 protein sequence. For instance, polypeptides that contain either the Arg and His residues at amino acid position 184, and polypeptides that contain either the Arg or Ile residue at amino acid position 293 of the SEQ ID NO: 4 in said contiguous stretch are particularly preferred embodiments of the invention and useful in the manufacture of antibodies to detect the presence and absence of these mutations. Similarly, polypeptides with a carboxy terminus at position 228 is a particularly preferred embodiment of the invention and useful in the manufacture of antibodies to detect the presence and absence of the mutation shown in SEQ ID NOs: 69 and 70. Similarly, polypeptides that that contain an peptide sequences of 8, 10, 12, 15, or 25 amino acids encoded over a naturally-occurring splice junction (the point at which two human PG1 exon (SEQ ID NOs: 100-111) are covalently linked) in said contiguous stretch are particularly preferred embodiments and useful in the manufacture of antibodies to detect the presence, localization, and quantity of the various protein products of the PG1 alternative splice species.



Alternatively, the antibodies is screened so as to isolate those which are capable of binding an epitope-containing fragment of at least 8, 10, 12, 15, 20, 25, or 30 amino acids of a human, mouse, or mammalian PG1 protein, preferably a sequence selected from SEQ ID NOs: 4, 5, 70, 74, or 125-136.

Antibodies may also be generated which are capable of specifically binding to a given isoform of the PG1 protein. For example, the antibodies is capable of specifically binding to an isoform of the PG1 protein which causes prostate cancer or another detectable phenotype which has been obtained as described above and expressed from an expression vector as described above. Alternatively, the antibodies is capable of binding to an isoform of the PG1 protein which does not cause prostate cancer. Such antibodies is used in diagnostic assays in which protein samples from an individual are evaluated for the presence of an isoform of the PG1 protein which causes cancer or another detectable phenotype using techniques such as Western blotting or ELISA assays.

Non-human animals or mammals, whether wild-type or transgenic, which express a different species of PG1 than the one to which antibody binding is desired, and animals which do not express PG1 (i.e. an PG1 knock out animal as described in Section VIII.) are particularly useful for preparing antibodies. PG1 knock out animals will recognize all or most of the exposed regions of PG1 as foreign antigens, and therefore produce antibodies with a wider array of PG1 epitopes. The humoral immune system of animals which produce a species of PG1 that resembles the antigenic sequence will preferentially recognize the differences between the animal's native PG1 species and the antigen sequence, and produce antibodies to these unique sites in the antigen sequence.

#### Example 18

Substantially pure protein or polypeptide is isolated from transfected or transformed cells containing an expression vector encoding the PG1 protein or a portion thereof as described in Example 11. The concentration of protein in the final preparation is adjusted, for example, by concentration on an Amicon filter device, to the level of a few micrograms/ml. Monoclonal or polyclonal antibody to the protein can then be prepared as follows:

##### A. Monoclonal Antibody Production by Hybridoma Fusion

Monoclonal antibody to epitopes in the PG1 protein or a portion thereof can be prepared from murine hybridomas according to the classical method of Kohler, G. and Milstein, C., Nature 256:495 (1975) or derivative methods thereof. Also see Harlow, E., and D. Lane. 1988. Antibodies A Laboratory Manual. Cold Spring Harbor Laboratory. pp. 53-242.

Briefly, a mouse is repetitively inoculated with a few micrograms of the PG1 protein or a portion thereof over a period of a few weeks. The mouse is then sacrificed, and the antibody producing cells of the spleen isolated. The spleen cells are fused by means of polyethylene glycol with mouse myeloma cells, and the excess unfused cells destroyed by growth of the system on selective media comprising aminopterin (HAT media). The successfully fused cells are diluted and aliquots of the dilution placed in wells of a microtiter plate where growth of the culture is continued. Antibody-producing clones are

identified by detection of antibody in the supernatant fluid of the wells by immunoassay procedures, such as ELISA, as originally described by Engvall, E., Meth. Enzymol. 70:419 (1980), and derivative methods thereof. Selected positive clones can be expanded and their monoclonal antibody product harvested for use. Detailed procedures for monoclonal antibody production are described in Davis, L. et al. Basic Methods in Molecular Biology Elsevier, New York. Section 21-2.

#### B. Polyclonal Antibody Production by Immunization

Polyclonal antiserum containing antibodies to heterogeneous epitopes in the PG1 protein or a portion thereof can be prepared by immunizing suitable non-human animal with the PG1 protein or a portion thereof, which can be unmodified or modified to enhance immunogenicity. A suitable non-human animal is preferably a non-human mammal is selected, usually a mouse, rat, rabbit, goat, or horse. Alternatively, a crude preparation which has been enriched for PG1 concentration can be used to generate antibodies. Such proteins, fragments or preparations are introduced into the non-human mammal in the presence of an appropriate adjuvant (e.g. aluminum hydroxide, RIBI, etc.) which is known in the art. In addition the protein, fragment or preparation can be pretreated with an agent which will increase antigenicity, such agents are known in the art and include, for example, methylated bovine serum albumin (mBSA), bovine serum albumin (BSA), Hepatitis B surface antigen, and keyhole limpet hemocyanin (KLH). Serum from the immunized animal is collected, treated and tested according to known procedures. If the serum contains polyclonal antibodies to undesired epitopes, the polyclonal antibodies can be purified by immunoaffinity chromatography.

Effective polyclonal antibody production is affected by many factors related both to the antigen and the host species. Also, host animals vary in response to site of inoculations and dose, with both inadequate or excessive doses of antigen resulting in low titer antisera. Small doses (ng level) of antigen administered at multiple intradermal sites appears to be most reliable. Techniques for producing and processing polyclonal antisera are known in the art, see for example, Mayer and Walker (1987). An effective immunization protocol for rabbits can be found in Vaitukaitis, J. et al. J. Clin. Endocrinol. Metab. 33:988-991 (1971).

Booster injections can be given at regular intervals, and antiserum harvested when antibody titer thereof, as determined semi-quantitatively, for example, by double immunodiffusion in agar against known concentrations of the antigen, begins to fall. See, for example, Ouchterlony, O. et al., Chap. 19 in: Handbook of Experimental Immunology D. Wier (ed) Blackwell (1973). Plateau concentration of antibody is usually in the range of 0.1 to 0.2 mg/ml of serum (about  $12 \cdot M$ ). Affinity of the antisera for the antigen is determined by preparing competitive binding curves, as described, for example, by Fisher, D., Chap. 42 in: Manual of Clinical Immunology, 2d Ed. (Rose and Friedman, Eds.) Amer. Soc. For Microbiol., Washington, D.C. (1980).

Antibody preparations prepared according to either the monoclonal or the polyclonal protocol are useful in quantitative immunoassays which determine concentrations of antigen-bearing substances

in biological samples; they are also used semi-quantitatively or qualitatively to identify the presence of antigen in a biological sample. The antibodies may also be used in therapeutic compositions for killing cells expressing the protein or reducing the levels of the protein in the body.

#### **VIII. VECTORS AND THE USES OF POLYNUCLEOTIDES IN CELLS, ANIMALS, AND**

##### **5 HUMANS**

The nucleic acids of the invention include expression vectors, amplification vectors, PCR-suitable polynucleotide primers, and vectors which are suitable for the introduction of a polynucleotide of the invention into an embryonic stem cells for the production of transgenic non-human animals. In addition, vectors which are suitable for the introduction of a polynucleotide of the invention into cells, organs and individuals, including human individuals, for the purposes of gene therapy to reduce the severity of or prevent genetic diseases associated with functional mutations in PG1 genes are encompassed by the present invention. Functional mutations in PG1 genes which are suitable as targets for the gene therapy and transgenic vectors and methods of the invention include, but are not limited to, mutations in the coding region of the PG1 gene which affect the amino acid sequence of the PG1 gene's product, mutations in the promoter or other regulatory regions which affect the levels of PG1 expression, mutations in the PG1 splice sites which affect length of the PG1 gene product or the relative frequency of PG1 alternative splicing species, and any other mutation which in any way affects the level or quality of PG1 expression or activity. The gene therapy methods can be achieved by targeting vectors and method for changing a mutant PG1 gene into a wild-type PG1 gene in a embryonic stem cell or somatic cell. Alternatively, the present invention also encompasses methods and vectors for introducing the expression of wild-type PG1 sequences without the disruption of any mutant PG1 which already reside in the cell, organ or individual.

The invention also embodies amplification vectors, which comprise a polynucleotide of the invention, and an origin of replication. Preferably, such amplification vectors further comprise restriction endonuclease sites flanking the polynucleotide, so as to facilitate cleavage and purification of the polynucleotides from the remainder of the amplification vector, and a selectable marker, so as to facilitate amplification of the amplification vector. Most preferably, the restriction endonuclease sites in the amplification vector are situated such that cleavage at those site would result in no other amplification vector fragments of a similar size.

Thus, such an amplification vector is transfected into a host cell compatible with the origin of replication of said amplification vector, wherein the host cell is a prokaryotic or eukaryotic cell, preferably a mammalian, insect, yeast, or bacterial cell, most preferably an Escherichia coli cell. The resulting transfected host cells is grown by culture methods known in the art, preferably under selection compatible with the selectable marker (e.g., antibiotics). The amplification vectors can be isolated and purified by methods known in the art (e.g., standard plasmid prep procedures). The polynucleotide of the invention can be cleaved with restriction enzymes that specifically cleave at the

restriction endonuclease sites flanking the polynucleotide, and the double-stranded polynucleotide fragment purified by techniques known in the art, including gel electrophoresis.

Alternatively linear polynucleotides comprising a polynucleotide of the invention is amplified by PCR. The PCR method is well known in the art and described in, e.g., U.S. Patent Nos. 4,683,195  
5 and 4,683,202 and Saiki, R et al. 1988. Science 239:487-491, and European patent applications 86302298.4, 86302299.2 and 87300203.4, as well as Methods in Enzymology 1987 155:335-350.

The polynucleotides of the invention can also be derivatized in various ways, including those appropriate for facilitating transfection and/or gene therapy. The polynucleotides can be derivatized by attaching a nuclear localization signal to it to improve targeted delivery to the nucleus. One well-  
10 characterized nuclear localization signal is the heptapeptide PKKKRKV (pro-lys-lys-lys-arg-lys-val). Preferably, in the case of polynucleotides in the form of a closed circle, the nuclear localization signal is attached via a modified loop nucleotide or spacer that forms a branching structure.

If it is to be used in vivo, the polynucleotide of the invention is derivatized to include ligands and/or delivery vehicles which provide dispersion through the blood, targeting to specific cell types,  
15 or permit easier transit of cellular barriers. Thus, the polynucleotides of the invention is linked or combined with any targeting or delivery agent known in the art, including but not limited to, cell penetration enhancers, lipofectin, liposomes, dendrimers, DNA intercalators, and nanoparticles. In particular, nanoparticles for use in the delivery of the polynucleotides of the invention are particles of less than about 50 nanometers diameter, nontoxic, non-antigenic, and comprised of albumin and  
20 surfactant, or iron as in the nanoparticle particle technology of SynGenix. In general the delivery vehicles used to target the polynucleotides of the invention may further comprise any cell specific or general targeting agents known in the art, and will have a specific trapping efficiency to the target cells or organs of from about 5 to about 35%.

The polynucleotides of the invention is used ex vivo in a gene therapy method for obtaining  
25 cells or organs which produce wild-type PG1 or PG1 proteins which have been selectively mutated. The cells are created by incubation of the target cell with one or more of the above-described polynucleotides under standard conditions for uptake of nucleic acids, including electroporation or lipofection. In practicing an ex vivo method of treating cells or organs, the concentration of polynucleotides of the invention in a solution prepare to treat target cells or organs is from about 0.1  
30 to about 100  $\mu$ M, preferably 0.5 to 50  $\mu$ M, most preferably from 1 to 10  $\mu$ M.

Alternatively, the oligonucleotides can be modified or co-administered for targeted delivery to the nucleus. Improved oligonucleotide stability is expected in the nucleus due to: (1) lower levels of DNases and RNases; and (2) higher oligonucleotide concentrations due to lower total volume.

Alternatively, the polynucleotides of the invention can be covalently bonded to biotin to form  
35 a biotin-polynucleotide prodrug by methods known in the art, and co-administered with a receptor ligand bound to avidin or receptor specific antibody bound to avidin, wherein the receptor is capable

of causing uptake of the resulting polynucleotide-biotin-avidin complex into the cells. Receptors that cause uptake are known to those of skill in the art.

The invention encompasses vectors which are suitable for the introduction of a polynucleotide of the invention into an embryonic stem cell for the production of transgenic non-human animals, which in turn result in the expression of recombinant PG1 in the transgenic animal. Any appropriate vector system can be used for the introduction and expression of PG1 in transgenic animals, including for example yeast artificial chromosomes (YAC), bacterial artificial chromosomes (BAC), bacteriophage P1, and other vectors known in the art which are able to accommodate sufficiently large inserts to encode the PG1 protein or desired fragments thereof. Selected alterations, additions and deletions in the PG1 gene may optionally be achieved by site-directed mutagenesis. Once an appropriate vector system is chosen, the site-directed mutagenesis process may then be conducted by techniques well known in the art, and the fragment be returned and ligated to the larger vector from which it was cleaved. For site directed mutagenesis methods see, for example, Kunkel, T. 1985. *Proc. Natl. Acad. Sci. U.S.A.* 82:488; Bandeyar, M. et al. 1988. *Gene* 65: 129-133; Nelson, M., and M. McClelland 1992. *Methods Enzymol.* 216:279-303; Weiner, M. 1994. *Gene* 151: 119-123; Costa, G. and M. Weiner. 1994. *Nucleic Acids Res.* 22: 2423; Hu, G. 1993. *DNA and Cell Biology* 12:763-770; and Deng, W. and J. Nickoff. 1992. *Anal. Biochem.* 200:81.

Briefly, the transgenic technology used herein involves the inactivation, addition or replacement of a portion of the PG1 gene or the entire gene. For example the present technology includes the addition of PG1 genes with or without the inactivation of the non-human animal's native PG1 genes, as described in the preceding two paragraphs and in the Examples. The invention also encompasses the use of vectors, and the vectors themselves which target and modify an existing human PG1 gene in a stem cell, whether it is contained in a non-human animal cell where it was previously introduced into the germ line by transgenic technology or it is a native PG1 gene in a human pluripotent or somatic cell. This transgene technology usually relies on homologous recombination in a pluripotent cell that is capable of differentiating into germ cell tissue. A DNA construct that encodes an altered region of the non-human animal's PG1 gene that contains, for instance a stop codon to destroy expression, is introduced into the nuclei of embryonic stem cells. Preferably mice are used for this transgenic work. In a portion of the cells, the introduced DNA recombines with the endogenous copy of the cell's gene, replacing it with the altered copy. Cells containing the newly engineered genetic alteration are injected in a host embryo of the same species as the stem cell, and the embryo is reimplanted into a recipient female. Some of these embryos develop into chimeric individuals that possess germ cells entirely derived from the mutant cell line. Therefore, by breeding the chimeric progeny it is possible to obtain a new strain containing the introduced genetic alteration. See Capecchi 1989. *Science*. 244:1288-1292 for a review of this procedure.

The present invention encompasses the polynucleotides described herein, as well as the methods for making these polynucleotides including the method for creating a mutation in a human PG1 gene. In addition, the present invention encompasses cells which comprise the polynucleotides of the invention, including but not limited to amplification host cells comprising amplification vectors of the invention. Furthermore the present invention comprises the embryonic stem cells and transgenic non-human animals and mammals described herein which comprise a gene encoding a human PG1 protein.

DNA construct that enables directing temporal and spatial gene expression in recombinant host cells and in transgenic animals

10 In order to study the physiological and phenotype consequences of a lack of synthesis of the PG1 protein, both at the cellular level and at the multi-cellular organism level, in particular as regards to disorders related to abnormal cell proliferation, notably cancers, the invention also encompasses DNA constructs and recombinant vectors enabling a conditional expression of a specific allele of the PG1 genomic sequence or cDNA and also of a copy of this genomic sequence or cDNA harboring  
15 substitutions, deletions, or additions of one or more bases as regards to the PG1 nucleotide sequence of SEQ ID NOs: 3, 112-125, 179, 182-184, or a fragment thereof, these base substitutions, deletions or additions being located either in an exon, an intron or a regulatory sequence, but preferably in a 5'-regulatory sequence of a mammalian PG1 gene, more preferably SEQ ID NO: 180 or in an exon of the PG1 genomic sequence or within the PG1 cDNA of SEQ ID NOs 3, 112-125, or 184.

20 A first preferred DNA construct is based on the tetracycline resistance operon tet from E. coli transposon Tn110 for controlling the PG1 gene expression, such as described by Gossen M. et al., 1992, Proc. Natl. Acad. Sci. USA, 89: 5547-5551; Gossen M. et al., 1995, Science, 268: 1766-1769; and Furth P.A. et al., 1994, Proc. Natl. Acad. Sci. USA, 91: 9302-9306. Such a DNA construct contains seven tet operator sequences from Tn10 (tetop) that are fused to either a minimal promoter or  
25 a 5'-regulatory sequence of the PG1 gene, said minimal promoter or said PG1 regulatory sequence being operably linked to a polynucleotide of interest that codes either for a sense or an antisense oligonucleotide or for a polypeptide, including a PG1 polypeptide or a peptide fragment thereof. This DNA construct is functional as a conditional expression system for the nucleotide sequence of interest when the same cell also comprises a nucleotide sequence coding for either the wild type (tTA) or the  
30 mutant (rTA) repressor fused to the activating domain of viral protein VP16 of herpes simplex virus, placed under the control of a promoter, such as the HCMVIE1 enhancer/promoter or the MMTV-LTR. Indeed, a preferred DNA construct of the invention will comprise both the polynucleotide containing the tet operator sequences and the polynucleotide containing a sequence coding for the tTA or the rTA repressor.

In the specific embodiment wherein the conditional expression DNA construct contains the sequence encoding the mutant tetracycline repressor rTA, the expression of the polynucleotide of interest is silent in the absence of tetracycline and induced in its presence.

DNA constructs allowing homologous recombination : replacement vectors

5 A second preferred DNA construct will comprise, from 5'-end to 3'-end : (a) a first nucleotide sequence that is comprised of a PG1 sequence preferably a PG1 genomic sequence; (b) a nucleotide sequence comprising a positive selection marker, such as the marker for neomycin resistance (neo); and (c) a second nucleotide sequence that comprised of a PG1 sequence preferably a PG1 genomic sequence, and is located on the genome downstream the first PG1 nucleotide sequence (a).

10 In a preferred embodiment, this DNA construct also comprises a negative selection marker located upstream the nucleotide sequence (a) or downstream the nucleotide sequence (b). Preferably, the negative selection marker consists of the thymidine kinase (tk) gene (Thomas K.R. et al., 1986, Cell, 44: 419-428), the hygromycin beta gene (Te Riele et al., 1990, Nature, 348: 649-651), the hprt gene (Van der Lugt et al., 1991, Gene, 105: 263-267; and Reid L.H. et al., 1990, Proc. Natl. Acad. Sci. USA, 87: 4299-4303) or the Diphtheria toxin A fragment (Dt-A) gene (Nada S. et al., 1993, Cell, 73: 1125-1135; Yagi T. et al., 1990, Proc. Natl. Acad. Sci. USA, 87: 9918-9922). Preferably, the positive selection marker is located within a PG1 exon sequence so as to interrupt the sequence encoding a PG1 protein.

20 These replacement vectors are described for example by Thomas K.R. et al., 1986, Cell, 44: 419-428; Thomas K.R. et al., 1987, Cell, 51: 503-512; Mansour S.L. et al., 1988, Nature, 336: 348-352; and Koller et al., 1992, Annu. Rev. Immunol., 10: 705-30.

The first and second nucleotide sequences (a) and (c) is located at any point within a PG1 regulatory sequence, an intronic sequence, an exon sequence or a sequence containing both regulatory and/or intronic and/or exon sequences. The length of nucleotide sequences (a) and (c) is determined empirically by one of ordinary skill in the art. Nucleotide sequences (a) and (c) or any length are specifically contemplated in the present invention, however, lengths ranging from 1 kb to 50 kb, preferably from 1 kb to 10 kb, more preferably from 2 kb to 6 kb and most preferably from 2 kb to 4 kb are normally used.

DNA constructs allowing homologous recombination : Cre-loxP system.

30 These new DNA constructs make use of the site-specific recombination system of the P1 phage. The P1 phage possesses a recombinase called Cre which interacts specifically with a 34 base pairs loxP site. The loxP site is composed of two palindromic sequences of 13 bp separated by a 8 bp conserved sequence (Hoess et al., 1986, Nucleic Acids Res., 14: 2287-2300). The recombination by the Cre enzyme between two loxP sites having an identical orientation leads to the deletion of the DNA fragment.

35

The Cre-loxP system used in combination with a homologous recombination technique has been first described by Gu H. et al., 1993, *Cell*, 73: 1155-1164 ; and Gu H. et al., 1994, *Science*, 265: 103-106. Briefly, a nucleotide sequence of interest to be inserted in a targeted location of the genome harbors at least two loxP sites in the same orientation and located at the respective ends of a nucleotide sequence to be excised from the recombinant genome. The excision event requires the presence of the recombinase (Cre) enzyme within the nucleus of the recombinant host cell. The recombinase enzyme is brought at the desired time either by (a) incubating the recombinant host cells in a culture medium containing this enzyme, by injecting the Cre enzyme directly into the desired cell, such as described by Araki K. et al., 1995, *Proc. Natl. Acad. Sci. USA*, 92: 160-164 ; or by lipofection of the enzyme into the cells, such as described by Baubonis et al., 1993, *Nucleic Acids Res.*, 21: 2025-2029; (b) transfecting the cell host with a vector comprising the Cre coding sequence operably linked to a promoter functional in the recombinant cell host, which promoter being optionally inducible, said vector being introduced in the recombinant cell host, such as described by Gu H. et al., 1993, *Cell*, 73: 1155-1164; and Sauer B. et al., 1988, *Proc. Natl. Acad. Sci. USA*, 85: 5166-5170; (c) introducing in the genome of the host cell a polynucleotide comprising the Cre coding sequence operably linked to a promoter functional in the recombinant cell host, which promoter is optionally inducible, and said polynucleotide being inserted in the genome of the cell host either by a random insertion event or an homologous recombination event, such as described by Gu H. et al., 1994, *Science*, 265: 103-106.

In the specific embodiment wherein the vector containing the sequence to be inserted in the PG1 gene by homologous recombination is constructed in such a way that selectable markers are flanked by loxP sites of the same orientation, it is possible, by treatment by the Cre enzyme, to eliminate the selectable markers while leaving the PG1 sequences of interest that have been inserted by an homologous recombination event. Again, two selectable markers are needed: a positive selection marker to select for the recombination event and a negative selection marker to select for the homologous recombination event. Vectors and methods using the Cre-loxP system are described by Zou Y.R. et al., 1994, *Curr. Biol.*, 4: 1099-1103.

Thus, a third preferred DNA construct of the invention comprises, from 5'-end to 3'-end: (a) a first nucleotide sequence that is comprised of a PG1 sequence, preferably a PG1 genomic sequence; (b) a nucleotide sequence comprising a polynucleotide encoding a positive selection marker, such as the marker for neomycin resistance (neo), said nucleotide sequence comprising additionally two sequences defining a site recognized by a recombinase, such as a loxP site, the two sites being placed in the same orientation; and (c) a second nucleotide sequence that is comprised of a PG1 sequence, preferably a PG1 genomic sequence, and is located on the genome downstream of the first PG1 nucleotide sequence (a).

The sequences defining a site recognized by a recombinase, such as a loxP site, are preferably located within the nucleotide sequence (b) at suitable locations bordering the nucleotide sequence for



which the conditional excision is sought. In one specific embodiment, two loxP sites are located at each side of the positive selection marker sequence, in order to allow its excision at a desired time after the occurrence of the homologous recombination event.

In a preferred embodiment of a method using the third DNA construct described above, the excision of the polynucleotide fragment bordered by the two sites recognized by a recombinase, preferably two loxP sites, is performed at a desired time, due to the presence within the genome of the recombinant host cell of a sequence encoding the Cre enzyme operably linked to a promoter sequence, preferably an inducible promoter, more preferably a tissue-specific promoter sequence and most preferably a promoter sequence which is both inducible and tissue-specific, such as described by Gu H. et al., 1994, Science, 265: 103-106.

The presence of the Cre enzyme within the genome of the recombinant cell host may result of the breeding of two transgenic animals, the first transgenic animal bearing the PG1-derived sequence of interest containing the loxP sites as described above and the second transgenic animal bearing the Cre coding sequence operably linked to a suitable promoter sequence, such as described by Gu H. et al., 1994, Science, 265: 103-106. Spatio-temporal control of the Cre enzyme expression may also be achieved with an adenovirus based vector that contains the Cre gene thus allowing infection of cells, or in vivo infection of organs, for delivery of the Cre enzyme, such as described by Anton M. et al., 1995, J. Virol., 69: 4600-4606; and Kanegae Y. et al., 1995, Nucl. Acids Res., 23: 3816-3821.

The DNA constructs described above is used to introduce a desired nucleotide sequence of the invention, preferably a PG1 genomic sequence or a PG1 cDNA sequence, and most preferably an altered copy of a PG1 genomic or cDNA sequence, within a predetermined location of the targeted genome, leading either to the generation of an altered copy of a targeted gene (knock-out homologous recombination) or to the replacement of a copy of the targeted gene by another copy sufficiently homologous to allow an homologous recombination event to occur (knock-in homologous recombination).

#### Nuclear antisense DNA constructs

Preferably, the antisense polynucleotides of the invention have a 3' polyadenylation signal that has been replaced with a self-cleaving ribozyme sequence, such that RNA polymerase II transcripts are produced without poly(A) at their 3' ends, these antisense polynucleotides being incapable of export from the nucleus, such as described by Liu Z. et al., 1994, Proc. Natl. Acad. Sci. USA, 91: 4528-4262. In a preferred embodiment, these PG1 antisense polynucleotides also comprise, within the ribozyme cassette, a histone stem-loop structure to stabilize cleaved transcripts against 3'-5' exonucleolytic degradation, such as described by Eckner R. et al., 1991, EMBO J., 10: 3513-3522.

#### Expression Vectors

The polynucleotides of the invention also include expression vectors. Expression vector systems, control sequences and compatible host are known in the art. For a review of these systems

see, for example, U.S. Patent No. 5,350,671, columns 45-48. Any of the standard methods known to those skilled in the art for the insertion of DNA fragments into a vector is used to construct expression vectors containing a chimeric gene consisting of appropriate transcriptional/translational control signals and the protein coding sequences. These methods may include in vitro recombinant DNA and  
5 synthetic techniques and in vivo recombinants (genetic recombination).

Expression of a polypeptide, peptide or derivative, or analogs thereof encoded by a polynucleotide sequence in SEQ ID NOs: 3, 69, 100-112, or 179-184 is regulated by a second nucleic acid sequence so that the protein or peptide is expressed in a host transformed with the recombinant DNA molecule. For example, expression of a protein or peptide is controlled by any  
10 promoter/enhancer element known in the art. Promoters which is used to control expression include, but are not limited to, the CMV promoter, the SV40 early promoter region (Bernoist and Chambon, 1981, *Nature* 290:304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, et al., 1980, *Cell* 22:787-797), the herpes thymidine kinase promoter (Wagner et al., 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:1441-1445), the regulatory sequences of the  
15 metallothionein gene (Brinster et al., 1982, *Nature* 296:39-42); prokaryotic expression vectors such as the beta-lactamase promoter (Villa-Kamaroff, et al., 1978, *Proc. Natl. Acad. Sci. U.S.A.* 75:3727-3731), or the tac promoter (DeBoer, et al., 1983, *Proc. Natl. Acad. Sci. U.S.A.* 80:21-25); see also "Useful proteins from recombinant bacteria" in *Scientific American*, 1980, 242:74-94; plant expression vectors comprising the nopaline synthetase promoter region (Herrera-Estrella et al., 1983,  
20 *Nature* 303:209-213) or the cauliflower mosaic virus 35S RNA promoter (Gardner, et al., 1981, *Nucl. Acids Res.* 9:2871), and the promoter of the photosynthetic enzyme ribulose biphosphate carboxylase (Herrera-Estrella et al., 1984, *Nature* 310:115-120); promoter elements from yeast or other fungi such as the Gal 4 promoter, the ADC (alcohol dehydrogenase) promoter, PGK (phosphoglycerol kinase) promoter, alkaline phosphatase promoter, and the following animal transcriptional control regions,  
25 which exhibit tissue specificity and have been utilized in transgenic animals: elastase I gene control region which is active in pancreatic acinar cells (Swift et al., 1984, *Cell* 38:639-646; Ornitz et al., 1986, *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409; MacDonald, 1987, *Hepatology* 7:425-515); insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, *Nature* 315:115-122), immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., 1984, *Cell* 38:647-658; Adames et al., 1985, *Nature* 318:533-538; Alexander et al., 1987, *Mol. Cell. Biol.* 7:1436-1444), mouse mammary tumor virus control region which is active in testicular,  
30 breast, lymphoid and mast cells (Leder et al., 1986, *Cell* 45:485-495), albumin gene control region which is active in liver (Pinkert et al., 1987, *Genes and Devel.* 1:268-276), alpha-fetoprotein gene control region which is active in liver (Krumlauf et al., 1985, *Mol. Cell. Biol.* 5:1639-1648; Hammer et al., 1987, *Science* 235:53-58; alpha 1-antitrypsin gene control region which is active in the liver  
35 (Kelsey et al., 1987, *Genes and Devel.* 1:161-171), beta-globin gene control region which is active in

myeloid cells (Mogram et al., 1985, Nature 315:338-340; Kollias et al., 1986, Cell 46:89-94; myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead et al., 1987, Cell 48:703-712); myosin light chain-2 gene control region which is active in skeletal muscle (Sani, 1985, Nature 314:283-286), and gonadotropic releasing hormone gene control region which is active in the hypothalamus (Mason et al., 1986, Science 234:1372-1378).

Other suitable vectors, particularly for the expression of genes in mammalian cells, is selected from the group of vectors consisting of P1 bacteriophages, and bacterial artificial chromosomes (BACs). These types of vectors may contain large inserts ranging from about 80-90 kb (P1 bacteriophage) to about 300 kb (BACs).

10

#### P1 bacteriophage

The construction of P1 bacteriophage vectors such as p158 or p158/neo8 are notably described by Sternberg N.L., 1992, Trends Genet., 8: 1-16; and Sternberg N.L., 1994, Mamm. Genome, 5: 397-404. Recombinant P1 clones comprising PG1 nucleotide sequences is designed for inserting large polynucleotides of more than 40 kb (Linton M.F. et al., 1993, J. Clin. Invest., 92: 3029-3037). To generate P1 DNA for transgenic experiments, a preferred protocol is the protocol described by McCormick et al., 1994, Genet. Anal. Tech. Appl., 11: 158-164. Briefly, E. coli (preferably strain NS3529) harboring the P1 plasmid are grown overnight in a suitable broth medium containing 25 µg/ml of kanamycin. The P1 DNA is prepared from the E. coli by alkaline lysis using the Qiagen Plasmid Maxi kit (Qiagen, Chatsworth, CA, USA), according to the manufacturer's instructions. The P1 DNA is purified from the bacterial lysate on two Qiagen-tip 500 columns, using the washing and elution buffers contained in the kit. A phenol/chloroform extraction is then performed before precipitating the DNA with 70% ethanol. After solubilizing the DNA in TE (10 mM Tris-HCl, pH 7.4, 1 mM EDTA), the concentration of the DNA is assessed by spectrophotometry.

When the goal is to express a P1 clone comprising PG1 nucleotide sequences in a transgenic animal, typically in transgenic mice, it is desirable to remove vector sequences from the P1 DNA fragment, for example by cleaving the P1 DNA at rare-cutting sites within the P1 polylinker (SfiI, NotI or SalI). The P1 insert is then purified from vector sequences on a pulsed-field agarose gel, using methods similar to those originally reported for the isolation of DNA from YACs (Schedl A. et al., 1993, Nature, 362: 258-261; and Peterson et al., 1993, Proc. Natl. Acad. Sci. USA, 90: 7593-7597). At this stage, the resulting purified insert DNA can be concentrated, if necessary, on a Millipore Ultrafree-MC Filter Unit (Millipore, Bedford, MA, USA – 30,000 molecular weight limit) and then dialyzed against microinjection buffer (10 mM Tris-HCl, pH 7.4; 250 µM EDTA) containing 100 mM NaCl, 30 µM spermine, 70 µM spermidine on a microdialysis membrane (type VS, 0.025 µM from Millipore). The intactness of the purified P1 DNA insert is assessed by electrophoresis on 1% agarose (Sea Kem GTG; FMC Bio-products) pulse-field gel and staining with ethidium bromide.

### Bacterial Artificial Chromosomes (BACs)

The bacterial artificial chromosome (BAC) cloning system (Shizuya et al., 1992, Proc. Natl. Acad. Sci. USA, 89: 8794-8797) has been developed to stably maintain large fragments of genomic DNA (100-300 kb) in *E. coli*. A preferred BAC vector consists of pBeloBAC11 vector that has been described Kim U. J., et al., 1996, Genomics, 34: 213-218. BAC libraries are prepared with this vector using size-selected genomic DNA that has been partially digested using enzymes that permit ligation into either the Bam HI or Hind III sites in the vector. Flanking these cloning sites are T7 and SP6 RNA polymerase transcription initiation sites that can be used to generate end probes by either RNA transcription or PCR methods. After the construction of a BAC library in *E. coli*, BAC DNA is purified from the host cell as a supercoiled circle. Converting these circular molecules into a linear form precedes both size determination and introduction of the BACs into recipient cells. The cloning site is flanked by two Not I sites, permitting cloned segments to be excised from the vector by Not I digestion. Alternatively, the DNA insert contained in the pBeloBAC11 vector is linearized by treatment of the BAC vector with the commercially available enzyme lambda terminase that leads to the cleavage at the unique cosN site, but this cleavage method results in a full length BAC clone containing both the insert DNA and the BAC sequences.

### Host Cells

The PG1 gene expression in human cells is rendered defective, or alternatively it is proceeded with the insertion of a PG1 genomic or cDNA sequence with the replacement of the PG1 gene counterpart in the genome of an animal cell by a PG1 polynucleotide according to the invention. These genetic alterations is generated by homologous recombination events using specific DNA constructs that have been previously described.

One kind of host cell that is used are mammal zygotes, such as murine zygotes. For example, murine zygotes may undergo microinjection with a purified DNA molecule of interest, for example a purified DNA molecule that has previously been adjusted to a concentration range from 1 ng/ml –for BAC inserts- 3 ng/μl –for P1 bacteriophage inserts- in 10 mM Tris-HCl, pH 7.4, 250 μM EDTA containing 100 mM NaCl, 30 μM spermine, and 70 μM spermidine. When the DNA to be microinjected has a large size, polyamines and high salt concentrations can be used in order to avoid mechanical breakage of this DNA, as described by Schedl et al., 1993, Nucleic Acids Res., 21: 4783-4787.

Anyone of the polynucleotides of the invention, including the DNA constructs described herein, is introduced in an embryonic stem (ES) cell line, preferably a mouse ES cell line. ES cell lines are derived from pluripotent, uncommitted cells of the inner cell mass of pre-implantation blastocysts. Preferred ES cell lines are the following : ES-E14TG2a (ATCC No. CRL-1821), ES-D3 (ATCC No. CRL1934 and No. CRL-11632), YS001 (ATCC No. CRL-11776), 36.5 (ATCC No. CRL-11116). To maintain ES cells in an uncommitted state, they are cultured in the presence of growth

inhibited feeder cells which provide the appropriate signals to preserve this embryonic phenotype and serve as a matrix for ES cell adherence. Preferred feeder cells consist of primary embryonic fibroblasts that are established from tissue of day 13- day 14 embryos of virtually any mouse strain, that are maintained in culture, such as described by Abbondanzo SJ et al., 1993, Methods in  
5 Enzymology, Academic Press, New York, pp. 803-823; and are inhibited in growth by irradiation, such as described by Robertson E., 1987, Embryo-derived stem cell lines. E.J. Robertson Ed. Teratocarcinomas and embryonic stem cells: a practical approach. IRL Press, Oxford, pp. 71, or by the presence of an inhibitory concentration of LIF, such as described by Pease S. and William R.S., 1990, Exp. Cell. Res., 190: 209-211.

10

#### Transgenic Animals

The terms "transgenic animals" or "host animals" are used herein designate non-human animals that have their genome genetically and artificially manipulated so as to include one of the nucleic acids according to the invention. Preferred animals are non-human mammals and include those belonging to a genus selected from Mus (e.g. mice), Rattus (e.g. rats) and Oryctogalus (e.g.  
15 rabbits) which have their genome artificially and genetically altered by the insertion of a nucleic acid according to the invention.

The transgenic animals of the invention all include within a plurality of their cells a cloned recombinant or synthetic DNA sequence, more specifically one of the purified or isolated nucleic acids comprising a PG1 coding sequence, a PG1 regulatory polynucleotide or a DNA sequence  
20 encoding an antisense polynucleotide such as described in the present specification.

Preferred transgenic animals according to the invention contains in their somatic cells and/or in their germ line cells a polynucleotide selected from the following group of polynucleotides:

- a) non-native, purified or isolated nucleic acid encoding a PG1 polypeptide, or a polypeptide fragment or variant thereof.
- 25 b) a non-native, purified or isolated nucleic acid comprising at least 8 consecutive nucleotides of the nucleotide sequence SEQ ID NOs: 179, 182, or 183, a nucleotide sequence complementary; in some embodiments, the length of the fragments can range from at least 8, 10, 15, 20 or 30 to 200 nucleotides, preferably from at least 10 to 50 nucleotides, more preferably from at least 40 to 50 nucleotides of SEQ ID NOs: 179, 182, or 183, or the sequence complementary thereto. In some  
30 embodiments, the fragments may comprise more than 200 nucleotides of SEQ ID NOs: 179, 182, or 183, or the sequence complementary thereto.
- c) a non-native, purified or isolated nucleic acid comprising at least 8 consecutive nucleotides of the nucleotide sequence SEQ ID NOs: 3, 69, 112-125 or 184, a sequence complementary thereto or a variant thereof; In some embodiments, the length of the fragments can range from at least 8, 10, 15,  
35 20 or 30 to 200 nucleotides, preferably from at least 10 to 50 nucleotides, more preferably from at least 40 to 50 nucleotides of SEQ ID NOs: 3, 69, 112-125 or 184, or the sequence complementary

thereto. In some embodiments, the fragments may comprise more than 200 nucleotides of SEQ ID NOs: 3, 69, 112-125 or 184, or the sequence complementary thereto.

d) a non-native, purified or isolated nucleic acid comprising a nucleotide sequence selected from the group of SEQ ID NOs: 100 to 111, a sequence complementary thereto or a fragment or a variant thereof.

e) a non-native, purified or isolated nucleic acid comprising a combination of at least two polynucleotides selected from the group consisting of SEQ ID NOs: 100 to 111, or the sequences complementary thereto wherein the polynucleotides are arranged within the nucleic acid, from the 5' end to the 3' end of said nucleic acid, in the same order than in SEQ NOs: 179, 182, or 183.

f) a non-native, purified or isolated nucleic acid comprising the nucleotide sequence SEQ ID NO: 180, or the sequences complementary thereto or a biologically active fragment or variant of the nucleotide sequence of SEQ ID NO: 180, or the sequence complementary thereto.

g) a non-native, purified or isolated nucleic acid comprising the nucleotide sequence SEQ ID NO: 181, or the sequence complementary thereto or a biologically active fragment or variant of the nucleotide sequence of SEQ ID NO: 181 or the sequence complementary thereto.

h) a polynucleotide consisting of :

(1) a nucleic acid comprising a regulatory polynucleotide of SEQ ID NO: 180 or the sequences complementary thereto or a biologically active fragment or variant thereof

(2) a polynucleotide encoding a desired polypeptide or nucleic acid.

(3) Optionally, a nucleic acid comprising a regulatory polynucleotide of SEQ NO: 181, or the sequence complementary thereto or a biologically active fragment or variant thereof.

i) a DNA construct as described previously in the present specification.

The transgenic animals of the invention thus contain specific sequences of exogenous genetic material or "non-native" such as the nucleotide sequences described above in detail.

In a first preferred embodiment, these transgenic animals is good experimental models in order to study the diverse pathologies related to cell differentiation, in particular concerning the transgenic animals within the genome of which has been inserted one or several copies of a polynucleotide encoding a native PG1 protein, or alternatively a mutant PG1 protein.

In a second preferred embodiment, these transgenic animals may express a desired polypeptide of interest under the control of the regulatory polynucleotides of the PG1 gene, leading to good yields in the synthesis of this protein of interest, and eventually a tissue specific expression of this protein of interest.

The design of the transgenic animals of the invention is made according to the conventional techniques well known from the one skilled in the art. For more details regarding the production of transgenic animals, and specifically transgenic mice, it is referred to Sandou et al. (1994) and also to US Patents Nos 4,873,191, issued Oct.10, 1989, 5,464,764 issued Nov 7, 1995 and 5,789,215, issued

Aug 4, 1998.

Transgenic animals of the present invention are produced by the application of procedures which result in an animal with a genome that has incorporated exogenous genetic material. The procedure involves obtaining the genetic material, or a portion thereof, which encodes either a PG1  
5 coding sequence, a PG1 regulatory polynucleotide or a DNA sequence encoding a PG1 antisense polynucleotide such as described in the present specification.

A recombinant polynucleotide of the invention is inserted into an embryonic or ES stem cell line. The insertion is preferably made using electroporation, such as described by Thomas K.R. et al., 1987, Cell, 51: 503-512. The cells subjected to electroporation are screened (e.g. by selection via  
10 selectable markers, by PCR or by Southern blot analysis) to find positive cells which have integrated the exogenous recombinant polynucleotide into their genome, preferably via an homologous recombination event. An illustrative positive-negative selection procedure that is used according to the invention is described by Mansour S.L. et al., 1988, Nature, 336: 348-352.

Then, the positive cells are isolated, cloned and injected into 3.5 days old blastocysts from  
15 mice, such as described by Bradley A., 1987, Production and analysis of chimaeric mice. In : E.J. Robertson (Ed.), Teratocarcinomas and embryonic stem cells : A practical approach. IRL Press, Oxford, pp.113. The blastocysts are then inserted into a female host animal and allowed to grow to term.

Alternatively, the positive ES cells are brought into contact with embryos at the 2.5 days old  
20 8-16 cell stage (morulae) such as described by Wood S.A. et al., 1993, Proc. Natl. Acad. Sci. USA, 90: 4582-4585; or by Nagy A. et al., 1993, Proc. Natl. Acad. Sci. USA, 90: 8424-8428. The ES cells being internalized to colonize extensively the blastocyst including the cells which will give rise to the germ line. The offspring of the female host are tested to determine which animals are transgenic e.g. include the inserted exogenous DNA sequence and which are wild-type.

25 Thus, the present invention also concerns a transgenic animal containing a nucleic acid, a recombinant expression vector or a recombinant host cell according to the invention.

Recombinant cell lines derived from the transgenic animals of the invention.

A further object of the invention consists of recombinant host cells obtained from a transgenic animal described herein.

30 Recombinant cell lines is established in vitro from cells obtained from any tissue of a transgenic animal according to the invention, for example by transfection of primary cell cultures with vectors expressing onc-genes such as SV40 large T antigen, as described by Chou J.Y., 1989, Mol. Endocrinol., 3: 1511-1514 ; and Shay J.W. et al., 1991, Biochem. Biophys. Acta, 1072: 1-7.

### Functional Analysis of the PG1 Polypeptides In Transgenic Animals

Using different BACs that contain the PG1 gene, we performed FISH experiment on the adenocarcinoma prostatic cell line PC3. Only one signal could be detected showing that this region of chromosome 8 is hemizygous in this tumoral cell line.

5 To study the function of PG1, it is inactivate by homologous recombination in the remaining allele of PG1 in the PC3 cell line. To inactivate the remaining PG1 allele, a knock-out targeting vector is generated by inserting two genomic DNA fragments of 3.0 and 4.3 kb (that correspond to a sequence upstream of the PG1 promoter and to part of intron 1, respectively) in the pKO Scrambler Neo TK vector (Lexicon ref V1901). Since the targeting vector contains the neomycine resistance  
10 gene as well as the Tk gene, homologous recombination is selected by adding geneticin and FIAU to the medium. The promoter, the transcriptional start site, and the first ATG contained in exon 1 on the recombinant allele is deleted by homologous recombination between the targeting vector and the remaining PG1 allele. Accordingly, no coding transcripts is initiated from the recombinant allele. The parental PC3 cells as well as cells hemizygous for the null allele are assessed for their phenotype,  
15 their growth rate in liquid culture, their ability to grow in agar (anchorage-independent growth) as well as their ability to form tumors and metastasis when injected subcutaneously in nude mice.

To determine the function of PG1 in the animal, and to generate an animal model for prostate tumorigenesis, mice in which tissue specific inactivation of the PG1 alleles can be induced are generated. For this purpose, the Cre-loxP system is utilized as described above to allow chromosome  
20 engineering to be perform directly in the animal.

First, to generate mice with a conditional null allele, two loxP sites are introduced in the murine genome, the first one 5' to the PG1 promoter and the second one 3' to the PG1 exon 1. Alternatively, to generate subtle mutations or to specifically mutate some isoforms, the loxP sites are introduced so that they flank any of the given exons or any potential set of exons. It is important to  
25 note that a functional PG1 messenger can be transcribed from these alleles until a recombination is triggered between the loxP sites by the Cre enzyme.

Second, to generate the inducer mice, the Cre gene is introduced in the mouse genome under the control of a tissue specific promoter, for example under the control of the PSA (prostate specific antigen) promoter.

30 Finally, tissue specific inactivation of the PG1 gene are induced by generating mice containing the Cre transgene that are homozygous for the recombinant PG1 allele.

### Gene Therapy

The present invention also comprises the use of the PG1 genomic DNA sequence of SEQ ID NO: 179, the PG1 cDNA of SEQ ID NO: 3, or nucleic acid encoding a mutant PG1 protein responsible  
35 for a detectable phenotype in gene therapy strategies, including antisense and triple helix strategies as described in Examples 19 and 20, below. In antisense approaches, nucleic acid sequences



complementary to an mRNA are hybridized to the mRNA intracellularly, thereby blocking the expression of the protein encoded by the mRNA. The antisense sequences may prevent gene expression through a variety of mechanisms. For example, the antisense sequences may inhibit the ability of ribosomes to translate the mRNA. Alternatively, the antisense sequences may block transport of the mRNA from the nucleus to the cytoplasm, thereby limiting the amount of mRNA available for translation. Another mechanism through which antisense sequences may inhibit gene expression is by interfering with mRNA splicing. In yet another strategy, the antisense nucleic acid is incorporated in a ribozyme capable of specifically cleaving the target mRNA.

#### Example 19

##### Preparation and Use of Antisense Oligonucleotides

The antisense nucleic acid molecules to be used in gene therapy is either DNA or RNA sequences. They may comprise a sequence complementary to the sequence of the PG1 genomic DNA of SEQ ID NO: 179, the PG1 cDNA of SEQ ID NO: 3, or a nucleic acid encoding a PG1 protein responsible for a detectable phenotype. The antisense nucleic acids should have a length and melting temperature sufficient to permit formation of an intracellular duplex having sufficient stability to inhibit the expression of the PG1 mRNA in the duplex. Strategies for designing antisense nucleic acids suitable for use in gene therapy are disclosed in Green et al., Ann. Rev. Biochem. 55:569-597 (1986) and Izant and Weintraub, Cell 36:1007-1015 (1984).

In some strategies, antisense molecules are obtained by reversing the orientation of the PG1 coding region with respect to a promoter so as to transcribe the opposite strand from that which is normally transcribed in the cell. The antisense molecules is transcribed using in vitro transcription systems such as those which employ T7 or SP6 polymerase to generate the transcript. Another approach involves transcription of PG1 antisense nucleic acids in vivo by operably linking DNA containing the antisense sequence to a promoter in an expression vector.

Alternatively, oligonucleotides which are complementary to the strand of the PG1 gene normally transcribed in the cell is synthesized in vitro. Thus, the antisense PG1 nucleic acids are complementary to the PG1 mRNA and are capable of hybridizing to the mRNA to create a duplex. In some embodiments, the PG1 antisense sequences may contain modified sugar phosphate backbones to increase stability and make them less sensitive to RNase activity. Examples of modifications suitable for use in antisense strategies are described by Rossi et al., Pharmacol. Ther. 50(2):245-254, (1991).

Various types of antisense oligonucleotides complementary to the sequence of the PG1 genomic DNA of SEQ ID NO: 179, the PG1 cDNA of SEQ ID NO: 3, or a nucleic acid encoding a PG1 protein responsible for a detectable phenotype is used. In one preferred embodiment, stable and semi-stable antisense oligonucleotides as described in International Application No. PCT WO94/23026, are used to inhibit the expression of the PG1 gene. In these molecules, the 3' end or both the 3' and 5' ends are engaged in intramolecular hydrogen bonding between complementary base pairs. These molecules are

better able to withstand exonuclease attacks and exhibit increased stability compared to conventional antisense oligonucleotides.

In another preferred embodiment, the antisense oligodeoxynucleotides described in International Application No. WO 95/04141, are used to inhibit expression of the PG1 gene.

5 In yet another preferred embodiment, the covalently cross-linked antisense oligonucleotides described in International Application No. WO 96/31523, are used to inhibit expression of the PG1 gene. These double- or single-stranded oligonucleotides comprise one or more, respectively, inter- or intra-oligonucleotide covalent cross-linkages, wherein the linkage consists of an amide bond between a primary amine group of one strand and a carboxyl group of the other strand or of the same strand,  
10 respectively, the primary amine group being directly substituted in the 2' position of the strand nucleotide monosaccharide ring, and the carboxyl group being carried by an aliphatic spacer group substituted on a nucleotide or nucleotide analog of the other strand or the same strand, respectively.

The antisense oligodeoxynucleotides and oligonucleotides disclosed in International Application No. WO 92/18522, may also be used to inhibit the expression of the PG1 gene. These molecules are  
15 stable to degradation and contain at least one transcription control recognition sequence which binds to control proteins and are effective as decoys therefor. These molecules may contain "hairpin" structures, "dumbbell" structures, "modified dumbbell" structures, "cross-linked" decoy structures and "loop" structures.

In another preferred embodiment, the cyclic double-stranded oligonucleotides described in  
20 European Patent Application No. 0 572 287 A2, are used to inhibit the expression of the PG1 gene. These ligated oligonucleotide "dumbbells" contain the binding site for a transcription factor which binds to the PG1 promoter and inhibits expression of the gene under control of the transcription factor by sequestering the factor.

Use of the closed antisense oligonucleotides disclosed in International Application No. WO  
25 92/19732, is also contemplated. Because these molecules have no free ends, they are more resistant to degradation by exonucleases than are conventional oligonucleotides. These oligonucleotides is multifunctional, interacting with several regions which are not adjacent to the target mRNA.

The appropriate level of antisense nucleic acids required to inhibit PG1 gene expression is determined using in vitro expression analysis. The antisense molecule is introduced into the cells by  
30 diffusion, injection, infection or transfection using procedures known in the art. For example, the antisense nucleic acids can be introduced into the body as a bare or naked oligonucleotide, oligonucleotide encapsulated in lipid, oligonucleotide sequence encapsidated by viral protein, or as an oligonucleotide operably linked to a promoter contained in an expression vector. The expression vector is any of a variety of expression vectors known in the art, including retroviral or viral vectors, vectors  
35 capable of extrachromosomal replication, or integrating vectors. The vectors is DNA or RNA.

The PG1 antisense molecules are introduced onto cell samples at a number of different concentrations preferably between  $1 \times 10^{-10} \text{M}$  to  $1 \times 10^{-4} \text{M}$ . Once the minimum concentration that can adequately control gene expression is identified, the optimized dose is translated into a dosage suitable for use in vivo. For example, an inhibiting concentration in culture of  $1 \times 10^{-7}$  translates into a dose of approximately 0.6 mg/kg bodyweight. Levels of oligonucleotide approaching 100 mg/kg bodyweight or higher is possible after testing the toxicity of the oligonucleotide in laboratory animals. It is additionally contemplated that cells from the vertebrate are removed, treated with the antisense oligonucleotide, and reintroduced into the vertebrate.

It is further contemplated that the PG1 antisense oligonucleotide sequence is incorporated into a ribozyme sequence to enable the antisense to specifically bind and cleave its target mRNA. For technical applications of ribozyme and antisense oligonucleotides see Rossi et al., supra.

In a preferred application of this invention, antibody-mediated tests such as RIAs and ELISA, functional assays, or radiolabeling are used to determine the effectiveness of antisense inhibition on PG1 expression.

The PG1 cDNA, the PG1 genomic DNA, and the PG1 alleles of the present invention may also be used in gene therapy approaches based on intracellular triple helix formation. Triple helix oligonucleotides are used to inhibit transcription from a genome. They are particularly useful for studying alterations in cell activity as it is associated with a particular gene. The PG1 cDNA, PG1 genomic DNA, or PG1 allele of the present invention or, more preferably, a portion of those sequences, can be used to inhibit gene expression in individuals suffering from prostate cancer or another detectable phenotype or individuals at risk for developing prostate cancer or another detectable phenotype at a later date as a result of their PG1 genotype. Similarly, a portion of the PG1 cDNA, the PG1 genomic DNA, or the PG1 alleles can be used to study the effect of inhibiting PG1 transcription within a cell. Traditionally, homopurine sequences were considered the most useful for triple helix strategies, such as those described in Example 20, below. However, homopyrimidine sequences can also inhibit gene expression. Such homopyrimidine oligonucleotides bind to the major groove at homopurine:homopyrimidine sequences. Thus, both types of sequences from the PG1 cDNA, the PG1 genomic DNA, and the PG1 alleles are contemplated within the scope of this invention.

#### Example 20

The sequences of the PG1 cDNA, the PG1 genomic DNA, and the PG1 alleles are scanned to identify 10-mer to 20-mer homopyrimidine or homopurine stretches which could be used in triple-helix based strategies for inhibiting PG1 expression. Following identification of candidate homopyrimidine or homopurine stretches, their efficiency in inhibiting PG1 expression is assessed by introducing varying amounts of oligonucleotides containing the candidate sequences into tissue culture cells which express the PG1 gene. The oligonucleotides is prepared on an oligonucleotide synthesizer or they is purchased

commercially from a company specializing in custom oligonucleotide synthesis, such as GENSET, Paris, France.

The oligonucleotides is introduced into the cells using a variety of methods known to those skilled in the art, including but not limited to calcium phosphate precipitation, DEAE-Dextran, electroporation, liposome-mediated transfection or native uptake.

Treated cells are monitored for altered cell function or reduced PG1 expression using techniques such as Northern blotting, RNase protection assays, or PCR based strategies to monitor the transcription levels of the PG1 gene in cells which have been treated with the oligonucleotide.

The oligonucleotides which are effective in inhibiting gene expression in tissue culture cells may then be introduced in vivo using the techniques described above and in Example 19 at a dosage calculated based on the in vitro results, as described in Example 19.

In some embodiments, the natural (beta) anomers of the oligonucleotide units can be replaced with alpha anomers to render the oligonucleotide more resistant to nucleases. Further, an intercalating agent such as ethidium bromide, or the like, can be attached to the 3' end of the alpha oligonucleotide to stabilize the triple helix. For information on the generation of oligonucleotides suitable for triple helix formation see Griffin et al. (Science 245:967-971 (1989)).

Alternatively, the PG1 cDNA, the PG1 genomic DNA, and the PG1 alleles of the present invention is used in gene therapy approaches in which expression of the PG1 protein is beneficial, as described in Example 21 below.

20

#### Example 21

The PG1 cDNA, the PG1 genomic DNA, and the PG1 alleles of the present invention may also be used to express the PG1 protein or a portion thereof in a host organism to produce a beneficial effect. In such procedures, the PG1 protein is transiently expressed in the host organism or stably expressed in the host organism. The expressed PG1 protein is used to treat conditions resulting from a lack of PG1 expression or conditions in which augmentation of existing levels of PG1 expression is beneficial.

A nucleic acid encoding the PG1 proteins of SEQ ID NO: 4, SEQ ID NO:5, or a PG1 allele is introduced into the host organism. The nucleic acid is introduced into the host organism using a variety of techniques known to those of skill in the art. For example, the nucleic acid is injected into the host organism as naked DNA such that the encoded PG1 protein is expressed in the host organism, thereby producing a beneficial effect.

Alternatively, the nucleic acid encoding the PG1 proteins of SEQ ID NO: 4, SEQ ID NO: 5, or a PG1 allele is cloned into an expression vector downstream of a promoter which is active in the host organism. The expression vector is any of the expression vectors designed for use in gene therapy, including viral or retroviral vectors.

The expression vector is directly introduced into the host organism such that the PG1 protein is expressed in the host organism to produce a beneficial effect. In another approach, the expression vector

is introduced into cells in vitro. Cells containing the expression vector are thereafter selected and introduced into the host organism, where they express the PG1 protein to produce a beneficial effect.

#### **IX. ISOLATION OF PG1 cDNA FROM NONHUMAN MAMMALS**

The present invention encompasses mammalian PG1 sequences including genomic and cDNA sequences, as well as polypeptide sequences. The present invention also encompasses the use of PG1 genomic and cDNA sequences of the invention, including SEQ ID NOs: 179, 3, 182, and 183, in methods of isolating and characterizing PG1 nucleotide sequences derived from nonhuman mammals, in addition to sequences derived from human sequences. The human and mouse PG1 nucleic acid sequences of the invention can be used to construct primers and probes for amplifying and identifying PG1 genes in other nonhuman animals particularly mammals. The primers and probes used to identify nonhuman PG1 sequences is selected and used for the isolation of nonhuman PG1 utilizing the same techniques described above in Examples 4, 5, 6, 12 and 13.

In addition, sequence analysis of other homologous proteins is used to optimize the sequences of these primers and probes. As described above in the Analysis of the PG1 Protein Sequence, three boxes of homology were identified in the structure of the PG1 protein product when compared to proteins from a diverse range of organisms. See Figure 9. Using the assumption that the nucleotide sequences for these homologous proteins also show a high degree of homology, it is possible to construct primers that are specific for the PCR amplification of PG1 cDNA in nonhuman mammals.

#### **Example 22**

The primers BOXIed: AATCATCAAAGCACAGTTGACTGGAT (SEQ ID NO: 77) and BOXIIIer: ATAAACCACCGTAACATCATAAATTGCATCTAA (SEQ ID NO: 78) were designed as PCR primers from the human PG1 sequences after comparison with the sequence homologies of Figure 9. The BOXIed (SEQ ID NO: 77) and BOXIIIer (SEQ ID NO: 78) primers were used to amplify a mouse PG1 cDNA sequence from mouse liver marathon-ready cDNA (Clontech) under the conditions described above in Example 4. This PCR reaction yielded a product of approximately 400 base pairs, the boxI-boxIII fragment, which was subjected to automated dideoxy terminator sequencing and electrophoresed on ABI 377 sequencers as described above. Sequence analysis confirmed very high homology to human PG1 both at the nucleic acid and protein levels.

Primers were designed for RACE analysis using the 400 base pair boxI-boxIII fragment. Further sequence information was obtained using 5' and 3' RACE reactions on mouse liver marathon cDNA using two sets of these nested PCR primers: moPG1RACE5.350: AATCAAAAGCAACGTGAGTGGC (SEQ ID NO: 94) and moPG1RACE5.276: GCAAATGCCTGACTGGCTGA (SEQ ID NO: 93) for the 5' RACE reaction and moPG1RACE3.18: CTGCCAGACAGGATGCCCTA (SEQ ID NO: 90) and moPG1RACE3.63: ACAAGTTAAAATGGCTTCCGCTG (SEQ ID NO: 91) for the 3' RACE reaction. The PCR products of the RACE reactions were sequenced by primer walking using the following primers:

moPGrace3S473: GAGATAAAAG ATAGGTTGCT CA (SEQ ID NO: 79);  
 moPGrace3S526: AAGAAACAAA TTTCCTGGG (SEQ ID NO: 80);  
 moPGrace3S597: TCTTGGGGAG TTTGACTG (SEQ ID NO: 81);  
 moPGrace5R323: GACCCCGGTG TAGTTCTC (SEQ ID NO: 82);  
 5 moPGrace5R372: CAGTAAAGCC GGTCGTC (SEQ ID NO: 83);  
 moPGrace5R444: CAGGCCAGCA GGTAGGT (SEQ ID NO: 84);  
 moPGrace5R492: AGCAGGTAGC GCATAGAGT (SEQ ID NO: 85).

Again a high degree of homology between the mouse sequence obtained from the primer walking and the human PG1 sequence was observed. An additional pair of nested primers were  
 10 designed and utilized to further extend the 3' mouse PG1 sequence in yet another RACE reaction, moPG3RACE2: TGGGCACCTG GTTGTATGGA (SEQ ID NO: 95) and moPG3RACE2n: TCCTTGGCTG CCTGTGTTT (SEQ ID NO: 96). The PCR product of this final RACE reaction was also sequenced by primer walking using the following primers:

moPG1RACE3R94: CAAATGCATG TTGGCTGT (SEQ ID NO: 92);  
 15 moPG3RACES20: GATGGCTACA CATTGTATCA C (SEQ ID NO: 97);  
 moPG3RACES5: TCCTGAATTA AATAAGGAGT TTTC (SEQ ID NO: 98);  
 moPG3RACES90: GTTGTTATT AAAGCATAAG CAAG (SEQ ID NO: 99).

The overlap in the 5' RACE, boxI-boxIII, and 3' RACE fragments allowed a single contiguous coding sequence for the mouse PG1 ortholog to be generated alignment of the three fragments. Primers  
 20 were chosen from near the 5' and 3' ends of this predicted contiguous sequence (contig) in order to confirm the existence of such a transcript. PCR amplification was performed again on mouse liver marathon-ready cDNA (Clontech) with the chosen primers, moPG15: TGGCGAGCCGAGAGGATG (SEQ ID NO: 87) and moPG13LR2: GGAAACAATGTGATACAATGTGTAGCC (SEQ ID NO: 86) under the PCR conditions described above in Example 4. The resulting PCR product was a roughly 1.2  
 25 kb DNA molecule and was shown to have an identical sequence to that of the deduced contig. Finally modified versions of the moPG15 and moPG13LR2 primers with the addition of EcoRI and BamHI sites, moPG15EcoRI: CGTGAATTCTGGCGAGCCGAGAGGATG (SEQ ID NO: 89) and moPG15BamI: CGTGGATCCGGAAACAATGTGATACAATGTGTAGCC (SEQ ID NO: 88) were used to obtain a PCR product that could be cloned into a pSKBluescript plasmid (Stratagene) cleaved  
 30 with EcoRI and BamHI restriction enzymes. The mouse PG1 cDNA in the resulting construct was subjected to automated dideoxy terminator sequencing and electrophoresed on ABI 377 sequencers as described above. The sequence for mouse PG1 cDNA is reported in SEQ ID NO: 72, and the deduced amino acid sequence corresponding to the cDNA is reported in SEQ ID NO: 74.

#### Example 23

35 A mouse BAC library was constructed by the cloning of BamHI partially digested DNA of pluripotent embryonic stem cells, cell line ES-E14TG2a (ATCC CRL-1821) into pBeloBACII vector

plasmid. Approximately fifty-six thousand clones with an average inset size of 120 kb were picked individually and pooled for PCR screening as described above for human BAC library screening. These pools were screened with STS g34292 derived from the region of the mouse PG1 transcript corresponding to exon6 of the human gene. The upstream and downstream primers defining this STS are: upstream amplification primer for g34292: ATTAAAACAC GTACTGACAC CA (SEQ ID NO: 75), and downstream amplification primer for g34292: AGTCATGGAT GGTGGATTT (SEQ ID NO: 76). BAC C0281H06 tested positive for hybridizing to g34292. This BAC was isolated and sequenced by sub-cloning into pGenDel sequencing vector. The resulting partial genomic sequence for mouse PG1 is reported in SEQ ID NO: 73. This process was repeated and the resulting partial genomic sequences for mouse PG1 is reported in SEQ ID NOs: 182 and 183.

Other mammalian PG1 cDNA and genomic sequences can be isolated by the methods of the present invention. PG1 genes in mammalian species have a region of at least 100, preferably 200, more preferably 500 nucleotides in each mammal's most abundant transcription species which has at least 75%, preferably 85%, more preferably 95% sequence homology to the most abundant human or mouse cDNA species (SEQ ID NO: 3). PG1 proteins in mammalian species have a region of at least 40, preferably 90, more preferably 160 amino acids in the deduced amino acid sequence of the most abundant PG1 transcription species which has at least 75%, preferably 85%, more preferably 95% sequence homology to the deduced amino acid sequence of the most abundant human or mouse translations species (SEQ ID NO: 4 or 74).

#### **X. METHODS FOR GENOTYPING AN INDIVIDUAL FOR BIALLELIC MARKERS**

Methods are provided to genotype a biological sample for one or more biallelic markers of the present invention, all of which is performed in vitro. Such methods of genotyping comprise determining the identity of a nucleotide at an PG1-related biallelic marker by any method known in the art. These methods find use in genotyping case-control populations in association studies as well as individuals in the context of detection of alleles of biallelic markers which, are known to be associated with a given trait, in which case both copies of the biallelic marker present in individual's genome are determined so that an individual is classified as homozygous or heterozygous for a particular allele.

These genotyping methods can be performed nucleic acid samples derived from a single individual or pooled DNA samples.

Genotyping can be performed using similar methods as those described above for the identification of the biallelic markers, or using other genotyping methods such as those further described below. In preferred embodiments, the comparison of sequences of amplified genomic fragments from different individuals is used to identify new biallelic markers whereas microsequencing is used for genotyping known biallelic markers in diagnostic and association study applications.

**X.A. Source of DNA for genotyping**

Any source of nucleic acids, in purified or non-purified form, can be utilized as the starting nucleic acid, provided it contains or is suspected of containing the specific nucleic acid sequence desired. DNA or RNA is extracted from cells, tissues, body fluids. As for the source of genomic  
5 DNA to be subjected to analysis, any test sample can be foreseen without any particular limitation. These test samples include biological samples, which can be tested by the methods of the present invention described herein, and include human and animal body fluids such as whole blood, serum, plasma, cerebrospinal fluid, urine, lymph fluids, and various external secretions of the respiratory, intestinal and genitourinary tracts, tears, saliva, milk, white blood cells, myelomas and the like;  
10 biological fluids such as cell culture supernatants; fixed tissue specimens including tumor and non-tumor tissue and lymph node tissues; bone marrow aspirates and fixed cell specimens. The preferred source of genomic DNA used in the present invention is from peripheral venous blood of each donor. Techniques to prepare genomic DNA from biological samples are well known to the skilled technician. While nucleic acids for use in the genotyping methods of the invention can be derived  
15 from any mammalian source, the test subjects and individuals from which nucleic acid samples are taken are generally understood to be human.

**X.B. Amplification Of DNA Fragments Comprising Biallelic Markers**

Methods and polynucleotides are provided to amplify a segment of nucleotides comprising one or more biallelic marker of the present invention. It will be appreciated that amplification of  
20 DNA fragments comprising biallelic markers is used in various methods and for various purposes and is not restricted to genotyping. Nevertheless, many genotyping methods, although not all, require the previous amplification of the DNA region carrying the biallelic marker of interest. Such methods specifically increase the concentration or total number of sequences that span the biallelic marker or include that site and sequences located either distal or proximal to it. Diagnostic assays may also rely  
25 on amplification of DNA segments carrying a biallelic marker of the present invention.

Amplification of DNA is achieved by any method known in the art. The established PCR (polymerase chain reaction) method or by developments thereof or alternatives. Amplification methods which can be utilized herein include but are not limited to Ligase Chain Reaction (LCR) as described in EP A 320 308 and EP A 439 182, Gap LCR (Wolcott, M.J., Clin. Microbiol. Rev. 5:370-  
30 386), the so-called "NASBA" or "3SR" technique described in Guatelli J.C. et al. (*Proc. Natl. Acad. Sci. USA* 87:1874-1878, 1990) and in Compton J. (*Nature* 350:91-92, 1991), Q-beta amplification as described in European Patent Application no 4544610, strand displacement amplification as described in Walker et al. (*Clin. Chem.* 42:9-13, 1996) and EP A 684 315 and, target mediated amplification as described in PCT Publication WO 9322461.

35 LCR and Gap LCR are exponential amplification techniques, both depend on DNA ligase to join adjacent primers annealed to a DNA molecule. In Ligase Chain Reaction (LCR), probe pairs are



used which include two primary (first and second) and two secondary (third and fourth) probes, all of which are employed in molar excess to target. The first probe hybridizes to a first segment of the target strand and the second probe hybridizes to a second segment of the target strand, the first and second segments being contiguous so that the primary probes abut one another in 5' phosphate-  
5 3'hydroxyl relationship, and so that a ligase can covalently fuse or ligate the two probes into a fused product. In addition, a third (secondary) probe can hybridize to a portion of the first probe and a fourth (secondary) probe can hybridize to a portion of the second probe in a similar abutting fashion. Of course, if the target is initially double stranded, the secondary probes also will hybridize to the target complement in the first instance. Once the ligated strand of primary probes is separated from  
10 the target strand, it will hybridize with the third and fourth probes which can be ligated to form a complementary, secondary ligated product. It is important to realize that the ligated products are functionally equivalent to either the target or its complement. By repeated cycles of hybridization and ligation, amplification of the target sequence is achieved. A method for multiplex LCR has also been described (WO 9320227). Gap LCR (GLCR) is a version of LCR where the probes are not adjacent  
15 but are separated by 2 to 3 bases.

For amplification of mRNAs, it is within the scope of the present invention to reverse transcribe mRNA into cDNA followed by polymerase chain reaction (RT-PCR); or, to use a single enzyme for both steps as described in U.S. Patent No. 5,322,770 or, to use Asymmetric Gap LCR (RT-AGLCR) as described by Marshall R.L. et al. (*PCR Methods and Applications* 4:80-84, 1994).  
20 AGLCR is a modification of GLCR that allows the amplification of RNA.

Some of these amplification methods are particularly suited for the detection of single nucleotide polymorphisms and allow the simultaneous amplification of a target sequence and the identification of the polymorphic nucleotide as it is further described in X.C.

The PCR technology is the preferred amplification technique used in the present invention. A  
25 variety of PCR techniques are familiar to those skilled in the art. For a review of PCR technology, see *Molecular Cloning to Genetic Engineering* White, B.A. Ed. in *Methods in Molecular Biology* 67: Humana Press, Totowa (1997) and the publication entitled "PCR Methods and Applications" (1991, Cold Spring Harbor Laboratory Press). In each of these PCR procedures, PCR primers on either side of the nucleic acid sequences to be amplified are added to a suitably prepared nucleic acid sample along  
30 with dNTPs and a thermostable polymerase such as Taq polymerase, Pfu polymerase, or Vent polymerase. The nucleic acid in the sample is denatured and the PCR primers are specifically hybridized to complementary nucleic acid sequences in the sample. The hybridized primers are extended. Thereafter, another cycle of denaturation, hybridization, and extension is initiated. The cycles are repeated multiple times to produce an amplified fragment containing the nucleic acid sequence between  
35 the primer sites. PCR has further been described in several patents including US Patents 4,683,195, 4,683,202 and 4,965,188.

The identification of biallelic markers as described above allows the design of appropriate oligonucleotides, which can be used as primers to amplify DNA fragments comprising the biallelic markers of the present invention. Amplification can be performed using the primers initially used to discover new biallelic markers which are described herein or any set of primers allowing the  
5 amplification of a DNA fragment comprising a biallelic marker of the present invention. Primers can be prepared by any suitable method. As for example, direct chemical synthesis by a method such as the phosphodiester method of Narang S.A. et al. (*Methods Enzymol.* 68:90-98, 1979), the phosphodiester method of Brown E.L. et al. (*Methods Enzymol.* 68:109-151, 1979), the diethylphosphoramidite method of Beaucage et al. (*Tetrahedron Lett.* 22:1859-1862, 1981) and the  
10 solid support method described in EP 0 707 592.

In some embodiments the present invention provides primers for amplifying a DNA fragment containing one or more biallelic markers of the present invention. It will be appreciated that the amplification primers listed in the present specification are merely exemplary and that any other set of primers which produce amplification products containing one or more biallelic markers of the present  
15 invention.

The primers are selected to be substantially complementary to the different strands of each specific sequence to be amplified. The length of the primers of the present invention can range from 8 to 100 nucleotides, preferably from 8 to 50, 8 to 30 or more preferably 8 to 25 nucleotides. Shorter primers tend to lack specificity for a target nucleic acid sequence and generally require cooler  
20 temperatures to form sufficiently stable hybrid complexes with the template. Longer primers are expensive to produce and can sometimes self-hybridize to form hairpin structures. The formation of stable hybrids depends on the melting temperature  $T_m$  of the DNA. The  $T_m$  depends on the length of the primer, the ionic strength of the solution and the G+C content. The higher the G+C content of the primer, the higher is the melting temperature because G:C pairs are held by three H bonds whereas  
25 A:T pairs have only two. The G+C content of the amplification primers of the present invention preferably ranges between 10 and 75 %, more preferably between 35 and 60 %, and most preferably between 40 and 55 %. The appropriate length for primers under a particular set of assay conditions is empirically determined by one of skill in the art.

The spacing of the primers determines the length of the segment to be amplified. In the  
30 context of the present invention amplified segments carrying biallelic markers can range in size from at least about 25 bp to 35 kbp. Amplification fragments from 25-3000 bp are typical, fragments from 50-1000 bp are preferred and fragments from 100-600 bp are highly preferred. It will be appreciated that amplification primers for the biallelic markers is any sequence which allow the specific amplification of any DNA fragment carrying the markers. Amplification primers is labeled or  
35 immobilized on a solid support as described in Section II.

#### **X.C. Methods of Genotyping DNA samples for Biallelic Markers**

Any method known in the art can be used to identify the nucleotide present at a biallelic marker site. Since the biallelic marker allele to be detected has been identified and specified in the present invention, detection will prove routine for one of ordinary skill in the art by employing any of a number of techniques. Many genotyping methods require the previous amplification of the DNA region carrying the biallelic marker of interest. While the amplification of target or signal is often preferred at present, ultrasensitive detection methods which do not require amplification are also encompassed by the present genotyping methods. Methods well-known to those skilled in the art that can be used to detect biallelic polymorphisms include methods such as, conventional dot blot analyzes, single strand conformational polymorphism analysis (SSCP) described by Orita et al. (*Proc. Natl. Acad. Sci. U.S.A* 86:27776-2770, 1989), denaturing gradient gel electrophoresis (DGGE), heteroduplex analysis, mismatch cleavage detection, and other conventional techniques as described in Sheffield, V.C. et al. (*Proc. Natl. Acad. Sci. USA* 49:699-706, 1991), White et al. (*Genomics* 12:301-306, 1992), Grompe, M. et al. (*Proc. Natl. Acad. Sci. USA* 86:5855-5892, 1989) and Grompe, M. (*Nature Genetics* 5:111-117, 1993). Another method for determining the identity of the nucleotide present at a particular polymorphic site employs a specialized exonuclease-resistant nucleotide derivative as described in US patent 4,656,127.

Preferred methods involve directly determining the identity of the nucleotide present at a biallelic marker site by sequencing assay, allele-specific amplification assay, or hybridization assay. The following is a description of some preferred methods. A highly preferred method is the microsequencing technique. The term "sequencing assay" is used herein to refer to polymerase extension of duplex primer/template complexes and includes both traditional sequencing and microsequencing.

#### 1) Sequencing assays

The nucleotide present at a polymorphic site can be determined by sequencing methods. In a preferred embodiment, DNA samples are subjected to PCR amplification before sequencing as described above. Methods for sequencing DNA using either the dideoxy-mediated method (Sanger method) or the Maxam-Gilbert method are widely known to those of ordinary skill in the art. Such methods are for example disclosed in Maniatis et al. (*Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Second Edition, 1989). Alternative approaches include hybridization to high-density DNA probe arrays as described in Chee et al. (*Science* 274, 610, 1996).

Preferably, the amplified DNA is subjected to automated dideoxy terminator sequencing reactions using a dye-primer cycle sequencing protocol. The products of the sequencing reactions are run on sequencing gels and the sequences are determined using gel image analysis.

The polymorphism detection in a pooled sample is based on the presence of superimposed peaks in the electrophoresis pattern resulting from different bases occurring at the same position. Because each dideoxy terminator is labeled with a different fluorescent molecule, the two peaks

corresponding to a biallelic site present distinct colors corresponding to two different nucleotides at the same position on the sequence. However, the presence of two peaks can be an artifact due to background noise. To exclude such an artifact, the two DNA strands are sequenced and a comparison between the peaks is carried out. In order to be registered as a polymorphic sequence, the polymorphism has to be detected on both strands.

The above procedure permits those amplification products, which contain biallelic markers to be identified. The detection limit for the frequency of biallelic polymorphisms detected by sequencing pools of 100 individuals is approximately 0.1 for the minor allele, as verified by sequencing pools of known allelic frequencies.

#### 10 Microsequencing assays

In microsequencing methods, the nucleotide at a polymorphic site in a target DNA is detected by a single nucleotide primer extension reaction. This method involves appropriate microsequencing primers which, hybridize just upstream of the polymorphic base of interest in the target nucleic acid. A polymerase is used to specifically extend the 3' end of the primer with one single ddNTP (chain terminator) complementary to the nucleotide at the polymorphic site. Next the identity of the incorporated nucleotide is determined in any suitable way.

Typically, microsequencing reactions are carried out using fluorescent ddNTPs and the extended microsequencing primers are analyzed by electrophoresis on ABI 377 sequencing machines to determine the identity of the incorporated nucleotide as described in EP 412 883. Alternatively capillary electrophoresis can be used in order to process a higher number of assays simultaneously.

Different approaches can be used to detect the nucleotide added to the microsequencing primer. A homogeneous phase detection method based on fluorescence resonance energy transfer has been described by Chen and Kwok (*Nucleic Acids Research* 25:347-353 1997) and Chen et al. (*Proc. Natl. Acad. Sci. USA* 94/20 10756-10761, 1997). In this method amplified genomic DNA fragments containing polymorphic sites are incubated with a 5'-fluorescein-labeled primer in the presence of allelic dye-labeled dideoxyribonucleoside triphosphates and a modified Taq polymerase. The dye-labeled primer is extended one base by the dye-terminator specific for the allele present on the template. At the end of the genotyping reaction, the fluorescence intensities of the two dyes in the reaction mixture are analyzed directly without separation or purification. All these steps can be performed in the same tube and the fluorescence changes can be monitored in real time. Alternatively, the extended primer is analyzed by MALDI-TOF Mass Spectrometry. The base at the polymorphic site is identified by the mass added onto the microsequencing primer (see Haff L.A. and Smirnov I.P., *Genome Research*, 7:378-388, 1997).

Microsequencing is achieved by the established microsequencing method or by developments or derivatives thereof. Alternative methods include several solid-phase microsequencing techniques. The basic microsequencing protocol is the same as described previously, except that the method is

conducted as a heterogenous phase assay, in which the primer or the target molecule is immobilized or captured onto a solid support. To simplify the primer separation and the terminal nucleotide addition analysis, oligonucleotides are attached to solid supports or are modified in such ways that permit affinity separation as well as polymerase extension. The 5' ends and internal nucleotides of synthetic oligonucleotides can be modified in a number of different ways to permit different affinity separation approaches, e.g., biotinylation. If a single affinity group is used on the oligonucleotides, the oligonucleotides can be separated from the incorporated terminator reagent. This eliminates the need of physical or size separation. More than one oligonucleotide can be separated from the terminator reagent and analyzed simultaneously if more than one affinity group is used. This permits the analysis of several nucleic acid species or more nucleic acid sequence information per extension reaction. The affinity group need not be on the priming oligonucleotide but could alternatively be present on the template. For example, immobilization can be carried out via an interaction between biotinylated DNA and streptavidin-coated microtitration wells or avidin-coated polystyrene particles. In the same manner oligonucleotides or templates is attached to a solid support in a high-density format. In such solid phase microsequencing reactions, incorporated ddNTPs can be radiolabeled (Syvänen, *Clinica Chimica Acta* 226:225-236, 1994) or linked to fluorescein (Livak and Hainer, *Human Mutation* 3:379-385, 1994). The detection of radiolabeled ddNTPs can be achieved through scintillation-based techniques. The detection of fluorescein-linked ddNTPs can be based on the binding of anti fluorescein antibody conjugated with alkaline phosphatase, followed by incubation with a chromogenic substrate (such as *p*-nitrophenyl phosphate). Other possible reporter-detection pairs include: ddNTP linked to dinitrophenyl (DNP) and anti-DNP alkaline phosphatase conjugate (Harju et al., *Clin. Chem.* 39/11 2282-2287, 1993) or biotinylated ddNTP and horseradish peroxidase-conjugated streptavidin with *o*-phenylenediamine as a substrate (WO 92/15712). As yet another alternative solid-phase microsequencing procedure, Nyren et al. (*Analytical Biochemistry* 208:171-175, 1993) described a method relying on the detection of DNA polymerase activity by an enzymatic luminometric inorganic pyrophosphate detection assay (ELIDA).

Pastinen et al. (*Genome research* 7:606-614, 1997) describe a method for multiplex detection of single nucleotide polymorphism in which the solid phase minisequencing principle is applied to an oligonucleotide array format. High-density arrays of DNA probes attached to a solid support (DNA chips) are further described in X.C.5.

In one aspect the present invention provides polynucleotides and methods to genotype one or more biallelic markers of the present invention by performing a microsequencing assay. It will be appreciated that any primer having a 3' end immediately adjacent to the polymorphic nucleotide is used. However, polynucleotides comprising at least 8, 12, 15, 20, 25, or 30 consecutive nucleotides of the sequence immediately adjacent to the biallelic marker and having a 3' terminus immediately

upstream of the corresponding biallelic marker are well suited for determining the identity of a nucleotide at biallelic marker site.

Similarly, it will be appreciated that microsequencing analysis is performed for any biallelic marker or any combination of biallelic markers of the present invention.

#### 5 Mismatch detection assays based on polymerases and ligases

In one aspect the present invention provides polynucleotides and methods to determine the allele of one or more biallelic markers of the present invention in a biological sample, by mismatch detection assays based on polymerases and/or ligases. These assays are based on the specificity of polymerases and ligases. Polymerization reactions places particularly stringent requirements on  
10 correct base pairing of the 3' end of the amplification primer and the joining of two oligonucleotides hybridized to a target DNA sequence is quite sensitive to mismatches close to the ligation site, especially at the 3' end. Methods, primers and various parameters to amplify DNA fragments comprising biallelic markers of the present invention are further described above in X.B.

#### Allele specific amplification

15 Discrimination between the two alleles of a biallelic marker can also be achieved by allele specific amplification, a selective strategy, whereby one of the alleles is amplified without amplification of the other allele. This is accomplished by placing the polymorphic base at the 3' end of one of the amplification primers. Because the extension forms from the 3' end of the primer, a mismatch at or near this position has an inhibitory effect on amplification. Therefore, under  
20 appropriate amplification conditions, these primers only direct amplification on their complementary allele. Designing the appropriate allele-specific primer and the corresponding assay conditions are well with the ordinary skill in the art.

#### Ligation/amplification based methods

The "Oligonucleotide Ligation Assay" (OLA) uses two oligonucleotides which are designed  
25 to be capable of hybridizing to abutting sequences of a single strand of a target molecules. One of the oligonucleotides is biotinylated, and the other is detectably labeled. If the precise complementary sequence is found in a target molecule, the oligonucleotides will hybridize such that their termini abut, and create a ligation substrate that can be captured and detected. OLA is capable of detecting single nucleotide polymorphisms and is advantageously combined with PCR as described by  
30 Nickerson D.A. et al. (*Proc. Natl. Acad. Sci. U.S.A.* 87:8923-8927, 1990). In this method, PCR is used to achieve the exponential amplification of target DNA, which is then detected using OLA.

Other methods which are particularly suited for the detection of single nucleotide polymorphism include LCR (ligase chain reaction), Gap LCR (GLCR) which are described above in X.B. As mentioned above LCR uses two pairs of probes to exponentially amplify a specific target.  
35 The sequences of each pair of oligonucleotides, is selected to permit the pair to hybridize to abutting sequences of the same strand of the target. Such hybridization forms a substrate for a template-

dependant ligase. In accordance with the present invention, LCR can be performed with oligonucleotides having the proximal and distal sequences of the same strand of a biallelic marker site. In one embodiment, either oligonucleotide will be designed to include the biallelic marker site. In such an embodiment, the reaction conditions are selected such that the oligonucleotides can be  
5 ligated together only if the target molecule either contains or lacks the specific nucleotide that is complementary to the biallelic marker on the oligonucleotide. In an alternative embodiment, the oligonucleotides will not include the biallelic marker, such that when they hybridize to the target molecule, a "gap" is created as described in WO 90/01069. This gap is then "filled" with complementary dNTPs (as mediated by DNA polymerase), or by an additional pair of  
10 oligonucleotides. Thus at the end of each cycle, each single strand has a complement capable of serving as a target during the next cycle and exponential allele-specific amplification of the desired sequence is obtained.

Ligase/Polymerase-mediated Genetic Bit Analysis<sup>TM</sup> is another method for determining the identity of a nucleotide at a preselected site in a nucleic acid molecule (WO 95/21271). This method  
15 involves the incorporation of a nucleoside triphosphate that is complementary to the nucleotide present at the preselected site onto the terminus of a primer molecule, and their subsequent ligation to a second oligonucleotide. The reaction is monitored by detecting a specific label attached to the reaction's solid phase or by detection in solution.

## 2) Hybridization assay methods

20 A preferred method of determining the identity of the nucleotide present at a biallelic marker site involves nucleic acid hybridization. The hybridization probes, which can be conveniently used in such reactions, preferably include the probes defined herein. Any hybridization assay is used including Southern hybridization, Northern hybridization, dot blot hybridization and solid-phase hybridization (see Sambrook et al., Molecular Cloning – A Laboratory Manual, Second Edition, Cold  
25 Spring Harbor Press, N.Y., 1989).

Hybridization refers to the formation of a duplex structure by two single stranded nucleic acids due to complementary base pairing. Hybridization can occur between exactly complementary nucleic acid strands or between nucleic acid strands that contain minor regions of mismatch. Specific probes can be designed that hybridize to one form of a biallelic marker and not to the other and  
30 therefore are able to discriminate between different allelic forms. Allele-specific probes are often used in pairs, one member of a pair showing perfect match to a target sequence containing the original allele and the other showing a perfect match to the target sequence containing the alternative allele. Hybridization conditions should be sufficiently stringent that there is a significant difference in hybridization intensity between alleles, and preferably an essentially binary response, whereby a  
35 probe hybridizes to only one of the alleles. Stringent, sequence specific hybridization conditions, under which a probe will hybridize only to the exactly complementary target sequence are well known

in the art (Sambrook et al., *Molecular Cloning – A Laboratory Manual*, Second Edition, Cold Spring Harbor Press, N.Y., 1989). Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point <sup>TM</sup> for the specific sequence at a defined ionic strength and pH. By way of example and not limitation, procedures using conditions of high stringency are as follows:

Prehybridization of filters containing DNA is carried out for 8 h to overnight at 65°C in buffer composed of 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 µg/ml denatured salmon sperm DNA. Filters are hybridized for 48 h at 65°C, the preferred hybridization temperature, in prehybridization mixture containing 100 µg/ml denatured salmon sperm DNA and 5-20 X 10<sup>6</sup> cpm of <sup>32</sup>P-labeled probe. Alternatively, the hybridization step can be performed at 65°C in the presence of SSC buffer, 1 x SSC corresponding to 0.15M NaCl and 0.05 M Na citrate. Subsequently, filter washes can be done at 37°C for 1 h in a solution containing 2X SSC, 0.01% PVP, 0.01% Ficoll, and 0.01% BSA, followed by a wash in 0.1X SSC at 50°C for 45 min. Alternatively, filter washes can be performed in a solution containing 2 x SSC and 0.1% SDS, or 0.5 x SSC and 0.1% SDS, or 0.1 x SSC and 0.1% SDS at 68°C for 15 minute intervals. Following the wash steps, the hybridized probes are detectable by autoradiography. By way of example and not limitation, procedures using conditions of intermediate stringency are as follows: Filters containing DNA are prehybridized, and then hybridized at a temperature of 60°C in the presence of a 5 x SSC buffer and labeled probe. Subsequently, filters washes are performed in a solution containing 2x SSC at 50°C and the hybridized probes are detectable by autoradiography. Other conditions of high and intermediate stringency which is used are well known in the art and as cited in Sambrook et al. (*Molecular Cloning - A Laboratory Manual*, Second Edition, Cold Spring Harbor Press, N.Y., 1989) and Ausubel et al. (*Current Protocols in Molecular Biology*, Green Publishing Associates and Wiley Interscience, N.Y., 1989).

Although such hybridizations can be performed in solution, it is preferred to employ a solid-phase hybridization assay. The target DNA comprising a biallelic marker of the present invention is amplified prior to the hybridization reaction. The presence of a specific allele in the sample is determined by detecting the presence or the absence of stable hybrid duplexes formed between the probe and the target DNA. The detection of hybrid duplexes can be carried out by a number of methods. Various detection assay formats are well known which utilize detectable labels bound to either the target or the probe to enable detection of the hybrid duplexes. Typically, hybridization duplexes are separated from unhybridized nucleic acids and the labels bound to the duplexes are then detected. Those skilled in the art will recognize that wash steps is employed to wash away excess target DNA or probe. Standard heterogeneous assay formats are suitable for detecting the hybrids using the labels present on the primers and probes.



Two recently developed assays allow hybridization-based allele discrimination with no need for separations or washes (see Landegren U. et al., *Genome Research*, 8:769-776,1998). The TaqMan assay takes advantage of the 5' nuclease activity of Taq DNA polymerase to digest a DNA probe annealed specifically to the accumulating amplification product. TaqMan probes are labeled with a donor-acceptor dye pair that interacts via fluorescence energy transfer. Cleavage of the TaqMan probe by the advancing polymerase during amplification dissociates the donor dye from the quenching acceptor dye, greatly increasing the donor fluorescence. All reagents necessary to detect two allelic variants can be assembled at the beginning of the reaction and the results are monitored in real time (see Livak et al., *Nature Genetics*, 9:341-342, 1995). In an alternative homogeneous hybridization based procedure, molecular beacons are used for allele discriminations. Molecular beacons are hairpin-shaped oligonucleotide probes that report the presence of specific nucleic acids in homogeneous solutions. When they bind to their targets they undergo a conformational reorganization that restores the fluorescence of an internally quenched fluorophore (Tyagi et al., *Nature Biotechnology*, 16:49-53, 1998).

The polynucleotides provided herein can be used in hybridization assays for the detection of biallelic marker alleles in biological samples. These probes are characterized in that they preferably comprise between 8 and 50 nucleotides, and in that they are sufficiently complementary to a sequence comprising a biallelic marker of the present invention to hybridize thereto and preferably sufficiently specific to be able to discriminate the targeted sequence for only one nucleotide variation. The GC content in the probes of the invention usually ranges between 10 and 75 %, preferably between 35 and 60 %, and more preferably between 40 and 55 %. The length of these probes can range from 10, 15, 20, or 30 to at least 100 nucleotides, preferably from 10 to 50, more preferably from 18 to 35 nucleotides. A particularly preferred probe is 25 nucleotides in length. Preferably the biallelic marker is within 4 nucleotides of the center of the polynucleotide probe. In particularly preferred probes the biallelic marker is at the center of said polynucleotide. Shorter probes may lack specificity for a target nucleic acid sequence and generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. Longer probes are expensive to produce and can sometimes self-hybridize to form hairpin structures. Methods for the synthesis of oligonucleotide probes have been described above and can be applied to the probes of the present invention.

Preferably the probes of the present invention are labeled or immobilized on a solid support. Labels and solid supports are further described in II. Detection probes are generally nucleic acid sequences or uncharged nucleic acid analogs such as, for example peptide nucleic acids which are disclosed in International Patent Application WO 92/20702, morpholino analogs which are described in U.S. Patents Numbered 5,185,444; 5,034,506 and 5,142,047. The probe may have to be rendered "non-extendable" in that additional dNTPs cannot be added to the probe. In and of themselves analogs usually are non-extendable and nucleic acid probes can be rendered non-extendable by

modifying the 3' end of the probe such that the hydroxyl group is no longer capable of participating in elongation. For example, the 3' end of the probe can be functionalized with the capture or detection label to thereby consume or otherwise block the hydroxyl group. Alternatively, the 3' hydroxyl group simply can be cleaved, replaced or modified, U.S. Patent Application Serial No. 07/049,061 filed 5 April 19, 1993 describes modifications, which can be used to render a probe non-extendable.

The probes of the present invention are useful for a number of purposes. They can be used in Southern hybridization to genomic DNA or Northern hybridization to mRNA. The probes can also be used to detect PCR amplification products. By assaying the hybridization to an allele specific probe, one can detect the presence or absence of a biallelic marker allele in a given sample.

10 High-Throughput parallel hybridizations in array format are specifically encompassed within "hybridization assays" and are described below.

#### **Hybridization to addressable arrays of oligonucleotides**

Hybridization assays based on oligonucleotide arrays rely on the differences in hybridization stability of short oligonucleotides to perfectly matched and mismatched target sequence variants. 15 Efficient access to polymorphism information is obtained through a basic structure comprising high-density arrays of oligonucleotide probes attached to a solid support (the chip) at selected positions. Each DNA chip can contain thousands to millions of individual synthetic DNA probes arranged in a grid-like pattern and miniaturized to the size of a dime.

The chip technology has already been applied with success in numerous cases. For example, 20 the screening of mutations has been undertaken in the BRCA1 gene, in *S. cerevisiae* mutant strains, and in the protease gene of HIV-1 virus (Hacia et al., *Nature Genetics*, 14(4):441-447, 1996; Shoemaker et al., *Nature Genetics*, 14(4):450-456, 1996; Kozal et al., *Nature Medicine*, 2:753-759, 1996). Chips of various formats for use in detecting biallelic polymorphisms can be produced on a customized basis by Affymetrix (GeneChip™), Hyseq (HyChip and HyGnostics), and Protogene 25 Laboratories.

In general, these methods employ arrays of oligonucleotide probes that are complementary to target nucleic acid sequence segments from an individual which, target sequences include a polymorphic marker. EP785280 describes a tiling strategy for the detection of single nucleotide polymorphisms. Briefly, arrays may generally be "tiled" for a large number of specific 30 polymorphisms. By "tiling" is generally meant the synthesis of a defined set of oligonucleotide probes which is made up of a sequence complementary to the target sequence of interest, as well as preselected variations of that sequence, e.g., substitution of one or more given positions with one or more members of the basis set of monomers, i.e. nucleotides. Tiling strategies are further described in PCT application No. WO 95/11995. In a particular aspect, arrays are tiled for a number of specific, 35 identified biallelic marker sequences. In particular the array is tiled to include a number of detection blocks, each detection block being specific for a specific biallelic marker or a set of biallelic markers.

For example, a detection block is tiled to include a number of probes, which span the sequence segment that includes a specific polymorphism. To ensure probes that are complementary to each allele, the probes are synthesized in pairs differing at the biallelic marker. In addition to the probes differing at the polymorphic base, monosubstituted probes are also generally tiled within the detection block. These monosubstituted probes have bases at and up to a certain number of bases in either direction from the polymorphism, substituted with the remaining nucleotides (selected from A, T, G, C and U). Typically the probes in a tiled detection block will include substitutions of the sequence positions up to and including those that are 5 bases away from the biallelic marker. The monosubstituted probes provide internal controls for the tiled array, to distinguish actual hybridization from artefactual cross-hybridization. Upon completion of hybridization with the target sequence and washing of the array, the array is scanned to determine the position on the array to which the target sequence hybridizes. The hybridization data from the scanned array is then analyzed to identify which allele or alleles of the biallelic marker are present in the sample. Hybridization and scanning is carried out as described in PCT application No. WO 92/10092 and WO 95/11995 and US patent No. 5,424,186.

#### 5) Integrated Systems

Another technique, which is used to analyze polymorphisms, includes multicomponent integrated systems, which miniaturize and compartmentalize processes such as PCR and capillary electrophoresis reactions in a single functional device. An example of such technique is disclosed in US patent 5,589,136, which describes the integration of PCR amplification and capillary electrophoresis in chips.

Integrated systems can be envisaged mainly when microfluidic systems are used. These systems comprise a pattern of microchannels designed onto a glass, silicon, quartz, or plastic wafer included on a microchip. The movements of the samples are controlled by electric, electroosmotic or hydrostatic forces applied across different areas of the microchip to create functional microscopic valves and pumps with no moving parts. Varying the voltage controls the liquid flow at intersections between the micro-machined channels and changes the liquid flow rate for pumping across different sections of the microchip.

For genotyping biallelic markers, the microfluidic system may integrate nucleic acid amplification, microsequencing, capillary electrophoresis and a detection method such as laser-induced fluorescence detection.

### **XI. METHODS OF GENETIC ANALYSIS USING THE BIALLELIC MARKERS OF THE PRESENT INVENTION**

The methods available for the genetic analysis of complex traits fall into different categories (see Lander and Schork, *Science*, 265, 2037-2048, 1994). In general, the biallelic markers of the present invention find use in any method known in the art to demonstrate a statistically significant

correlation between a genotype and a phenotype. The biallelic markers is used in linkage analysis and in allele-sharing methods. Preferably, the biallelic markers of the present invention are used to identify genes associated with detectable traits using association studies, an approach which does not require the use of affected families and which permits the identification of genes associated with  
5 complex and sporadic traits.

The genetic analysis using the biallelic markers of the present invention is conducted on any scale. The whole set of biallelic markers of the present invention or any subset of biallelic markers of the present invention is used. In some embodiments, any additional set of genetic markers including a biallelic marker of the present invention is used. As mentioned above, it should be noted that the  
10 biallelic markers of the present invention is included in any complete or partial genetic map of the human genome. These different uses are specifically contemplated in the present invention and claims.

#### **XI.A. Linkage Analysis**

Until recently, the identification of genes linked with detectable traits has mainly relied on a  
15 statistical approach called linkage analysis. Linkage analysis involves proposing a model to explain the inheritance pattern of phenotypes and genotypes observed in a pedigree. Linkage analysis is based upon establishing a correlation between the transmission of genetic markers and that of a specific trait throughout generations within a family. In this approach, all members of a series of affected families are genotyped with a few hundred markers, typically microsatellite markers, which  
20 are distributed at an average density of one every 10 Mb. By comparing genotypes in all family members, one can attribute sets of alleles to parental haploid genomes (haplotyping or phase determination). The origin of recombined fragments is then determined in the offspring of all families. Those that co-segregate with the trait are tracked. After pooling data from all families, statistical methods are used to determine the likelihood that the marker and the trait are segregating  
25 independently in all families. As a result of the statistical analysis, one or several regions having a high probability of harboring a gene linked to the trait are selected as candidates for further analysis. The result of linkage analysis is considered as significant (i.e. there is a high probability that the region contains a gene involved in a detectable trait) when the chance of independent segregation of the marker and the trait is lower than 1 in 1000 (expressed as a LOD score  $> 3$ ). Generally, the length  
30 of the candidate region identified as having a LOD score of greater than 3 using linkage analysis is between 2 and 20Mb. Once a candidate region is identified as described above, analysis of recombinant individuals using additional markers allows further delineation of the candidate region. Linkage analysis studies have generally relied on the use of a maximum of 5,000 microsatellite markers, thus limiting the maximum theoretical attainable resolution of linkage analysis to about 600  
35 kb on average.

Linkage analysis has been successfully applied to map simple genetic traits that show clear Mendelian inheritance patterns and which have a high penetrance (i.e., the ratio between the number of trait positive carriers of allele  $a$  and the total number of  $a$  carriers in the population). About 100 pathological trait-causing genes were discovered using linkage analysis over the last 10 years. In most of these cases, the majority of affected individuals had affected relatives and the detectable trait was rare in the general population (frequencies less than 0.1%). In about 10 cases, such as Alzheimer's Disease, breast cancer, and Type II diabetes, the detectable trait was more common but the allele associated with the detectable trait was rare in the affected population. Thus, the alleles associated with these traits were not responsible for the trait in all sporadic cases.

Linkage analysis suffers from a variety of drawbacks. First, linkage analysis is limited by its reliance on the choice of a genetic model suitable for each studied trait. Furthermore, as already mentioned, the resolution attainable using linkage analysis is limited, and complementary studies are required to refine the analysis of the typical 2Mb to 20Mb regions initially identified through linkage analysis. In addition, linkage analysis approaches have proven difficult when applied to complex genetic traits, such as those due to the combined action of multiple genes and/or environmental factors. In such cases, too large an effort and cost are needed to recruit the adequate number of affected families required for applying linkage analysis to these situations, as recently discussed by Risch, N. and Merikangas, K. (*Science*, 273:1516-1517, 1996). Finally, linkage analysis cannot be applied to the study of traits for which no large informative families are available. Typically, this will be the case in any attempt to identify trait-causing alleles involved in sporadic cases, such as alleles associated with positive or negative responses to drug treatment.

#### XI.B. Allele-Sharing methods

Whereas linkage analysis involves proposing a model to explain the inheritance pattern of phenotypes and genotypes in a pedigree, allele-sharing methods are not based on constructing a model, but rather on rejecting a model (see Lander and Schork, *Science*, 265, 2037-2048, 1994). More specifically, one tries to prove that the inheritance pattern of a chromosomal region is not consistent with random Mendelian segregation by showing that affected relatives inherit identical copies of the region more often than expected by chance. Because allele-sharing methods are nonparametric (that is, assume no model for the inheritance of the trait), they tend to be more useful for the analysis of complex traits than linkage analysis. Affected relatives should show excess allele sharing even in the presence of incomplete penetrance and polygenic inheritance. Allele-Sharing methods involve studying affected relatives in a pedigree to determine how often a particular copy of a chromosomal region is shared identical-by-descent (IBD), that is, is inherited from a common ancestor within the pedigree. The frequency of IBD sharing at a locus can then be compared with random expectation. Affected sib pair analysis is a well-known special case and is the simplest form of this method.

However, as allele-sharing methods analyze affected relatives, they tend to be of limited value in the genetic analysis of drug responses or in the analysis of side effects to treatments. This type of analysis is impractical in such cases due to the lack of availability of familial cases. In fact, the likelihood of having more than one individual in a family being exposed to the same drug at the same time is very low.

#### XI.C. Association Studies

The present invention comprises methods for identifying one or several genes among a set of candidate genes that are associated with a detectable trait using the biallelic markers of the present invention. In one embodiment the present invention comprises methods to detect an association between a biallelic marker allele or a biallelic marker haplotype and a trait. Further, the invention comprises methods to identify a trait causing allele in linkage disequilibrium with any biallelic marker allele of the present invention.

As described above, alternative approaches can be employed to perform association studies: genome-wide association studies, candidate region association studies and candidate gene association studies. In a preferred embodiment, the biallelic markers of the present invention are used to perform candidate gene association studies. The candidate gene analysis clearly provides a short-cut approach to the identification of genes and gene polymorphisms related to a particular trait when some information concerning the biology of the trait is available. Further, the biallelic markers of the present invention is incorporated in any map of genetic markers of the human genome in order to perform genome-wide association studies. Methods to generate a high-density map of biallelic markers has been described in US Provisional Patent application serial number 60/082,614. The biallelic markers of the present invention may further be incorporated in any map of a specific candidate region of the genome (a specific chromosome or a specific chromosomal region for example).

As mentioned above, association studies is conducted within the general population and are not limited to studies performed on related individuals in affected families. Linkage disequilibrium and association studies are extremely valuable as they permit the analysis of sporadic or multifactor traits. Moreover, association studies represent a powerful method for fine-scale mapping enabling much finer mapping of trait causing alleles than linkage studies. Studies based on pedigrees often only narrow the location of the trait causing allele. Association studies and Linkage Disequilibrium mapping methods using the biallelic markers of the present invention can therefore be used to refine the location of a trait causing allele in a candidate region identified by Linkage Analysis or by Allele-Sharing methods. Moreover, once a chromosome segment of interest has been identified, the presence of a candidate gene such as a candidate gene of the present invention, in the region of interest can provide a shortcut to the identification of the trait causing allele. Biallelic markers of the

present invention can be used to demonstrate that a candidate gene is associated with a trait. Such uses are specifically contemplated in the present invention and claims.

**1) Case-control populations (inclusion criteria)**

Association studies do not concern familial inheritance and do not involve the analysis of large family pedigrees but compare the prevalence of a particular genetic marker, or a set of markers, in case-control populations. They are case-control studies based on comparison of unrelated case (affected or trait positive) individuals and unrelated control (random or unaffected or trait negative) individuals. The control group is composed of individuals chosen randomly or of unaffected (trait negative) individuals, preferably the control group is composed of unaffected or trait negative individuals. Further, the control group is preferably both ethnically- and age-matched to the case population. In the following "trait positive population", "case population" and "affected population" are used interchangeably.

An important step in the dissection of complex traits using association studies is the choice of case-control populations (see Lander and Schork, *Science*, 265, 2037-2048, 1994). Narrowing the definition of the disease and restricting the patient population to extreme phenotypes allows one to work with a trait that is more nearly Mendelian in its inheritance pattern and more likely to be homogeneous (patients suffer from the disease for the same genetic reasons). Therefore, a major step in the choice of case-control populations is the clinical definition of a given trait or phenotype. Four criteria are often useful: clinical phenotype, age at onset, family history and severity. Preferably, in order to perform efficient and significant association studies, such as those described herein, the trait under study should preferably follow a bimodal distribution in the population under study, presenting two clear non-overlapping phenotypes (trait positive and trait negative). Nevertheless, even in the absence of such bimodal distribution (as may in fact be the case for more complex genetic traits), any genetic trait may still be analyzed by the association method proposed here by carefully selecting the individuals to be included in the trait positive and trait negative phenotypic groups. The selection procedure involves selecting individuals at opposite ends of the non-bimodal phenotype spectra of the trait under study, so as to include in these trait positive and trait negative populations individuals which clearly represent extreme, preferably non-overlapping phenotypes. This is particularly useful for continuous or quantitative traits (such as blood pressure for example). Selection of individuals at extreme ends of the trait distribution increases the ability to analyze these complex traits. The definition of the inclusion criteria for the case-control populations is an important aspect of association studies. The selection of those drastically different but relatively uniform phenotypes enables efficient comparisons in association studies and the possible detection of marked differences at the genetic level, provided that the sample sizes of the populations under study are significant enough.

Preferably, case-control populations to be included in association studies such as those proposed in the present invention consist of phenotypically homogeneous populations of individuals each representing 100% of the corresponding phenotype if the trait distribution is bimodal. If the trait distribution is non-bimodal, trait positive and trait negative populations consist of phenotypically uniform populations of individuals representing each between 1 and 98%, preferably between 1 and 80%, more preferably between 1 and 50%, and more preferably between 1 and 30%, most preferably between 1 and 20% of the total population under study, and selected among individuals exhibiting non-overlapping phenotypes. In some embodiments, the trait positive and trait negative groups consist of individuals exhibiting the extreme phenotypes within the studied population. The clearer the difference between the two trait phenotypes, the greater the probability of detecting an association with biallelic markers.

In preferred embodiments, a first group of between 50 and 300 trait positive individuals, preferably about 100 individuals, are recruited according to their phenotypes. A similar number of trait negative individuals are included in such studies.

In the present invention, typical examples of inclusion criteria include a diagnosis of cancer or prostate cancer or the evaluation of the response to anti-cancer or anti-prostate cancer agent or side effects to treatment with anti-cancer or anti-prostate cancer agents.

Suitable examples of association studies using biallelic markers including the biallelic markers of the present invention, are studies involving the following populations:

- a case population suffering from a form of cancer and a healthy unaffected control population, or
- a case population suffering from a form of prostate cancer and a healthy unaffected control population, or
- a case population treated with anticancer agents suffering from side-effects resulting from the treatment and a control population treated with the same agents showing no side-effects, or
- a case population treated with anti-prostate cancer agents suffering from side-effects resulting from the treatment and a control population treated with the same agents showing no side-effects, or
- a case population treated with anti-cancer agents showing a beneficial response and a control population treated with same agents showing no beneficial response, or
- a case population treated with anti-prostate cancer agents showing a beneficial response and a control population treated with same agents showing no beneficial response.

## **2) Determining the frequency of an allele in case-control populations**

Allelic frequencies of the biallelic markers in each of the populations can be determined using one of the methods described above under the in Section X. under the heading "Methods for genotyping an individual for biallelic markers", or any genotyping procedure suitable for this intended purpose. The frequency of a biallelic marker allele in a population can be determined by genotyping pooled samples or individual samples. One way to reduce the number of genotypings required is to



use pooled samples. A major obstacle in using pooled samples is in terms of accuracy and reproducibility for determining accurate DNA concentrations in setting up the pools. Genotyping individual samples provides higher sensitivity, reproducibility and accuracy and; is the preferred method used in the present invention. Preferably, each individual is genotyped separately and simple  
5 gene counting is applied to determine the frequency of an allele of a biallelic marker or of a genotype in a given population.

### 3) Determining the frequency of a haplotype in case-control populations

The gametic phase of haplotypes is usually unknown when diploid individuals are heterozygous at more than one locus. Different strategies for inferring haplotypes is used to partially  
10 overcome this difficulty (see Excoffier L. and Slatkin M., *Mol. Biol. Evol.*, 12(5): 921-927, 1995). One possibility is that the multiple-site heterozygous diploids can be eliminated from the analysis, keeping only the homozygotes and the single-site heterozygote individuals, but this approach might lead to a possible bias in the sample composition and the underestimation of low-frequency haplotypes. Another possibility is that single chromosomes can be studied independently, for  
15 example, by asymmetric PCR amplification (see Newton et al., *Nucleic Acids Res.*, 17:2503-2516, 1989; Wu et al., *Proc. Natl. Acad. Sci. USA*, 86:2757, 1989) or by isolation of single chromosome by limit dilution followed by PCR amplification (see Ruano et al., *Proc. Natl. Acad. Sci. USA*, 87:6296-6300, 1990). Further, multiple haplotypes can sometimes be inferred using genealogical information in families (Perlin et al., *Am. J. Hum. Genet.*, 55:777-787, 1994). A sample is haplotyped for  
20 sufficiently close biallelic markers by double PCR amplification of specific alleles (Sarkar, G. and Sommer S.S., *Biotechniques*, 1991). These approaches are not entirely satisfying either because of their technical complexity, the additional cost they entail, their lack of generalization at a large scale, or the possible biases they introduce. To overcome these difficulties, an algorithm based on Hardy-Weinberg equilibrium (random mating) to infer the phase of PCR-amplified DNA genotypes  
25 introduced by Clark A.G. (*Mol. Biol. Evol.*, 7:111-122, 1990) is used. Briefly, the principle is to start filling a preliminary list of haplotypes present in the sample by examining unambiguous individuals, that is, the complete homozygotes and the single-site heterozygotes. Then other individuals in the same sample are screened for the possible occurrence of previously recognized haplotypes. For each positive identification, the complementary haplotype is added to the list of recognized haplotypes,  
30 until the phase information for all individuals is either resolved or identified as unresolved. This method assigns a single haplotype to each multiheterozygous individual, whereas several haplotypes are possible when there are more than one heterozygous site. Any other method known in the art to determine the frequency of a haplotype in a population is used. Preferably, an expectation-maximization (EM) algorithm (Dempster et al., *J. R. Stat. Soc.*, 39B:1-38, 1977) leading to maximum-  
35 likelihood estimates of haplotype frequencies under the assumption of Hardy-Weinberg proportions is used (see Excoffier L. and Slatkin M., *Mol. Biol. Evol.*, 12(5): 921-927, 1995). The EM algorithm is

used to estimate haplotype frequencies in the case when only genotype data from unrelated individuals are available. The EM algorithm is a generalized iterative maximum-likelihood approach to estimation that is useful when data are ambiguous and/or incomplete. The EM algorithm is used to resolve heterozygotes into haplotypes. Haplotype estimations are further described below under the heading "Statistical methods".

#### 4) Genetic Analysis based on Linkage Disequilibrium

Linkage disequilibrium is the non-random association of alleles at two or more loci and represents a powerful tool for genetic mapping of complex traits (see Jorde L.B., *Am. J. Hum. Genet.*, 56:11-14, 1995). Biallelic markers, because they are densely spaced in the human genome and can be genotyped in large numbers, are particularly useful in genetic analysis based on linkage disequilibrium.

When a disease mutation is first introduced into a population (by a new mutation or the immigration of a mutation carrier), it necessarily resides on a single chromosome and thus on a single "background" or "ancestral" haplotype of linked markers. Consequently, there is complete disequilibrium between these markers and the disease mutation: one finds the disease mutation only in the presence of a specific set of marker alleles. Through subsequent generations recombinations occur between the disease mutation and these marker polymorphisms, and the disequilibrium gradually dissipates. The pace of this dissipation is a function of the recombination frequency, so the markers closest to the disease gene will manifest higher levels of disequilibrium than those that are further away. When not broken up by recombination, "ancestral" haplotypes and linkage disequilibrium between marker alleles at different loci can be tracked not only through pedigrees but also through populations.

The pattern or curve of disequilibrium between disease and marker loci will exhibit a single maximum that occurs at the disease locus. Consequently, the amount of linkage disequilibrium between a disease allele and closely linked genetic markers may yield valuable information regarding the location of the disease gene. For fine-scale mapping of a disease locus, it is useful to have some knowledge of the patterns of linkage disequilibrium that exist between markers in the studied region. As mentioned above the mapping resolution achieved through the analysis of linkage disequilibrium is much higher than that of linkage studies. The high density of biallelic markers combined with linkage disequilibrium analysis provide powerful tools for fine-scale mapping. Different methods to calculate linkage disequilibrium are described below under the heading "Statistical Methods". Moreover, association studies as a method of mapping genetic traits rely on the phenomenon of linkage disequilibrium.

#### 3) Association studies

As mentioned above, the occurrence of pairs of specific alleles at different loci on the same chromosome is not random, and the deviation from random is called linkage disequilibrium. If a

specific allele in a given gene is directly involved in causing a particular trait, its frequency will be statistically increased in an affected (trait positive) population when compared to the frequency in a trait negative population or in a random control population. As a consequence of the existence of linkage disequilibrium, the frequency of all other alleles present in the haplotype carrying the trait-causing allele will also be increased in trait positive individuals compared to trait negative individuals or random controls. Therefore, association between the trait and any allele (specifically a biallelic marker allele) in linkage disequilibrium with the trait-causing allele will suffice to suggest the presence of a trait-related gene in that particular allele's region. Association studies focus on population frequencies. Case-control populations can be genotyped for biallelic markers to identify associations that narrowly locate a trait causing allele. Moreover, any marker in linkage disequilibrium with one given marker associated with a trait will be associated with the trait. Linkage disequilibrium allows the relative frequencies in case-control populations of a limited number of genetic polymorphisms (specifically biallelic markers) to be analyzed as an alternative to screening all possible functional polymorphisms in order to find trait-causing alleles. Association studies compare the frequency of marker alleles in unrelated case-control populations, and represent powerful tools for the dissection of complex traits.

#### Association analysis

The general strategy to perform association studies using biallelic markers derived from a candidate gene is to scan two groups of individuals (case-control populations) in order to measure and statistically compare the allele frequencies of the biallelic markers of the present invention in both groups.

If a statistically significant association with a trait is identified for at least one or more of the analyzed biallelic markers, one can assume that: either the associated allele is directly responsible for causing the trait (the associated allele is the trait causing allele), or more likely the associated allele is in linkage disequilibrium with the trait causing allele. The specific characteristics of the associated allele with respect to the candidate gene function usually gives further insight into the relationship between the associated allele and the trait (causal or in linkage disequilibrium). If the evidence indicates that the associated allele within the candidate gene is most probably not the trait causing allele but is in linkage disequilibrium with the real trait causing allele, then the trait causing allele can be found by sequencing the vicinity of the associated marker.

Association studies are usually run in two successive steps. In a first phase, the frequencies of a reduced number of biallelic markers from one or several candidate genes are determined in the trait positive and trait negative populations. In a second phase of the analysis, the identity of the candidate gene and the position of the genetic loci responsible for the given trait is further refined using a higher density of markers from the relevant gene. However, if the candidate gene under study is relatively

small in length, as it is the case for many of the candidate genes analyzed included in the present invention, a single phase is sufficient to establish significant associations.

#### **Haplotype analysis**

As described above, when a chromosome carrying a disease allele first appears in a population as a result of either mutation or migration, the mutant allele necessarily resides on a chromosome having a unique set of linked markers: the ancestral haplotype. This haplotype can be tracked through populations and its statistical association with a given trait can be analyzed. The statistical power of association studies is increased by complementing single point (allelic) association studies with multi-point association studies also called haplotype studies. Thus, a haplotype association study allows one to define the frequency and the type of the ancestral carrier haplotype. A haplotype analysis is important in that it increases the statistical significance of an analysis involving individual markers. Indeed, by performing an association study with a set of biallelic markers, it increases the value of the results obtained through the study, allowing false positive and/or negative data that may result from the single marker studies to be eliminated.

In a first stage of a haplotype frequency analysis, the frequency of the possible haplotypes based on various combinations of the identified biallelic markers of the invention is determined. The haplotype frequency is then compared for distinct populations of trait positive and control individuals. The number of trait positive individuals which should be subjected to this analysis to obtain statistically significant results usually ranges between 30 and 300, with a preferred number of individuals ranging between 50 and 150. The same considerations apply to the number of random control or unaffected individuals used in the study. The results of this first analysis provide haplotype frequencies in case-control populations, the relative risk for an individual carrying a given haplotype of being affected with the given trait under study and the estimated p value for each evaluated haplotype.

#### **Interaction Analysis**

The biallelic markers of the present invention may also be used to identify patterns of biallelic markers associated with detectable traits resulting from polygenic interactions. The analysis of genetic interaction between alleles at unlinked loci requires individual genotyping using the techniques described herein. The analysis of allelic interaction among a selected set of biallelic markers with appropriate level of statistical significance can be considered as a haplotype analysis, similar to those described in further details within the present invention. Preferably, genotyping typing is performed using the microsequencing technique.

Methods to test for association between a trait and a biallelic marker allele or a haplotype of biallelic marker alleles are described below.

#### **XI.D. Statistical methods**

In general, any method known in the art to test whether a trait and a genotype show a statistically significant correlation is used.

#### Methods to estimate haplotype frequencies in a population

As described above, when genotypes are scored, it is often not possible to distinguish heterozygotes so that haplotype frequencies cannot be easily inferred. When the gametic phase is not known, haplotype frequencies can be estimated from the multilocus genotypic data. Any method known to person skilled in the art can be used to estimate haplotype frequencies (see Lange K., *Mathematical and Statistical Methods for Genetic Analysis*, Springer, New York, 1997; Weir, B.S., *Genetic data Analysis II: Methods for Discrete population genetic Data*, Sinauer Assoc., Inc., Sunderland, MA, USA, 1996) Preferably, maximum-likelihood haplotype frequencies are computed using an Expectation- Maximization (EM) algorithm (see Dempster et al., *J. R. Stat. Soc.*, 39B:1-38, 1977; Excoffier L. and Slatkin M., *Mol. Biol. Evol.*, 12(5): 921-927, 1995). This procedure is an iterative process aiming at obtaining maximum-likelihood estimates of haplotype frequencies from multi-locus genotype data when the gametic phase is unknown. Haplotype estimations are usually performed by applying the EM algorithm using for example the EM-HAPLO program (Hawley M.E. et al., *Am. J. Phys. Anthropol.*, 18:104, 1994) or the Arlequin program (Schneider et al., *Arlequin: a software for population genetics data analysis*, University of Geneva, 1997). The EM algorithm is a generalized iterative maximum likelihood approach to estimation and is briefly described below.

In the following part of this text, phenotypes will refer to multi-locus genotypes with unknown phase. Genotypes will refer to known-phase multi-locus genotypes.

Suppose a sample of N unrelated individuals typed for K markers. The data observed are the unknown-phase K-locus phenotypes that can be categorized in F different phenotypes. Suppose that we have H underlying possible haplotypes (in case of K biallelic markers,  $H=2^K$ ).

For phenotype j, suppose that  $c_j$  genotypes are possible. We thus have the following equation

$$P_j = \sum_{i=1}^{c_j} pr(genotype_i) = \sum_{i=1}^{c_j} pr(h_k, h_l) \quad \text{Equation 1}$$

where  $P_j$  is the probability of the phenotype j,  $h_k$  and  $h_l$  are the two haplotypes constituent the genotype i. Under the Hardy-Weinberg equilibrium,  $pr(h_k, h_l)$  becomes :

$$pr(h_k, h_l) = pr(h_k)^2 \text{ if } h_k = h_l, pr(h_k, h_l) = 2pr(h_k).pr(h_l) \text{ if } h_k \neq h_l. \quad \text{Equation 2}$$

The successive steps of the E-M algorithm can be described as follows:

Starting with initial values of the of haplotypes frequencies, noted,  $p_1^{(0)}, p_2^{(0)}, \dots, p_T^{(0)}$ .

these initial values serve to estimate the genotype frequencies (Expectation step) and then estimate

another set of haplotype frequencies (Maximization step):  $p_1^{(1)}, p_2^{(1)}, \dots, p_T^{(1)}$ .

these two steps are iterated until change in the sets of haplotypes frequency are very small.

A stop criterion can be that the maximum difference between haplotype frequencies between two iterations is less than  $10^{-7}$ . These values can be adjusted according to the desired precision of estimations.

In detail, at a given iteration  $s$ , the Expectation step consists in calculating the genotypes frequencies by the following equation:

$$\begin{aligned} pr(genotype_i)^{(s)} &= pr(phenotype_j) \cdot pr(genotype_i | phenotype_j)^{(s)} \\ &= \frac{n_j}{N} \cdot \frac{pr(h_k, h_l)^{(s)}}{P_j^{(s)}} \end{aligned} \quad \text{Equation 3}$$

where genotype  $i$  occurs in phenotype  $j$ , and where  $hk$  and  $hl$  constitute genotype  $i$ . Each probability are derived according to equations 1 and 2 above.

Then the Maximization step simply estimates another set of haplotype frequencies given the genotypes frequencies. This approach is also known as gene-counting method (Smith, *Ann. Hum. Genet.*, 21:254-276, 1957).

$$p_i^{(s+1)} = \frac{1}{2} \sum_{j=1}^F \sum_{i=1}^{c_j} \delta_{it} \cdot pr(genotype_i)^{(s)} \quad \text{Equation 4}$$

where  $\delta_{it}$  is an indicator variable which count the number of time haplotype  $t$  in genotype  $i$ . It takes the values of 0, 1 or 2.

To ensure that the estimation finally obtained are the maximum-likelihood estimations several values of departures are required. The estimations obtained are compared and if they differ the estimations leading to the best likelihood are kept. The term "haplotype determination method" is used to refer to all methods for determinin haplotypes known in the art including expectation-maximization algorithms.

## 20 Methods to calculate linkage disequilibrium between markers

A number of methods can be used to calculate linkage disequilibrium between any two genetic positions, in practice, linkage disequilibrium is measured by applying a statistical association test to haplotype data taken from a population.

Linkage disequilibrium between any pair of biallelic markers comprising at least one of the biallelic markers of the present invention ( $M_i, M_j$ ) can be calculated for every allele combination ( $M_{i1}, M_{j1}; M_{i1}, M_{j2}; M_{i2}, M_{j1}$  and  $M_{i2}, M_{j2}$ ), according to the Piazza formula :

$$\Delta M_{ik}, M_{jl} = \sqrt{\theta 4 - \sqrt{(\theta 4 + \theta 3)(\theta 4 + \theta 2)}}, \text{ where :}$$

$\theta 4 = - -$  = frequency of genotypes not having allele  $k$  at  $M_i$  and not having allele  $l$  at  $M_j$

$\theta 3 = - +$  = frequency of genotypes not having allele  $k$  at  $M_i$  and having allele  $l$  at  $M_j$

$\theta 2 = + -$  = frequency of genotypes having allele  $k$  at  $M_i$  and not having allele  $l$  at  $M_j$

Linkage disequilibrium (LD) between pairs of biallelic markers ( $M_i, M_j$ ) can also be calculated for every allele combination ( $M_{i1}, M_{j1}; M_{i1}, M_{j2}; M_{i2}, M_{j1}$  and  $M_{i2}, M_{j2}$ ), according to the maximum-likelihood estimate (MLE) for delta (the composite linkage disequilibrium coefficient), as described by Weir (B.S. Weir, *Genetic Data Analysis*, Sinauer Ass. Eds, 1996). This formula allows linkage disequilibrium between alleles to be estimated when only genotype, and not haplotype, data are available. This LD composite test makes no assumption for random mating in the sampled population, and thus seems to be more appropriate than other LD tests for genotypic data.

Another means of calculating the linkage disequilibrium between markers is as follows. For a couple of biallelic markers,  $M_i (a/b_i)$  and  $M_j (a/b_j)$ , fitting the Hardy-Weinberg equilibrium, one can estimate the four possible haplotype frequencies in a given population according to the approach described above.

The estimation of gametic disequilibrium between  $a_i$  and  $a_j$  is simply:

$$D_{aiaj} = pr(haplotype(a_i, a_j)) - pr(a_i) \cdot pr(a_j).$$

Where  $pr(a_i)$  is the probability of allele  $a_i$  and  $a_j$  is the probability of allele  $a_j$ . and where  $pr(haplotype(a_i, a_j))$  is estimated as in Equation 3 above.

For a couple of biallelic marker only one measure of disequilibrium is necessary to describe the association between  $M_i$  and  $M_j$ .

Then a normalized value of the above is calculated as follows:

$$\begin{aligned} D'_{aiaj} &= D_{aiaj} / \max (pr(a_i) \cdot pr(a_j), pr(b_i) \cdot pr(b_j)) && \text{with } D_{aiaj} < 0 \\ D'_{aiaj} &= D_{aiaj} / \max (pr(b_i) \cdot pr(a_j), pr(a_i) \cdot pr(b_j)) && \text{with } D_{aiaj} > 0 \end{aligned}$$

The skilled person will readily appreciate that other LD calculation methods can be used without undue experimentation.

Linkage disequilibrium among a set of biallelic markers having an adequate heterozygosity rate can be determined by genotyping between 50 and 1000 unrelated individuals, preferably between 75 and 200, more preferably around 100.

#### Testing for association

Methods for determining the statistical significance of a correlation between a phenotype and a genotype, in this case an allele at a biallelic marker or a haplotype made up of such alleles, is determined by any statistical test known in the art and with any accepted threshold of statistical significance being required. The application of particular methods and thresholds of significance are well within the skill of the ordinary practitioner of the art.

Testing for association is performed by determining the frequency of a biallelic marker allele in case and control populations and comparing these frequencies with a statistical test to determine if there is a statistically significant difference in frequency which would indicate a correlation between the trait and the biallelic marker allele under study. Similarly, a haplotype analysis is performed by estimating the frequencies of all possible haplotypes for a given set of biallelic markers in case and

control populations, and comparing these frequencies with a statistical test to determine if there is a statistically significant correlation between the haplotype and the phenotype (trait) under study. Any statistical tool useful to test for a statistically significant association between a genotype and a phenotype is used. Preferably the statistical test employed is a chi square test with one degree of freedom. A P-value is calculated (the P-value is the probability that a statistic as large or larger than the observed one would occur by chance).

#### Statistical significance

In preferred embodiments, significance for diagnosis purposes, either as a positive basis for further diagnostic tests or as a preliminary starting point for early preventive therapy, the p value related to a biallelic marker association is preferably about  $1 \times 10^{-2}$  or less, more preferably about  $1 \times 10^{-4}$  or less, for a single biallelic marker analysis and about  $1 \times 10^{-3}$  or less, still more preferably  $1 \times 10^{-6}$  or less and most preferably of about  $1 \times 10^{-8}$  or less, for a haplotype analysis involving several markers. These values are believed to be applicable to any association studies involving single or multiple marker combinations.

The skilled person can use the range of values set forth above as a starting point in order to carry out association studies with biallelic markers of the present invention. In doing so, significant associations between the biallelic markers of the present invention and cancer and prostate cancer can be revealed and used for diagnosis and drug screening purposes.

Using the method described above and evaluating the associations for single marker alleles or for haplotypes permits an estimation of the risk a corresponding carrier has to develop a given trait, and particularly in the context of the present invention, a disease, preferably cancer, more preferably prostate cancer. Significance thresholds of relative risks are to be adapted to the reference sample population used.

In this regard, among all the possible marker combinations or haplotypes which are evaluated to determine the significance of their association with a given trait, for example a form of cancer or prostate cancer, a response to treatment with anti-cancer or anti-prostate cancer agents or side effects related to treatment with anti-cancer or anti-prostate cancer agents, it is believed that those displaying a coefficient of relative risk above 1, preferably about 5 or more, preferably of about 7 or more are indicative of a "significant risk" for the individuals carrying the identified haplotype to develop the given trait. It is difficult to evaluate accurately quantified boundaries for the so-called "significant risk". Indeed, and as it has been demonstrated previously, several traits observed in a given population are multifactorial in that they are not only the result of a single genetic predisposition but also of other factors such as environmental factors or the presence of further, apparently unrelated, haplotype associations. Thus, the evaluation of a significant risk must take these parameters into consideration in order to, in a certain manner, weigh the potential importance of external parameters in the development of a given trait. Without wishing to be bound to any invariable model or theory



based on the above statistical analyses, the inventors believe that a "significant risk" to develop a given trait is evaluated differently depending on the trait under consideration.

It will of course be understood by practitioners skilled in the treatment or diagnosis of cancer and prostate cancer that the present invention does not intend to provide an absolute identification of  
5 individuals who could be at risk of developing a particular form of cancer or who will or will not respond or exhibit side effects to treatment with anti-cancer or anti-prostate cancer agents but rather to indicate a certain degree or likelihood of developing a disease or of observing in a given individual a response or a side effect to treatment with a particular agent or set of agents.

However, this information is extremely valuable as it can, in certain circumstances, be used to  
10 initiate preventive treatments or to allow an individual carrying a significant haplotype to foresee warning signs such as minor symptoms. In the case of cancer, the knowledge of a potential predisposition, even if this predisposition is not absolute, might contribute in a very significant manner to treatment, or allow for suggestions in changes in diet or the reduction of risky behaviors, e.g. smoking. Similarly, a diagnosed predisposition to a potential side effect could immediately direct  
15 the physician toward a treatment, for which such side effects have not been observed during clinical trials.

#### **Phenotypic randomization**

In order to confirm the statistical significance of the first stage haplotype analysis described above, it might be suitable to perform further analyses in which genotyping data from case-control  
20 individuals are pooled and randomized with respect to the trait phenotype. Each individual genotyping data is randomly allocated to two groups which contain the same number of individuals as the case-control populations used to compile the data obtained in the first stage. A second stage haplotype analysis is preferably run on these artificial groups, preferably for the markers included in the haplotype of the first stage analysis showing the highest relative risk coefficient. This experiment  
25 is reiterated between 50 and 200 times, preferably between 75 and 125 times. The repeated iterations allow the determination of the percentage of obtained haplotypes with a significant p-value level below about  $1 \times 10^{-3}$ .

#### Example 24

##### Detailed Association Studies

30 The initial association studies between the 8p23 locus and prostate cancer described in Section I.D. were repeated at a higher level of sophistication.

##### Collection of DNA samples from affected and non-affected individuals

Prostate cancer patients were recruited according to clinical inclusion criteria based on pathological or radical prostatectomy records as described above in Section I. However, the pool of  
35 individuals suffering from prostate cancer described in Section I was augmented from the original 185 individuals to a range of between 275 and 491 individuals depending on the marker tested. Similarly,

the control pool of non-diseased individuals described in Section I was augmented from the original 104 individuals to a range of between 130 and 313 individuals depending on the marker tested.

#### Genotyping Affected and Control Individuals

As for Section I.D., allelic frequencies of the biallelic markers in each population were  
5 determined by performing microsequencing reactions on amplified fragments obtained by genomic PCR performed on the DNA samples from each individual as described in Example 5.

#### Association Studies

Association results were obtained using markers spanning a 650 kb region of the 8p28 locus around PG1 both using single point analysis and haplotyping studies. See Figure 16. As compared  
10 with the earlier representation of the initial association results for this region shown in Figure 2, Figure 16 is to scale, since the entire region has now been sequenced. In addition, more markers were generated around the association peak in the area of PG1; each of which has been tested in single point analysis (hence the density of data within this subregion). The haplotyping curve in Figure 16 represents, for each marker considered, the maximum p-value for haplotypes obtained using this  
15 marker and any number from all markers harbored by the same BAC and being in Hardy Weinberg Disequilibrium with said marker.

The data presented in Figure 16 shows a strong association between this specific region within 8p23 locus, especially in the area that has been identified as being the PG1 gene, and prostate cancer. The maximum p-value in single point analysis, for the PG1 sub-region, is  $3.10^{-3}$ , while outside  
20 of the PG1 subregion, most of the p-values obtained for single point associations are less significant than  $1.10^{-1}$ . The maximum p-value obtained for haplotyping studies is the one obtained for a marker inside PG1's BAC, and equals  $3.10^{-6}$ .

Figure 17 is a graph showing an enlarged view of the single point association results within a 160 kb region comprising the PG1 gene. Markers involved in this enlargement were all located on  
25 BAC B0463F01 (see Figure 16), except marker 4-14, which lies in very close proximity, on BAC B0189E08. Figure 17 shows all of the markers which made up the maximum haplotype shown in Figure 16. Some of these markers were later revealed to lie within the promoter, exonic or intronic regions of the PG1 gene. The markers outside the gene were all informative biallelic markers with a least frequent allele present at a frequency of more than 20%, while markers within the gene were a  
30 mix of such informative markers and markers whose least frequent allele's frequency is less than 20%. These data confirm and narrow the previous peak of association values seen in Figure 16, to a 40 kb harboring the PG1 gene. Significant associations are obtained for markers starting at the promoter site with marker No. 99-1485, and ending at the 3' UTR site with marker No. 5-66.

Figure 18A is a graph showing an enlarged view of the single point association results of 40  
35 kb within the PG1 gene. These data confirm that seven markers within the PG1 gene have one allele associated with prostate cancer, with p-values all similar and more significant than  $1.10^{-2}$ , specifically

markers 99-622 ; 4-77 ; 4-71 ; 4-73 ; 99-598 ; 99-576 ; 4-66. Figure 18B is a table listing the location of markers within PG1 gene, the two possible alleles at each site. For each marker, the disease-associated allele is indicated first ; its frequencies in cases and controls as well as the difference between both are shown ; the odd-ratio and the p-value of each individual marker association are also shown.

The data in Figures 17, 18A, and 18B demonstrate that the markers in the PG1 gene have an association with prostate cancer that is valid, and exhibits similar significance values, regardless whether the considered cases are sporadic or familial cases. Therefore, some PG1 alleles must be general risk factors for any type of prostate cancer, whether familial or sporadic. The fact that several p-values for associated alleles are around  $1.10^{-2}$  suggests that all these markers are in linkage disequilibrium to one another, and can all be used individually to assess PG1 associated prostate cancer susceptibility risk. The prostate cancer associated alleles of the 7 markers discussed above, all exhibit an odd-ratio of about 1.5, which means for each of them that an individual carrying such allele has 1.5 more chances to be susceptible to prostate cancer than not.

In order to confirm the significance of the association results found for markers on the BAC harboring PG1, a novel statistical method was performed as described in provisional patent application serial no. 60/107,986, filed November 10, 1998.

#### Haplotype analysis

The results of a haplotype analysis study using 4 markers (marker Nos. 4-14, 99-217, 4-66 and 99-221) within the 160 kb region shown in Figure 17 are shown in Figure 19A. These 4 markers have each been shown to be strongly associated with prostate cancer, i.e. with p-values more significant than  $1.10^{-3}$  on approximately 150 cases and 130 controls. All haplotypes using 2, 3, or 4 markers among the 4 above cited were analyzed using 491 case patients and 317 control individuals. Figure 19A shows the most significant haplotypes obtained, as well as the individual odd-ratios for each. Haplotype 11 is the most significant (p-value of ca.  $3.10^{-6}$ ), and is related to haplotype 5, shown in Figure 4 in that three of the four marker alleles (4-14 C, 99-217 T and 99-221 A) are common to both haplotypes, and both cover a similar region. Differences in p-values are explained both by the addition of markers and of more case or control individuals. Haplotype 11 has an highly informative odd-ratio (of above 3) ; it is present in 3% of the controls and almost 10% of the cases.

Figure 19B is a table showing the segmented haplotyping results according to the age of the subjects, and whether the prostate cancer cases were sporadic or familial, using the same markers 4 markers and the same individuals as were used to generate the results in Figure 19A. Figure 19B shows equivalent results for all segments of the population analyzed, demonstrating that the PG1 associated alleles are general risk factors for prostate cancer, regardless of the age of onset of the disease.

The haplotyping results and odd ratios for all of the combinations of the 7 markers (99-622; 4-77; 4-71; 4-7 ; 99-598; 99-576; and 4-66) within PG1 gene that were shown in Figure 18 to have p-values more significant than  $1 \times 10^{-2}$  were computed. A portion of these data are shown in Figure 20. All of the 2-, 3-, 4-, 5-, 6- and 7-marker haplotypes were tested. Figure 20 identifies for each x-marker  
 5 haplotype category, the most significant haplotype. Among all these, the most significant haplotype is the two-marker haplotype 1, which shows a p-value of approximately  $6.10^{-5}$ , with an odd ratio of 2. The frequency of haplotype 1 among the control individuals is 15%, while it is 26% among the case patients. It is worth noting that these frequencies are very similar for all haplotypes presented on Figure 20. It will thus be sufficient to test this two marker haplotype for prognosis/diagnosis on risk  
 10 patients, as opposed to having a more complex test of a haplotype comprising 3 or more makers.

Finally, Figure 21 is a graph showing the distribution of statistical significance, as measured by Chi-square values, for each series of possible x-marker haplotypes, ( $x = 2, 3$  or  $4$ ) using all of the 19 markers found in PG1 gene. These data confirm that testing 2-marker haplotypes within PG1 is sufficient because the testing 3- or 4-marker haplotypes does not increase the statistical relevance of  
 15 the analysis.

#### Example 25

##### Attributable Risk

Attributable risk describes the proportion of individuals in a population exhibiting a phenotype due to exposure to a particular factor. For further discussion of attributable risk values, see  
 20 *Holland, Bart K., Probability without Equations – Concepts for Clinicians*; The Johns Hopkins University Press, pp. 88-90. In the present case the phenotype examined was prostate cancer, and the exposure was either one single allele of an individual PG1-related marker, or a haplotype thereof in an individual's genome.

The formula used for calculating attributable risk values in the present study was the following:

25  $AR = P_E (RR-1) / [P_E (RR-1)+1]$ , where:

AR was the attributable risk of allele or haplotype ;

$P_E$  was the frequency of exposure to allele or haplotype within the population at large, in the present study a random male Caucasian population ; and

RR was the relative risk, in the present study relative risk is approximated with the odd-ratio,  
 30 because of the relatively low incidence of prostate cancer in populations at large (values for the odd ratios are found in Figures 18B and 20).

In this case,  $P_E$  was estimated using a dominant transmission model for prostate cancer:

$P_E = (N_{AA} + N_{AB}) / N$ , where:

$N_{AA}$  was the number of homozygous individuals harboring the disease associated allele or  
 35 haplotype within a given random population, and  $N_{AB}$  was the number of heterozygous individuals in said random population.  $N_{AA}$  and  $N_{AB}$  were calculated using the allele frequencies in the random

population as indicated in Figures 18B and 20, and N was the number of individuals in total random population.

We calculated the attributable risks of disease-associated alleles for markers within PG1 gene and presented these results in Figure 18B. In Figure 20, the attributable risk for the two-marker  
 5 haplotypes present in the figure as shown as well. These data demonstrate that disease-associated alleles of PG1 are present in approximately 20% of prostate cancer patients in the Caucasian population at large, and therefor represent prognostic tools of significant value.

#### SEQUENCE LISTING FREE TEXT

10 The following free text appears in the accompanying Sequence Listing:

identification method Proscan

potential start codon

exon1

Tyr phos

15 upstream amplification primer

polymorphic fragment

polymorphic base

downstream amplification primer

complement

20 upstream amplification primer 99-217-PU, extracted from SEQ ID1 34216 34234

Klein, Kanehisa and DeLisi identification method, potential helix

Eisenberg, Schwarz, Komarony, Wall identification method, potential helix

Prosite match

potential Tyrosine kinase site, Prosite match

25 potential caseine kinase II site, Prosite match

potential Leucine zipper site, Prosite match

potential site, Prosite match

potential protein kinase C, Prosite match

potential cAMP and cGMP dependant protein kinase site, Prosite match

30 primer oligonucleotide

box2 from SEQID4, present in AF003136, P33333, P26647, U89336, U56417,

AB005623

box2 from Z72511

box3 from SEQID4, present in AF003136

35 potential microsequencing oligo

complement potential microsequencing oligo

polymorphic fragment 4-77, extracted from SEQ ID1 12057 12103

polymorphic fragment 99-123, variant version of SEQ ID21

base A ; G in SEQ ID22

downstream amplification primer 99-217-RP, extracted from SEQ ID1 34625 34645

5 complement

polymorphic base C in PG1 (13680) SEQ ID1

stop codon

potential

amplification oligonucleotide

10 sequencing oligonucleotide

Box II

Box III

Box I

upstream amplification primer for SEQ 188, SEQ 265, SEQ 189, SEQ 266

15 downstream amplification primer for SEQ 185, SEQ 262, SEQ 186, SEQ 263, SEQ 187, SEQ

264

microsequencing oligo for 4-20-149.mis1

Although this invention has been described in terms of certain preferred embodiments, other  
20 embodiments which will be apparent to those of ordinary skill in the art in view of the disclosure  
herein are also within the scope of this invention. Accordingly, the scope of the invention is intended  
to be defined only by reference to the appended claims.

Table 1

marker	polymorphism	most frequent	less frequent	cases		controls	
				p *	q **	p *	q **
99-123	C/T	C	T	0,65	0,35	0,7	0,3
4-26	A/G	A	G	0,61	0,39	0,55	0,45
4-14	C/T	C	T	0,65	0,35	0,59	0,41
4-77	C/G	C	G	0,67	0,33	0,76	0,24
99-217	C/T	C	T	0,69	0,31	0,77	0,23
4-67	C/T	C	T	0,74	0,26	0,84	0,16
99-213	A/G	A	G	0,55	0,45	0,62	0,38
99-221	C/A	C	A	0,43	0,57	0,43	0,57
99-135	A/G	A	G	0,75	0,25	0,7	0,3

\*: frequency of most frequent base within each sub-population

\*\* : frequency of least frequent base within each sub-population ( $p+q=1$ )

standard deviations ~0,023 to 0,031 for controls

standard deviations ~0,018 to 0,021 for cases

WHAT IS CLAIMED IS:

1. A recombinant, purified or isolated polynucleotide comprising a mammalian PG1 gene, cDNA, complement thereof, or fragment thereof having at least 10 nucleotides in length.
- 5 2. The polynucleotide according to claim 1, wherein said mammalian PG1 gene or cDNA is human or mouse.
3. The polynucleotide according to claim 2, wherein the polynucleotide is selected from  
10 SEQ ID NOS: 3, 69, 112-124, 179, and 182-184.
4. A polynucleotide selected from SEQ ID NOS: 185-578.
5. A purified or isolated polypeptide comprising a mammalian PG1 protein, or fragment  
15 thereof having at least 8 amino acids in length.
6. The polypeptide according to claim 5, wherein said mammalian PG1 protein is human or mouse.
- 20 7. The polypeptide according to claim 6, wherein said polypeptide is selected from SEQ ID NOS: 4, 5, 70, 74, and 125-136.
8. The polypeptide according to claim 5, wherein said polypeptide consists of said mammalian PG1 protein, or fragment thereof having at least 8 amino acids in length.
- 25 9. A polynucleotide comprising a nucleic acid sequence encoding a polypeptide according to claim 8.
10. An antibody composition capable of selectively binding to an epitope-containing  
30 fragment of a polypeptide according to claim 8, wherein said antibody is either polyclonal or monoclonal.
11. A vector comprising a polynucleotide according to any one of claims 1, 4, and 9.
- 35 12. A host cell comprising a polynucleotide according to claim 11.



- 13 A nonhuman host animal or mammal comprising a vector according to claim 11.
14. A mammalian host cell comprising a PG1 gene disrupted by homologous recombination with a knock out vector.
- 5 15. A nonhuman host mammal comprising a PG1 gene disrupted by homologous recombination with a knock out vector.
- 10 16. A polynucleotide according to any one of claims 1, 4, and 9, further comprising a label.
17. A polynucleotide according to any one of claims 1, 4, and 9, attached to a solid support.
- 15 18. A random or addressable array of polynucleotides comprising at least one polynucleotide according to any one of claims 1, 4, and 9.
19. A method of determining whether an individual is at risk of developing cancer or prostate cancer, or whether said individual suffers from cancer or prostate cancer as a result of a  
20 mutation in the PG1 gene comprising:  
obtaining a nucleic acid sample from said individual; and  
determining whether the nucleotides present at one or more PG1-related biallelic marker are indicative of a risk of developing cancer or prostate cancer or indicative of cancer or prostate cancer resulting from a mutation in the PG1 gene.
- 25 20. A method of determining whether an individual is at risk of developing cancer or prostate cancer or whether said individual suffers from cancer or prostate cancer as a result of a mutation in the PG1 gene comprising:  
obtaining a nucleic acid sample from said individual; and  
30 determining whether the nucleotides present at one or more PG1-related biallelic marker are indicative of a risk of developing cancer or prostate cancer or indicative of cancer or prostate cancer resulting from a mutation in the PG1 gene.
21. A method according to either one of claims 19 and 20, wherein said PG1-related  
35 biallelic is a PG1-related biallelic markers positioned in SEQ ID NO: 179; a PG1-related biallelic marker selected from the group consisting of 99-1485/251, 99-622/95, 99-619/141, 4-76/222, 4-

77/151, 4-71/233, 4-72/127, 4-73/134, 99-610/250, 99-609/225, 4-90/283, 99-602/258, 99-600/492, 99-598/130, 99-217/277, 99-576/421, 4-61/269, 4-66/145, and 4-67/40; or a PG1-related biallelic marker selected from the group consisting of 99-622, 4-77, 4-71, 4-73, 99-598, 99-576, and 4-66.

5           22.     A method of obtaining an allele of the PG1 gene which is associated with a detectable phenotype comprising:

obtaining a nucleic acid sample from an individual expressing said detectable phenotype;

contacting said nucleic acid sample with an agent capable of specifically detecting a nucleic acid encoding the PG1 protein; and

10           isolating said nucleic acid encoding the PG1 protein.

23.     A method of obtaining an allele of the PG1 gene which is associated with a detectable phenotype comprising:

obtaining a nucleic acid sample from an individual expressing said detectable phenotype;

15           contacting said nucleic acid sample with an agent capable of specifically detecting a sequence within the 8p23 region of the human genome;

identifying a nucleic acid encoding the PG1 protein in said nucleic acid sample; and

isolating said nucleic acid encoding the PG1 protein.

20           24.     A method of categorizing the risk of prostate cancer in an individual comprising the step of assaying a sample taken from the individual to determine whether the individual carries an allelic variant of PG1 associated with an increased risk of prostate cancer.

25           25.     The method of Claim 24 wherein said sample is a nucleic acid sample.

26.     The method of Claim 24 wherein said sample is a protein sample.

27.     The method of Claim 26, further comprising determining whether the PG1 protein in said sample binds an antibody that binds specifically to a PG1 isoform associated with prostate  
30 cancer.

28.     A method of genotyping comprising determining the identity of a nucleotide at a PG1-related biallelic marker in a biological sample.

29. A method of estimating the frequency of an allele in a population comprising determining the proportional representation of a nucleotide at a PG1-related biallelic marker in a pooled biological sample derived from said population.

5 30. A method of detecting an association between a genotype and a phenotype, comprising the steps of:

- a) genotyping at least one PG1-related biallelic marker in a trait positive population;
- b) genotyping said PG1-related biallelic marker in a control population; and
- c) determining whether a statistically significant association exists between said genotype and

10 said phenotype.

31. A method of estimating the frequency of a haplotype for a set of biallelic markers in a population, comprising:

- a) genotyping at least one PG1-related biallelic marker;
- 15 b) genotyping a second biallelic marker by determining the identity of the nucleotides at said second biallelic marker for both copies of said second biallelic marker present in the genome of each individual in said population; and
- c) applying an haplotype determination method to the identities of the nucleotides determined in steps a) and b) to obtain an estimate of said frequency.

20 32. A method of detecting an association between a haplotype and a phenotype, comprising the steps of:

- a) estimating the frequency of at least one haplotype in a trait positive population according to the method of claim 31;
- 25 b) estimating the frequency of said haplotype in a control population according to the method of claim 31; and
- c) determining whether a statistically significant association exists between said haplotype and said phenotype.

30 33. A method according to claim 31, wherein said PG1-related biallelic marker and said second biallelic marker are 4-77/151 and 4-66/145,

34. A method according to claim 32, wherein said haplotype exhibits a p-value of  $< 1 \times 10^{-3}$  in an association with a trait positive population with cancer, or prostate cancer.

35. A method according to any one of claims 29 to 31, wherein said PG1-related biallelic is a PG1-related biallelic markers positioned in SEQ ID NO: 179; a PG1-related biallelic marker selected from the group consisting of 99-1485/251, 99-622/95, 99-619/141, 4-76/222, 4-77/151, 4-71/233, 4-72/127, 4-73/134, 99-610/250, 99-609/225, 4-90/283, 99-602/258, 99-600/492, 99-598/130, 5 99-217/277, 99-576/421, 4-61/269, 4-66/145, and 4-67/40; or a PG1-related biallelic marker selected from the group consisting of 99-622, 4-77, 4-71, 4-73, 99-598, 99-576, and 4-66.

36. A method according to either one of claims 30 and 32, wherein said control population is a trait negative population or a random population.  
10

37. A method according to any one of claims 22, 23, 30, and 32, wherein said phenotype is a disease, cancer or prostate cancer; a response to an anti-cancer agent or an anti-prostate cancer agent; or a side effect to an anti-cancer or anti-prostate cancer agent.

15 38. A polynucleotide for use in a hybridization assay for determining the identity of the nucleotide at an PG1-related biallelic marker; for use in a sequencing assay for determining the identity of the nucleotide at an PG1-related biallelic marker; for use in an allele-specific amplification assay for determining the identity of the nucleotide at an PG1-related biallelic marker; or for use in amplifying a segment of nucleotides comprising an PG1-related biallelic marker.

20

39. The polynucleotide according to claim 38, wherein said PG1-related biallelic is a PG1-related biallelic markers positioned in SEQ ID NO: 179; a PG1-related biallelic marker selected from the group consisting of 99-1485/251, 99-622/95, 99-619/141, 4-76/222, 4-77/151, 4-71/233, 4-72/127, 4-73/134, 99-610/250, 99-609/225, 4-90/283, 99-602/258, 99-600/492, 99-598/130, 99-217/277, 99-576/421, 4-61/269, 4-66/145, and 4-67/40; or a PG1-related biallelic marker selected from the group consisting of 99-622, 4-77, 4-71, 4-73, 99-598, 99-576, and 4-66.  
25




Figure 1

Figure 2

# ASSOCIATION STUDIES (FIRST SCREENING)

Population	AFFECTED	NON AFFECTED
Sample size	CASES = 112	CONTROLS=76
Population	35 sporadic cases + 77 familial cases	> 65 years PSA<4
Characteristics		





Figure 3

# ASSOCIATION STUDIES (ZOOM)

characteristics of populations	AFFECTED CASES (185)	UNAFFECTED CONTROLS (104)
	47 sporadic cases + 138 familial cases	> 65 years PSA<4







5/24

Figure 5

## HAPLOTYPE SIMULATIONS (100 ITERATIONS)

markers	4-14	4-77	99-217	4-67	99-213	99-221	haplotype frequencies		relativ risk	pvalue
							cases	controls		
haplotype	C	G	T	T	G	A	0,117	0,013	10,06	9,00E-07





Figure 6A

BAC	Marker	SEQ ID N°	SEQ ID N° (mut)	PU sequence	SEQ ID N°	RP sequence	SEQ ID N°	Polymorphism position*	Base	Microseq. oligos positions*
228	99-123	21	30	AAAGCAGGACTAGAAGG	39	TATTCAGAAAGGAGTGGG	48	24	C/T	25-47 (complementary)
189	4-26	22	31	TACAGCCCTGTAAAGAC	40	TGAGGACTGCTAGGAAAG	49	24	A/G	25-47 (complementary)
228/189	4-14	23	32	TCTAACCTCTCATCCAAC	41	GACTGTATCTCTTTGATGCAC	50	24	C/T	25-47 (complementary)
189/463	4-77	24	33	TGTTGATTTACAGCGGC	42	GGAAAGGTACTCATTCTAG	51	24	G/C	25-47 (complementary)
463	99-217	25	34	GGTGGAAATTTACTATATG	43	GTTTATTTTGTGTGAGCTTTG	52	24	C/T	25-47 (complementary)
189/463	4-67	26	35	AAGTCCACTTCTCAAGC	44	TGAAAGAGTTTATCTCTGG	53	24	C/T	25-47 (complementary)
463	99-213	27	36	ATACTGCGACGCGTGTGCTTC	45	TTATTGCCCCACATGCTTGAG	54	24	C/T	25-47 (complementary)
463	99-221	28	37	CCCTTTTCTTCACTGTTTC	46	TCATTCTGCTGCTAGGTC	55	24	A/C	25-47 (complementary)
725	99-135	29	38	TGGAAGTTGTTATTGCCCC	47	AAACACCTCCCATTTGTGC	56	24	A/G	25-47 (complementary)

Figure 6B

BAC	Marker	SEQ ID N°	SEQ ID N° (mut)	PU sequence	SEQ ID N°	RP sequence	SEQ ID N°	Polymorphism position*	Base	Microseq. oligos positions*
189/463	99-1482	57	60	ATCAAATCAGTGAAGTCTGAG	63	ACAAATCTATATAAGGCTGG	66	24	A/C	25-47 (complementary)
463	4-73	58	61	ATCGCTGGAACATTTCTGG	64	CTCTTGTTAAACAGCAGTG	67	24	G/C	25-47 (complementary)
463	4-65	59	62	GATTTAAGCTACGCTATTAG	65	TGGCTCTGCATTTCTTCC	68	24	C/T	25-47 (complementary)

\*, positions are given relative to the sequence of the corresponding marker (i.e. SEQ ID N° 21-38 and 57-62)

7/24

**Figure 7**

<b>EXON Phase</b>	<b>START</b>	<b>END</b>	<b>5' SPsite</b>	<b>PHASE</b>	<b>3' SPsite</b>
Ex1 +0	2001	2216			GTGAGC
Ex2 +1 GTTTGTA	18196	18265	TAG	+0	
Ex3 +0	23717	23832	CAG	+2	GTAAGT
Ex4 +0 GTAAGA	25571	25660	CAG	+0	
Ex5 GTAAGT +2	34669	34759	CAG	+0	
Ex6 +1	40688	40846	TAG	+1	GTAAGT
Ex7 +2	48070	48193	TAG	+2	GTGAGT
Ex8	50182	54523	TAG	+1	
ATG codon	2031	2033			
STOP codon	50405	50407			
POLY Ad site	54445	54450			

8/24

# PG1 genome structure and cDNAs





Figure 8A

9/24

# Prostate Gene 1

Figure 8B



10/24

		box1	box2	box3
PG1	Hs	NHQ 81-83	FPEGTR 160-165	LDAIYDVTV 211-219
AF003136 (Genbank)	Ce	NHQ 630-632	FPEGTR 712-717	LDAIYDVTV 762-770
Z72511 (Genbank)	Ce	48 NHR 50	FPEGTD 129-134	VEYIYDITI 204-212
P38226 (Swissprot)	Sc	111 NHQ 113	FPEGTN 223-228	IESLYDITI 271-279
P33333 (Swissprot)	Sc	81 NHQ 83	FPEGTR 154-159	-
Z49770 (Genbank)	Sc	116 NHQ 118	FPEGTN 215-220	LDAIYDVTI 265-273
P26647 (Swissprot)	Ec	72 NHQ 74	FPEGTR 145-150	-
Z49860 (Genbank)	Bn	-	FVEGTR 90-95	VPAIYDMTV 138-146
U89336 (Genbank)	Hs	95 NHQ 97	FPEGTR 168-173	-
U56417 (Genbank)	Hs	103 NHQ 105	FPEGTR 176-181	-
AB005623 (Genbank)	Mm	100 NHQ 102	FPEGTR 173-178	-
Z29518 (Genbank)	Zm	91 NHR 93	FVEGTR 170-175	VPAIYDITV 218-226


Hs = Homo sapiens, Ce = Caenorabibtis elegans, Ec = Escherichia coli; Sc = Saccharomyces cerevisiae, Bn = Brassica napus, Zm = Zea maize, Mm = Mus musculus

- = pattern absent from protein sequence

Note : Functional acyl glycerol transferases all contain boxes 1 and 2 and not box 3.  
Proteins most related to PG1 contain the 3 boxes with a high degree of conservation.

**Figure 9**

Figure 10



12/24






Figure 11



Figure 12






PNT2 PG1 mut229

Figure 13

Figure 14

## Alternative splicing



**Figure 15**

16/24

### Combinations of exons of PG1 gene discovered by PCR with primers specific for exon borders


Primer	Clones	prostate	PNT1A	PNT1B	PNT2	LNCaPFCG	LNCaPJMB	CoHPV	Du145	PC3	ECF5	ECF6	ECF7	ECF8	ECF9	ECF10	ECF11	ECF12	ECF13	ECF14	ECF15	ECF16	ECF17	ECF18	ECF19	ECF20	ECF21	ECF22	ECF23	ECF24
PGlexon13	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
PGlexon14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
PGlexon15	+	-	-	-	-	-	-	NT	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
PGlexon16	-	+	-	+	+	-	-	NT	-	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
PGlexon17	+	+	+	-	+	+	+	NT	+	+	-	-	-	-	-	-	+	+	+	+	+	+	+	-	-	-	+	-	-	-
PGlexon18	+	+	+	+	+	-	-	NT	+	+	-	-	-	-	-	-	-	-	-	+	+	+	+	-	-	-	+	+	-	+
PGlexon24	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
PGlexon25	+	+	-	+	+	-	-	NT	+	-	-	-	-	-	-	+	-	-	-	-	+	-	-	-	-	-	-	-	-	-
PGlexon26	-	-	-	-	-	-	-	NT	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
PGlexon27	-	-	-	+	+	-	-	NT	+	-	-	-	-	-	-	-	-	-	+	-	+	-	-	-	-	-	-	-	-	+
PGlexon28	-	-	+	-	-	-	-	NT	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
PGlexon35	-	+	+	+	+	+	+	NT	+	+	-	-	-	-	-	-	+	+	-	+	+	+	-	-	-	+	+	+	+	-
PGlexon36	-	-	-	-	-	-	-	NT	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
PGlexon37	-	-	-	-	-	-	-	NT	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
PGlexon38	-	-	-	-	-	-	-	NT	-	+	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-	-	-	-	-
PGlexon46	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
PGlexon47	+	+	+	+	+	-	-	NT	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
PGlexon48	-	-	-	-	+	-	-	NT	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
PGlexon57	+	+	+	+	+	-	-	NT	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
PGlexon58	-	-	-	-	-	+	+	NT	-	-	-	-	-	-	-	+	+	-	-	-	-	-	-	-	-	-	-	+	-	+
PGlexon68	-	+	+	-	+	+	+	NT	+	+	-	-	-	-	-	+	+	-	-	+	+	-	+	-	-	-	-	+	-	+
PGlexon11b	+	+	+	+	+	+	+	NT	+	+	-	+	-	-	+	+	+	+	+	-	-	-	+	-	-	-	+	+	+	-
PGlexon1b2	+	+	+	+	+	+	+	NT	+	+	-	-	-	-	+	+	+	+	+	+	-	-	+	-	-	-	+	+	+	-

**[+]** alternative splicing form with combination of exons 13478 instead of 1345678

# ASSOCIATION STUDIES HAPLOTYPE FREQUENCY ANALYSIS (chrom 8p23)

Figure 16


POPULATIONS	AFFECTED N= [275;491]	UNAFFFECTED N=[130;313]
	ALL CASES	CONTROLS
characteristics of populations	[275,491] all cases	<28 unaffected ( 65 years or older) & <289 controls (65 years or older with PSA<4)



# ASSOCIATION STUDIES PG1 (8p23)

POPULATIONS		AFFECTED N= [275;491]		UNAFFECTED N=[130;313]	
ALL CASES		SPORADIC CASES		FAMILIAL CASES	
characteristics of populations		<=491 all cases		<=294 sporadic cases	
		<=197 familial cases		<=289 controls (65 years or older)	
		& <=289 controls (65 years or older with PSA<4)			


Figure 17



# ASSOCIATION STUDIES PG1 (8p23)

Figure 18A

POPULATIONS		AFFECTED N= [275;491]		UNAFFECTED N=[130;313]	
ALL CASES		SPORADIC CASES		FAMILIAL CASES	
characteristics of populations		<=294 sporadic cases		<=197 familial cases	
		<=289 controls (65 years or older with PSA<4)		<=28 unafflicted (65 years or older)	



**Figure 18B**

name of markers	PG1	Polym.	Freq(cas)	Freq(controls)	abs diff % ( (freq(cases) - freq(controls) )	Odd Ratio	Pvalue	Freq(randoms)	Attributable Risk
99-1485/251	prom	G/T	0,32	0,24	7,4	1,44	2,53E-02	0,29	17,58
99-622/95	in1	G/T	0,52	0,42	10,1	1,51	9,64E-03	ND \$	ND
99-619/141	in1	C/T	0,28	0,22	5,8	1,37	6,93E-02	ND	ND
4-76/222	in1	G/A	0,43	0,38	5	1,23	1,57E-01	0,41	13,15
4-77/151	in1	G/C	0,34	0,26	7,4	1,43	1,35E-02	0,31	18,16
4-71/233	in1	A/G	0,34	0,26	8,3	1,49	1,43E-02	0,28	18,64
4-72/127	in1	A/G	0,36	0,30	5,7	1,29	9,43E-02	0,31	13,25
4-73/134	in1	G/C	0,52	0,42	9,7	1,48	7,29E-03	0,52	26,76
99-610/250	in1	G/A	0,43	0,37	6,2	1,30	8,33E-02	ND	ND
99-609/225	in1	A/T	0,37	0,30	7	1,36	4,83E-02	ND	ND
4-90/283	in2	A/C	0,29	0,25	4,4	1,25	1,68E-01	0,28	9,32
99-602/258	in2	A/G	0,33	0,25	7,4	1,44	2,69E-02	ND	ND
99-600/492	in3	T/A	0,34	0,34	0,3	1,01	7,52E-01	ND	ND
99-598/130	in4	G/A	0,35	0,25	9,2	1,55	7,29E-03	ND	ND
99-217/277	in4	T/C	0,31	0,28	3,8	1,20	1,07E-01	0,28	8,46
99-576/421	in6	G/C	0,27	0,17	9,2	1,72	3,18E-03	0,24	18,40
4-61/269	3'UTR	G/A	0,01	0,00	0,3	1,76	0,527 \$	ND	ND
4-66/145	3'UTR	C/T	0,25	0,19	6,2	1,43	4,68E-03	0,24	13,16
4-67/40	3'UTR	T/C	0,25	0,20	4,9	1,33	2,39E-02	0,24	10,97

\$ Test Fisher - \$ ND: Not done - \* disease associated allele / not associated allele





22/24

Figure 19-B

# **HAPLOTYPE FREQUENCY ANALYSIS** **PG1 (8p23)**

markers of haplotype Max	4-14/240	99-217/277 ln4	4-66/145 3'UTR	99-221/377
	C	T	C	A
distance between mks	<100kb>		<17kb>	
			<43kb>	

PG1	sample sizes	haplotype frequencies		odd ratio	chl-S	P value
		cases	controls			
cases vs controls	cases vs controls					
cases (<=65 years) vs controls	445 vs 294	0,095	0,032	3,18	21,59	3,40E-06 *****
cases (>65 years) vs controls	171 vs 294	0,105	0,032	3,56	20,91	4,60E-06 *****
	271 vs 294	0,079	0,032	2,60	12,13	4,80E-04 ****
sporadic cases vs controls	266 vs 294	0,096	0,032	3,23	19,73	8,60E-06 *****
sporadic cases (<=65 years) vs controls	85 vs 294	0,095	0,032	3,20	12,04	5,00E-04 ****
sporadic cases (>65 years) vs controls	178 vs 294	0,085	0,032	2,82	12,75	3,50E-04 ****
informative sporadic cases vs controls	67 vs 294	0,062	0,032	2,00	2,70	9,40E-02 **
familial cases vs controls	179 vs 294	0,098	0,032	3,32	18,33	1,80E-05 *****
familial cases (<=65 years) vs controls	86 vs 294	0,112	0,032	3,83	17,98	2,20E-05 *****
familial cases (>65 years) vs controls	93 vs 294	0,075	0,032	2,48	6,59	1,00E-02 **
familial cases (>=3 caP) vs controls	79 vs 294	0,123	0,032	4,26	21,33	3,70E-06 *****

Figure 20

## HAPLOTYPE FREQUENCY ANALYSIS (PG1)

Markers in PG1									
size (cases vs controls)									
allelic frequency % (cases / controls)									
allelic frequency % (randoms)									
diff freq. all. % (cases-controls)									
pvalue (cases vs controls)									
Odd Ratio									
Attributable Risk %									
Hardy Weinberg Disequilibrium	cases								
	controls								
haplotype 1	2 MKS	339 vs 167							
haplotype 2	3 MKS	330 vs 122							
haplotype 3	4 MKS	312 vs 122							
haplotype 4	5 MKS	311 vs 121							
haplotype 5	6 MKS	309 vs 121							
haplotype 6	7 MKS	290 vs 99							
			G						
			C						
			A						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						
			G						


24/24

Figure 21

Comparison of Pvalue between nb of mks for haplotype  
(19 mks of PG1)

GENE	# of markers	# of 2 mks combinations	# of 3 mks combinations	# of 4 mks combinations
PG1	19	171	969	3876

CUMULATIVE FREQUENCY HISTOGRAM OF Chi-square statistics (Chi-S) for PG1



<110> Cohen, Daniel  
Blumenfeld , Marta  
Chumakov, Ilya  
Bougueleret, Lydie  
<120> Prostate cancer gene  
<130> PG1  
<150> 08/996,806  
<151> 1997-12-22  
<150> 60/099,658  
<151> 1998/09/09  
<160> 578  
<170> Patent.pm  
<210> 1  
<211> 56516  
<212> DNA  
<213> Homo sapiens  
<220>  
<221> promoter  
<222> 1629..1870  
<223> identification method Proscan  
<221> misc\_feature  
<222> 1998..2000  
<223> potential ATG  
<221> exon  
<222> 2001..2216  
<223> exon1  
<221> misc\_feature  
<222> 2031..2033  
<223> ATG  
<221> misc\_feature  
<222> 11694..14332  
<223> Tyr Phos  
<221> primer\_bind  
<222> 11930..11947  
<223> upstream amplification primer 4-77 SEQ ID42  
<221> allele  
<222> 12057..12103  
<223> polymorphic fragment 4-77 SEQ ID24  
<221> primer\_bind  
<222> 12339..12358  
<223> downstream amplification primer 4-77 SEQ ID51, complement  
<221> primer\_bind  
<222> 13547..13564  
<223> upstream amplification primer 4-73 SEQ ID64  
<221> allele  
<222> 13657..13703  
<223> polymorphic fragment 4-73 SEQ ID58  
<221> primer\_bind  
<222> 13962..13981  
<223> downstream amplification primer 4-73 SEQ ID67, complement  
<221> exon  
<222> 18196..18265  
<223> exon 2  
<221> exon  
<222> 23717..23832  
<223> exon 3  
<221> exon  
<222> 25571..25660  
<223> exon 4  
<221> primer\_bind  
<222> 34216..34234

<223> upstream amplification primer 99-217 SEQ ID43  
 <221> allele  
 <222> 34469..34515  
 <223> polymorphic fragment 99-217 SEQ ID25  
 <221> primer\_bind  
 <222> 34625..34645  
 <223> downstream amplification primer 99-217 SEQ ID52, complement  
 <221> exon  
 <222> 34669..34759  
 <223> exon 5  
 <221> exon  
 <222> 40688..40846  
 <223> exon 6  
 <221> exon  
 <222> 48070..48193  
 <223> exon 7  
 <221> exon  
 <222> 50182..54523  
 <223> exon 8  
 <221> primer\_bind  
 <222> 51149..51168  
 <223> upstream amplification primer 4-65 SEQ ID65  
 <221> allele  
 <222> 51448..51494  
 <223> polymorphic fragment 4-65 SEQ ID59  
 <221> primer\_bind  
 <222> 51482..51499  
 <223> downstream amplification primer 4-65 SEQ ID68, complement  
 <221> primer\_bind  
 <222> 51596..51613  
 <223> upstream amplification primer 4-67 SEQ ID44  
 <221> allele  
 <222> 51612..51658  
 <223> polymorphic fragment 4-67 SEQ ID26  
 <221> primer\_bind  
 <222> 51996..52015  
 <223> downstream amplification primer 4-67 SEQ ID53, complement  
 <221> polyA\_signal  
 <222> 54445..54450  
 <223> AATAAA  
 <400> 1  
 gtggatctgt gactgttcgc aggaagagag gagcgggagc aggacagaca ataactgata 60  
 gtcaggagct gggtttggag ataaagaggg aacaagagaa agttaagtgc tgtgttttca 120  
 tggcaaacat tgcacaaaag ttacaactt cgtgactaac agtaatctgg ggtgattcac 180  
 aacaaattta cacataaaca catatttact gactttatac acagcaatcc taactgaac 240  
 acagaacctg ctttatcttt tcgcacactg ttctagtgtg gagatgtctg gtctcagtta 300  
 aagaaagcat aaggagcatt agttgtgcac actgtccaca cccgtgactt tttccacca 360  
 gtactaaacc tagtgcttct tacagtacag ggcaatgaca gccacagaaa gagagaagct 420  
 ccttttactg tgtaatgctt cctgctggcc ttcaaatact tgttacttga gagatctcca 480  
 ttcacctggc ttgtgtccca aaggtcatca tctaccaatg atgttggtat ttgatgttaa 540  
 tcatgtataa agaaagtagc taccatcctg gccctgatta gaacttccca ctgaaatacc 600  
 gtcctgccta aaggtagcac aggtttccat tatgggtggg gtggggaggg ggcgggaata 660  
 tatatatata tatatatata tatatatatg gtaaaagcatt cggcattctt ttaaagtaca 720  
 actatccttg aaaagggtta catattaaac catttttacc acagccaaag gggaggagaa 780  
 agatccaaaa gtcctgtgga tctgctttaa catcaataaa acagttatcc acccttcgta 840  
 gcttttagtg aagggtacaa aagtatgctt tttatggatt acacatgtgc acgcaactac 900  
 tttaattact acagaaaaaa acgaggctcc ttattaaaaa aaatcagaa acaagtccaa 960  
 cagactctga ggaaatgaag caagagtga ttctgaaaag gtctaataaa cagtatggaa 1020  
 atatccttgt gggattgttc ttcagctatg cataaacatg taattatcat cattactgtg 1080  
 atggggaaaa acacggaccc taattctgaa acaccctggt agcgagagac gggcaggagg 1140  
 ggctgctgcg cactcagagc ggaggctgag gaggcggcgt ccccttgcaa aggactggca 1200

gtgagcagat	gggacactc	gagctgcccc	gcgacctggg	ccgagctgcc	tacaacctgg	1260
gcccaggtgc	ctgcaagaat	tagacctccg	ataacgttaa	caccactttt	ctcactgctc	1320
taattgtgtg	catcccggcg	cccaggggct	tgtgagcagc	aggtgcgcgt	tccaggcagc	1380
tccagcgacc	cttaaacctg	accgcgcgca	cgteccggccc	gagggagcag	aacaagaggc	1440
acccggaccc	tcctccggcc	agcaccacc	ttcaccaggt	tccgtcagtc	gccaccacct	1500
cccttcccgc	gtccgcagcc	ggcccagctg	gggagcatgc	gcagtggccg	gagccgggtt	1560
gcccgcgcca	cagcaggtag	ctgtactgca	actgtcggcc	caaaccaacc	aatcaagaga	1620
cgtgttattg	ccgcccaggt	ggaactatgg	caacggggcg	ccaatcagaa	ggcgcgttgt	1680
tgccgcggag	ccccctgccc	cggcaggggg	atgtggcgat	gggtgagggt	catggggtgt	1740
gagcatccct	gagccatcga	tccgggaggg	ccgcgggttc	ccttgctttg	ccgcggggag	1800
cggcgcacgc	agccccgcac	tcgcctaccc	ggccccgggc	ggcggcgcg	cccatgcggc	1860
tggggggcga	ggctgggagc	gggtggcggg	cgcggcgccc	cgggcccggg	cgggtgattg	1920
ccgcctgctg	gcccgcactg	aggcccggga	ggcggggcgg	gagcgcaggc	ggagctcgct	1980
gcccgcgagc	tgagaagatg	ctgctgtccc	tggtgctcca	cacgtactcc	atgcgctacc	2040
tgctgcccag	cgtegtgctc	ctgggcacgg	cgcccaccta	cgtgttggcc	tggggggtct	2100
ggcggtctgt	ctcgccttc	ctgcctcgcc	gcttctacca	agcgtggac	gaccggctgt	2160
actgcgtcta	ccagagcatg	gtgctcttct	tcttcgagaa	ttacaccggg	gtccagggtga	2220
gcccctctcc	gctcccggtt	ctcggcgctc	acccgagctc	ccgggggcgc	ggacctctcc	2280
gctccccac	agctggcgag	ggtcacccgg	ccggcccggc	ggaccagca	cggagagcac	2340
gtgcgcctc	ccgccttcc	tctccgcatg	cttctgccc	ttctgcccag	atcgctctct	2400
aggaagctgt	ggctgcgtcg	tccgtaggct	acgagtggga	cccgcgccc	ctttcccgcg	2460
ccctgcctg	ggctgatgc	tgcttagcaa	agtgggtgca	gatgcacgtt	ttaaataata	2520
gggcacgcgt	ttagcagttt	ctggcctttg	gtccaaagag	gtggtcatgt	tggaacagat	2580
cggagacgtc	tacactccga	agtgcgcttt	tacagtgacc	tcttgaacaa	gaagtacaat	2640
tcggtcttgt	gttctttccc	ctggacaagt	gaaagctggg	cgaagaaatg	aatacathtt	2700
ttaaccgtag	aagcctaact	agatacaatt	cttgccaact	ttaactgggc	ttgaatgtgt	2760
gggtgatctg	ttgtctgatt	actttctttc	tggtactgtt	tctctgtaga	gattggattc	2820
gtagattaaa	cttgagaaac	aaaccataaa	agtgggaagg	cctctttaac	agtaggtatt	2880
tgaagtgtta	taaaaaaaaa	aaaggtgaat	ttttctttta	tttctcagtt	tgaaagaaca	2940
gctttattct	tggttattcc	taatgtccac	ctagtcctct	tttacttttc	ttggtagggg	3000
tagggtagga	tggggaaatg	ggacggatc	attttgtctt	tttaactttt	tttttttcca	3060
cctacagcag	ctgtttttac	cctgtggtca	gtcaggtact	atatttagtt	tgcagttgca	3120
ctgctgatcg	acccttgatg	gccccagttg	gaagttgttt	ggggggaagg	aactaggaga	3180
ggccagggcc	tccatttaaa	ccagtgtctg	taagtgtctc	cttggaagga	aaaaaagata	3240
ctgttccagg	tcatggtttc	ctggtagttg	acgttttaaa	tgggcctcat	ttaaaaattt	3300
caataattca	ggctaatttt	ttccctttat	atggtaactc	caccaagttt	gtctaaatgt	3360
atgattttta	tcatgattaa	gtttttactt	ccacatcatg	tgacaactgg	cctgggtagg	3420
gatataagct	cagaacacaa	agtcattcac	ctgttataaa	aataattcta	tctgtggcgg	3480
gttatgttat	ttttgttcaa	agaggacaca	atatgatgca	gaatacacca	ttgaaggatt	3540
ttttggtttg	gcaagttctt	atttttttta	atggctgtaa	aacctagcag	tgtttctgaa	3600
attgcatacc	ttacctgatg	ttcagagatc	cgatttactt	cttgatttcc	cagcaagtga	3660
ttttgaaaac	atttaactca	atcattcccc	ccaccgtctg	ttcaaatcaa	aggaagtggc	3720
atccagcact	aattttcatg	catttatgaa	aggatgcctg	aggaccctta	agtataattc	3780
aaaattttgt	ttaatgtgtg	ttccttgatg	aagttcttta	ggagtcgtag	aacgaactga	3840
ttgcccactg	atcatcaaat	gcaagttatg	aacattttaat	aaaaatttaa	aaccaagagt	3900
ttcttgttcc	tgcattttta	tttttattgt	atggagggga	caaataatta	ttttctgttt	3960
agtaacagag	cagggtattt	tgaatttatt	agggctcttt	tctgcagtct	gggtttcctg	4020
tgtaacacaa	gctacctttc	aatatttttt	attgtttctg	ttaagattaa	atcaatagag	4080
gaataaatag	ctatcttcaa	acataagacc	caaaggaaaa	agatttatag	tgatgttctg	4140
tcaccttatt	ttttacctgt	gactttgtac	cattaacttt	gtcactgaga	tgttttgatt	4200
aaaattttta	gcttgctttt	cttgttttgt	taggacactc	tttttttctt	gaattgtttt	4260
tatcagcttt	cgtttgcaag	gctagtgatg	attctcttgt	tctgtataaa	gtattgttga	4320
ctcatttctg	aaggaggttt	tagtaattta	agaggttata	agttttttaa	taaaagggtt	4380
attaatttat	aatatttaaa	gaggcatttt	aaaaataaat	ttttttttaa	tgacattttt	4440
acacctttca	actctaggtt	taaaaaataa	gtggttcaca	gtagttcttg	cagaagaata	4500
ttttctttta	catagaattt	ttaagctgaa	gagaagtagt	agtaggtcca	tgagattttat	4560
gatctgtgct	tggcaggtaa	acctgcttcc	aacaaattta	gttggttttt	tcttggtatt	4620
tggtgaaata	cctttttctt	ccccagtttc	actactttat	tttcatatgt	atctctgaga	4680
tagagaaata	tttcagtcag	tgctgctaaa	attgttccct	ataactcggt	tatcctttta	4740
ggctcttcca	gaatctctca	ttgggtactga	aactcaaatg	gggtactttct	tcaccattta	4800
tttctttaga	ataagtaata	agaattttat	aagctttttt	atatttcacg	taatttgaga	4860

ctattgaaaa	tccagtttaag	tctctctact	gtgttgagag	gcattgattc	aagtacctgt	4920
gttactttcc	tgtgctgcca	aaacagatca	cctcaaacta	agcggcttaa	aataatagaa	4980
cttaagttct	cgtgattctg	gaggccagca	ctttgaaatc	aagggttagg	ctcaatttta	5040
ctccctctgg	aggccctagg	gggaatctgt	tcttgtgggt	ttcaacttct	ggtgactggg	5100
ggcattcctt	ggcttggggc	cccatcactt	caacctctgc	cttacagtcc	ttgctgccac	5160
ctcttctgtc	tcacatctca	ctctcccttt	ctcttagaag	gatgcttgtc	attgggttta	5220
gagcccacct	ggatattccg	ggatgatctc	ttcatctcaa	gacccctaat	tataactgca	5280
aagagccttt	ttocaaataa	gaaaacattc	acaggttcca	gggcttagga	tgtggacaca	5340
ttttttgagg	ggctgccctt	cattccccca	caacaatgaa	ctccatagtt	ctgcctattc	5400
agtattttgt	agttatttcg	tagtttaact	tgccctattt	cttttaggtat	ttacgtatta	5460
aagcattttg	gtctctgctt	tctttaacag	agaacctggg	tttctgtaat	aagtttactt	5520
actttcccat	aatcttttag	tttcttattt	acagattttac	cttcacatat	cccttaagta	5580
gaacattttg	ttactgtttt	tatttttcgga	acaaatctgc	attctgtata	ataaccaact	5640
tattcatatt	tcgggtattct	tttaattctt	atctgattct	gaaattacca	tcttgtgatt	5700
atatatatat	atatatggaa	ataactgaaa	tcttgataaa	ttaaaggtga	tataacttct	5760
aagacaattt	attatgtatg	atgtgggtgaa	tatactgggt	tttgggttgt	ttgccactta	5820
aaagccctat	ctataggata	ggaagtaact	tgaatgtgga	atgcttagag	actcagagta	5880
agaggccgta	tatatatcct	tgagctggag	tttaaggaaa	acttatggga	aattaaaagg	5940
aaagtgtggg	tactgacaga	ggattgcgta	ggactcatga	aaaaggaatg	aagttacctt	6000
aaattctatc	atcgtgagtt	aacgtgaaac	tagattttatg	ttagttttata	gcctagaatt	6060
ctatccctagg	aatctagata	tatcctaact	gttgagatag	ctgcataaac	aataactgta	6120
atcgttatga	taaataatga	caaatctttt	tagcatgttt	tgtgaagctg	ataaatgtta	6180
ataggatgtc	ttcaaatgtc	agaattcttt	tttctttgct	tcttttttaa	aaaatttctt	6240
ttcccccat	cctatgcaat	acactgaaaa	ctgatcattg	aaattttag	gccaaaaaat	6300
taatcaacac	gtaatagatt	gggggtttggg	tttttttgag	tcagggtctt	cttctgtcac	6360
ccaggctctg	gtgcggtggc	accatcatgg	ctcattgcag	ccttgaatgc	ctgggttcaa	6420
gtgatctctc	ggagtagctg	cctgtgccatt	attctagct	aatttttaaa	agtttttgt	6480
gaaatggggg	ctttctgtgt	tgcccaggct	ggctttgaat	tcctggcctc	aggatgacct	6540
tctgccttgg	cctcccaaag	tgctgggatt	acaggtgtga	gccaccatgc	ctagccccta	6600
ataaatattc	taattaccga	tttatcttgc	ttaaatcagt	tggttaacact	tggaatttac	6660
ttcagaatat	attttacatt	agtggctctg	actgctaatt	cccccttctc	caaagtctaa	6720
tgtaatataa	caataaaaatg	cacagttcct	aagtttatat	aaaataaaca	ggttttcagt	6780
tgacctgctt	taagtgtaaa	atagtgtgaa	aaacacaaga	aagaagataa	agaatttaag	6840
attttgacat	ttctctaata	tgcccttaac	ttctccaagg	attcatactt	ttttttgtaa	6900
gacagaatct	cacactgttg	cccaaaccag	aggtgcagtg	gtgcagcttc	cactcactgc	6960
aacctctgcc	ccggggtcca	agcggctctc	ccacctcagc	ctcctgagta	gctgggacta	7020
caggtaacaca	gcacctgcc	cagctaattt	ttttttttgg	tatttttttag	tgggggtaga	7080
gacgagattt	tgccatattg	cccagctctg	ttttgagctc	ctgggctcaa	gtgatccgtc	7140
cttgatccac	catgcttagc	tgattcatac	tcttaactga	aacattgttc	caagtttctc	7200
agaaacagtc	aaggcttttt	atctagagaa	catttataac	tggatctttc	tttgtgtagc	7260
actgattcat	caaactaatc	ctaaactcct	aatgagttaa	atttatatte	tgaatcttgc	7320
tgtaaaagca	gcpattcatt	agaatgaaac	atgtttactt	agaattggag	aagggtgctt	7380
ataagtcata	tagtctactc	ccttttatga	cacttctaca	ttctttctgc	acttctgcc	7440
aaatgttgcc	cagcgtcgtc	tctgatacct	atagtcctaa	caagaatatg	aatcatacct	7500
tgtatcctta	attttactct	tctctgctta	tttgccattc	atgtgaagac	cttaaataga	7560
tcttaaatgg	cttccttcac	tttagctgag	agtgacagga	ctgtgtaggt	gtgggtgtgt	7620
ttctgcattt	gcttatttta	gcaggataat	aaaaactttt	actataggaa	atataacatt	7680
tcccaatcaa	atacaattcc	agtctaacac	aattaaatc	tggttaggga	actgcttaac	7740
ttactagact	tataggaaaa	tactaaaaaa	atgttaactag	aactctattt	ttacacttta	7800
taaatataaa	cctctgtgaa	caaaccagtt	atttcaggtt	gcatttgtgt	atagtttttt	7860
aatgcctgat	ttttctattt	taaaatcaca	gatgcaatta	tacattcaaa	cactgccaca	7920
atactttgag	aaagttaaag	tttcccttac	tcctacactg	cgtacacctt	tcctaggtac	7980
atcccagttt	ggtgtgtaac	tttagatttc	ttccaagagc	ttttgagtaa	gtgtttgaa	8040
tgtgggaagg	ttcttttagt	aaatgaactt	cttacagatc	agtttttttag	tacagttagca	8100
cgaaatatac	ctgcatacct	atgggggatac	ctctgtgcca	ttacgatgga	aggcacggga	8160
aaacagcact	ccgtatatac	ctagtttact	ttccctcttt	tgtatatattg	tctgattttg	8220
tggagctgat	gcttctcaag	tggaaatcaga	agtttaacttt	tcctttacta	ttttctcatt	8280
ttattatggg	ttcttaacta	gaggttgatg	ttagtgggtg	gaccattcaa	tagtaagttaa	8340
tgactttttc	ttaaggggac	tctagaacct	agatccctta	attcctgcaa	tattcccgtg	8400
tgtacattgt	tccaggtgct	gtcctgggta	ccaaggggata	caatgtttga	tagacaatgt	8460
acctgccatt	atggagggtca	cattctagtg	tgggaagaca	aacaataaca	agaaaatgaa	8520



aatttactgt	gccatgccag	gttggttagc	ctgggtgggtg	agaggtaggg	gtttggaaaa	8580
tcttactgag	caagtgcacat	ttgtgtggag	ctctgtaaaa	gggccagctt	ggaaggtaat	8640
gtagtcatcc	aggtgagaaa	tgatggtagc	gggagtggaa	agagtggatg	ttaagattga	8700
aaagaattcc	aatcttattt	tagtggtagc	tgatagggct	ttgtgattga	atgtggagga	8760
aaaagaagag	ggfgggttag	taacacactc	agtcgcagtt	agtgagtgtc	gctgtgtgca	8820
agtattgttc	tattatgtaa	ataattccat	ctttacaaag	taggcaccat	tcttcctctt	8880
ttacagacaa	ggaaaaggga	acacccatgg	ttcacatctg	tagtagccta	gccaggagtt	8940
tcaggcactt	attttctgaa	gatgctctgc	ctggcaatgt	ggttatattg	gttgaaatga	9000
gacccccctac	tttcaaggta	ttcatctagg	aaagacatga	actgccatt	acaatatagg	9060
ataacactga	aaftagagac	gtgtttatta	actttgccat	acagaggtaa	agtaactctt	9120
taaagtaact	ctttgtctgg	gttagtgagg	aaggctataa	aaattacttg	gagtttttac	9180
tttgaacatg	cgtaattaac	atggaatgtt	tagggaaaag	aggttttcaa	ttgataacat	9240
aataaacatg	aggagtgttg	agcatggcat	tcaaggtttt	ctaaattctg	ccccggttaa	9300
cttttccatt	cgttgggttc	attctagtct	agcttttcc	tctggggccgc	ccctccccac	9360
attagaccgc	tcctctctgg	aattccaact	caagcccttg	cttttctcca	tctgtcatga	9420
tgttacccca	tctcatgtgc	agggtaactt	ttatgtaata	ttaacatata	taatactgat	9480
ataacattag	catattttaa	tgtatggatc	atctcctctg	caacattgta	acctcttgga	9540
gatggcaata	atgggaagaa	tgacttgatt	ttactttttc	ttttaacaaa	aatggtggag	9600
tagtctgggc	acggtgtggc	tcatgcctgt	aatcccagca	ttttgggagg	ccaaggaggg	9660
tggatcactt	gaggtcaggc	attcgagacc	agtcgggcca	acattgtgaa	accccatctc	9720
taccaaaaaa	ataccaaacac	ttactgggca	tggtgtgtgtg	tgcctgtagt	cctagctact	9780
caggaggctg	aggtgggaga	atcacttgaa	catgggaggt	agaggctcca	gcttgggcga	9840
cagagtgaga	ccctgtctca	aaagaaaaaa	aaggtaaaag	ggccagggtc	ggaggctcac	9900
gctggtaatc	caagcacttt	gggaggctga	ggcaatggat	cacctgaggt	cgggagttcg	9960
agatcagcct	gaccaacatg	gagaaaacccc	ttctctacta	aaaatacaaa	attagccggg	10020
cgtgggtggtg	cgctcctgta	atctaagcta	catgggaggc	tgaggcagga	gaatcacttg	10080
aaccagggag	acagaggttg	tggtgagcca	agatggcacc	attgcactcc	cgactgggca	10140
acaagagcga	aattccgtct	caaaacaaac	aaacaaacaa	aacaaaacag	agagaaaagg	10200
cagagtactc	tagggaattc	tagtctgtgt	ttctgtggaa	atgtatatga	atctcacttt	10260
taagggatgg	agatttttga	atggcataac	tagttgataa	gttttgctct	aacagggtac	10320
ccaagtctag	tgagtcgat	tcattctttc	cttaaataga	tgaaggagga	agaaacatga	10380
ctccaccctc	aagagttaagg	cagaatgagc	aaagtcagag	aagttaaaaa	agaattctca	10440
cgcagccagc	agtgacagaga	aaccttggtt	tagttgtgaa	tcaaaaccag	tactttttgt	10500
aatttttgag	cctatgcaat	tctccaaggt	tttatgttgt	ttcttctgtt	tctctgtagg	10560
caccagaaat	caaaacccca	aataagaaag	tgttacttga	agattttaga	gtacttattt	10620
gtgtataagt	gtaagtgata	tttggaagac	gactttactg	cgctcctcca	gcttggcatg	10680
agaattccag	gggpcggaag	aaaggagggt	gatgttacct	ggaaaggaga	gtcatgttaa	10740
gtcccagcca	catattaagt	gctaaccacc	tactgttaaa	aggtgtaatg	ttctagactg	10800
acaaaataca	tagtctctac	cgtaaagtaa	cacataattt	agcagtgcag	aaagatgtca	10860
cttaaaagaa	aacttgaata	tatgctgaga	tagttcacaa	attaaagaaa	tgaacaaaga	10920
actgagggaa	taaaggagga	atacaactgt	gtccaaatga	atacttaact	gggtgggagc	10980
tgttgcatat	gtaagcaggt	ggttcaccta	aaagtggat	gtaacgtagt	taacggcagc	11040
tcttgggtgca	cttacatatt	gcattgcttc	cgggcttaat	ttgtgttcat	ataggaataa	11100
attttttgtt	ggtttttaat	tttactcctt	gtaattccgt	ggttgatatt	caaagtgaag	11160
aaaattacat	aagcttctaa	tatatgagaa	gtcttctcac	ttgacatttt	ttatttggaa	11220
tttttgcaga	gagtagtttt	gtcacagtca	aaagattttg	ggatcttgca	gtgagaaacc	11280
taggtgtaat	tcctattttc	ctgccattcc	gtatgtcatc	tggattaagt	gtcaacttct	11340
cagtctcaag	attctcgtcc	ttaaatggaa	tactttttgt	catgctattt	tgaagacaaa	11400
atgagataat	acgtgaaact	gcctagctca	gtgaatggta	catcatagat	actcagaaaa	11460
aacacaccct	ctaaaataag	aacagtacca	aaagacagga	tgtaaaaata	gggcagtagc	11520
aaaagacaca	tgcatgctga	gtgtatgaga	aagaactttg	tggccttctt	gggtggcaca	11580
ggccatggca	gttccacagc	atgacgtggg	tgctgtgggt	ggtagagcag	acatgccgct	11640
cccgttact	gcctggcttt	gatgcttgc	ttcttcagct	gagaggacgc	agctgtgata	11700
tgaaggtctt	gtgtgtacag	tcgtgacctc	acatttccaa	tttctgtctg	gcagaaccca	11760
cagtctacaa	cgtacgagca	ccagagttag	cgtgagacag	acagcataca	gaggcttgta	11820
acatccttct	ggpaaacact	gtgtaagctt	tcagtgcgaa	taaacaatgat	cagtggcaag	11880
ttctgttaga	tgtagtctgc	aagcatcctg	attttactgg	gcaagactat	gttgatttac	11940
aggcggctga	tgtatccatg	gatagccac	tactagtatt	ttcacaatt	tcacaagaca	12000
ttcttactgg	aagattgccc	gtttcttatg	atactgctgc	ccttttagct	tcatttgctg	12060
ttcagactaa	actgggagac	tacagtgcgt	cagagaactt	gctaggccac	ctctcagggt	12120
attctttcat	tcttgatcat	cctcaaaatt	ttgaaaaaga	aattgtaaaa	attacatcag	12180

caacatatag	gcctatgtcc	ttgagaagca	gcagttaatt	acctaaacac	agcaagtacc	12240
ttagaactct	gtggagttga	attgcactat	gcaagggatc	aagtaacaat	aaaattatga	12300
ttggaatgat	gtcaagagga	attctgattt	ataacaggct	atgaatgagt	acctttccat	12360
ggtcgaagat	tgtaaaaatt	tgttttaagt	gcaaacagtt	ttttattcag	ctttgaaaat	12420
gacttgcata	aatctggaga	aagattatca	ggatttaata	tggtgaatta	tatggcatgt	12480
aaacatttgt	ggaaagcaag	tttagaacat	cacatattct	tctgtttgga	cagaccactt	12540
ccaactagaa	agaatTTTTT	tgcacattat	tttacattag	gttcaaaatt	cctaattgat	12600
ggtgggagaa	ctgaagttca	gttagttcag	tatggcaaag	aaaaggcaaa	taaagacaga	12660
ctacttgcag	gacctctcaag	taagccattg	acgtggaaat	taatagtttg	ggaagtagta	12720
ggcaggaatt	caatatctga	tgaaaagatt	agaaacataa	agccttccat	cacaattccc	12780
acccggaaca	ggaattccta	ctcatcaaaa	ttctgcattc	atacaagagg	gaacctgatt	12840
atgaccatct	tctgttggtc	atgttgtaga	ttatgtgggt	cacacttctt	ccaaatattt	12900
gcaaactcaga	catcaccatt	atcagcacia	gctaatagca	tcattctgga	atcatcacta	12960
ttacaggaca	cccctggaga	tgggtagcct	ccagctttac	cacccaaaca	agctaagaaa	13020
aactgttggg	accaaattca	ttattttacat	tttcaacaag	atctggaaga	tcatattaat	13080
gaaacgttga	tgttctatct	tctcttaaaa	aatctgtctc	taatgggtgg	attctacatg	13140
ataatcgtgt	tctaattccga	gtgaacctga	cgaaaattgga	aggtttgagg	tcaatgcaaa	13200
gggggatatg	atcagaagat	gtctgtgatc	gtgtcctgag	aagcaccagg	aacacctttg	13260
acctcagtga	ctctcgattg	aagagaagac	caagttgtat	tgatcagtgg	ttgggacttt	13320
acagaacaca	cccatgattg	ggttgctctg	ctttttaaag	ccaactgtga	gagacattct	13380
ggggaactca	tgcttctagt	tctacctatg	ctgcataatga	tgtagtggaa	gaagtgtctag	13440
aaaatgagac	agacttccag	tacattctgg	agaaagcccc	actagatagt	gtccaccagg	13500
atgaccatgt	gctgtgggag	tcagtgatcc	agctaaccga	gggcttatcg	ctggaacatt	13560
ctggacacaa	tttgatcaac	ttatcaaaaa	aaaacttggg	atgacaattt	ctgggtgccag	13620
attaccttag	aacttttgca	aaaatagata	gagatagttt	tccttatgat	gttacatggc	13680
ttatttttaa	aggtaatgaa	aactacatca	gtgtaattcc	agcatcataa	gtcagaacag	13740
tgcttgtcaa	ggggcggtac	cacacacttg	aacagatttt	tggcagatga	cttggaaca	13800
aggctcctcc	atgtttgtaa	tgttgaccac	acaagttgaa	tgtggcagag	ttaaattgacc	13860
ccaatattgg	ccagaaccca	caggaagttc	atcctatgga	tgctaccaag	ccttctgcc	13920
ctgagaagaa	ggaagcactg	tctttatctt	caggaagatc	acactgctgt	ttaaccaaga	13980
gaaaaattag	agpgtcatca	atcacgcaga	tccagtacag	aggggtggcct	gacctgggag	14040
accctgatga	ttcagtgaat	ttctggattt	tgtttttcat	atgcaaaata	agagggtctag	14100
caaggaaaaa	ccccttgttg	tttcttgtag	tgtcttgagt	ggaagaacca	gcgttcttaa	14160
tactatggaa	acagccatgt	gtctcattga	tctcattgaa	tgtagtcagc	cagtttattc	14220
actagacatg	gtaagaacaa	tgagagagca	gtgagccgtg	atgggtccaa	cacctagtca	14280
ttacagtttt	gcgtgtgaa	tactattttg	aaagcttatg	aagaaggctt	tgctgaagaa	14340
agcaaaaagg	aaaaaagaac	tttgtcatct	gttaggttcc	atattattgca	tgataattgt	14400
gtttgtattg	attattgggc	aagtagctgt	ttgctatttt	gatcttattt	cagaagggca	14460
taataatttt	actattcaat	gaaacgtttt	aaacggggtg	gaaaaagact	agtttttgta	14520
tgctttacag	cagaaatctt	ataatgatta	actggtaata	tatttcgttg	gcataaaaaat	14580
acatttaaaa	gttcaagtaa	ttataaacat	tgtaaatgt	atatgtaatc	atattgaaat	14640
tgaaattctt	tatagctgta	cttctgtgta	atcaaagact	ggggagagat	agactagcta	14700
gctctttctc	ttatccatta	atcacttaac	agagttttga	ataaaaagtt	ccatttcatg	14760
ggataagaat	aatgacaggt	taacctattt	tagttgggtt	ctatgttcta	ggtgttgat	14820
gaagttagtt	acatagtttc	actgatttca	ctacaatccc	aggaggagta	gttactatta	14880
ttacactcat	tttacaggca	aagaaatagg	tttgaggggg	ttgggtgttt	tgcccaagtt	14940
ctcatcgtaa	aatgacagat	gaggattcaa	attcaagtct	taattgaaat	ccattacttt	15000
agaacctacc	tcttagtggc	tcttatgtta	cagtataagg	gagagcagac	tgttccttta	15060
cccttgtagg	gtagctaggg	cttggtgaatt	aagagactga	ttaacaggag	aagaggcata	15120
cacattttat	tgacgttagt	atttttacat	gcacagggaa	ggagggtttt	atttttattt	15180
ttatttttat	ctttatttta	aagagacagg	ggtcttgctg	tggtgccagg	gctggactca	15240
aactcctgaa	gcgaagcgat	tcttctgctt	gagattcctg	agtagcaggg	actataggtg	15300
tgctcctctg	tgcttggtga	aagaagggtg	ttgtatgtga	tttttaacaa	aggctgataa	15360
attgtgaaga	agtgaactag	caaaggagaa	gaggatttca	gctcccaggg	gtggtaaat	15420
gtgggaagat	gactaggaaa	tgtatagtaa	taaggtttgc	tatgcaggtt	tattttgcc	15480
gtttctggtc	tcctaataag	ggacagggaa	acacctttac	agatggaaat	tcatatcacc	15540
tttccacagg	gaaatttatg	tcctgcctta	ggcagttagg	ggaagggcag	agaattcttc	15600
ctgtatctgc	tgtgtctcag	gtgccttcag	ctcaaaataa	tccttatgcc	aaagtagcat	15660
atttgggtgt	ggcatattct	ctgatctctt	tcaacagcat	catctatact	taacaacagc	15720
aaaagttttt	tttaaaaaat	catgtttcaa	gatttgcag	tggaagacaa	atggacatga	15780
ttgagataaa	tgaagaatat	atatttttta	acaaagaatg	ctgtatatatt	atgtctctgt	15840

gacatttgtgt	tatggagggt	aagggtgttaa	gcatgtgatt	acttttagatg	ccgtatgact	15900
acctgtttttt	aagattaaaa	agaatcaat	aggcagttta	tatgcatggg	agcaagttaa	15960
aaacaacaca	gatgtgatga	aggcgagggtg	aaactgggtcc	gcatctaatt	caggccttct	16020
cctgaaagcc	agtgtgtgca	agataaataa	gtttgtttga	cgaaagcaga	ataactagtt	16080
tgctcctttgt	gatgaagata	gttattcaga	aatcattttt	attggctacc	tctgaattaa	16140
taaatgaaaa	gagaaatttt	ttttctgtga	ggggatgtct	gatgagttct	taaaaagtg	16200
atgaacctga	aattatcatg	aacaagcaat	tataatgaac	ttaaaattac	ttaaagagtt	16260
atgaaaaaca	aaaagaaaaag	ccgtatgttt	tcttgtgcct	tattttgaag	tgacaaatta	16320
tttgagggt	acattttag	acggaactaa	tgtgatttaa	aaaaatgagta	ctagattttac	16380
agaatgatgc	ctttaaaaag	tcactgggtgc	actttaatta	ttttatttat	gtttattctg	16440
aaactacctt	tattttgaaa	atgagggtata	gctttgccta	ctggtgacaa	aagtgtaaat	16500
aattcagtaa	acatctgtta	aaaaccagct	tgggtgctagg	ctcttggggt	agaaaactga	16560
tcaggccatt	gaggagctca	tagtccctaa	ggggctgggg	acttgtcatt	aggtgtgcag	16620
tgtgttctgg	atgctcctga	aggagtgtgg	gcaggtgcgc	accaccatgc	ctggctaata	16680
tttttataat	tatgtagaga	cagggtctgg	ctgtgctgcc	catgctgggt	ttgaacttct	16740
gggcttaaga	gatcttcctt	ccctgccct	accgaccctg	cccgcctact	ccacctcagc	16800
ctcccaaaag	cactgggatt	gcaggcatgg	gccactatgc	ctgggctgtg	caaaaactttt	16860
aaatcagtg	atctcaatg	gtcttgatgc	aattctggct	tgttggttaag	agaatgggga	16920
tttactcaca	agccacgatg	tcacttttaa	ctctgaacag	atcaagctat	tggtattact	16980
catttatgtc	atcgataaac	tttatgaata	aaaactcatt	gtgcaaatat	ttaaacatac	17040
tacatacata	gcactgtgca	gtttctaagg	aaagtaatgg	aaacctttgt	cacatccctg	17100
gcttccagaa	ctttatgtta	tctaagtgc	ttgtctgca	aagttgttgg	gttaattgct	17160
cctttctttt	ttctcttttt	aagatattaa	taaatagtgt	catgaccaa	agataatcct	17220
tatggacaag	atagatctaa	aaagccttag	ctaatttata	atcttgcata	atccatgatg	17280
acaagatgca	gaaacaaaaa	tgcccagaat	aaaaacttag	caccattagc	agccatttcc	17340
ttttaagtct	ttacaagtat	actcccagtt	tcttgaaaaa	tttattctaa	aatatgtaag	17400
acacacaaaa	cagcagaagg	actaatacag	gtacatcgaa	cacctgtgtg	cctaccgccc	17460
agtttaaaaa	taaactggaa	tgatgtttct	ctcactacta	cagaataaag	ttttaatctt	17520
tagcatggaa	ttcaaaagac	ttctgccatt	ccagttcaga	gccacccttc	tggctctcct	17580
gctcctcagc	cgcgacactg	cccatgtacc	caacaggcct	ccagggttac	tgcttccatt	17640
cgttcttatt	ctcatgaaca	ttttccttca	tctcatctgc	cagaatccta	cctaataata	17700
ctcctgtctt	gcagtttaca	gttctttaaa	attaaaaaag	gttgtgtacc	ctttagtgtc	17760
ctgaaaaaag	aaaaaacaaa	tttaaaacct	taaaaaggta	ccataatttc	atagtatttg	17820
cgttatgtct	cattacagtt	cctgtggaca	tgtctgtctc	ttttactaga	ttgattgtgg	17880
gctctttgaa	ggaagatata	tcttatgaac	agtgttttat	atattgttag	caatcaatga	17940
atgcttgcta	tatttttctc	atgaggatat	tgattattct	attttaattt	attaccnnnn	18000
nnntgtacta	tacataactg	ctttctgtac	ctgagctatt	tatgatctct	gaggctcctg	18060
tgagaaatct	aatttttgtt	aatcatggat	ggaatatattc	acaacatcat	tcgtcagttt	18120
cttcacattg	tcttcccttg	tattattacag	atgtttttaa	atatcaaagt	aatgtttttt	18180
tgttttatct	tttagatatt	gctatatgga	gatttgccaa	aaaaataaga	aaatataata	18240
tatttagcaa	atcatcaaag	cacagggtttg	tatttcattt	gcatgaaacc	taggtttttc	18300
tacagatggc	acatgggcat	tcaaaatacc	gttcttatat	ttaaatgaag	tgggtttttt	18360
aaaacagcaa	ttttctgtgc	agatattaca	cctgttcttg	tatttttgtg	attttacttt	18420
ttggaaagtc	agaaacttga	aagctatgaa	ttttcctaaa	cttaccttct	ccctctgttg	18480
gatgtaagta	agctatcttc	ttacttgctt	gctttgtttt	tcctttgtgt	agctctttaa	18540
agagtgtatt	cattcttttt	gtaagtgtatg	tttctagaag	tagcattggg	gggtcgaagt	18600
gtgtatacat	tttacatttt	tgatttgctaa	gctgcagaaa	agctgtattg	gtatgtaagt	18660
actcgtttcc	ttactatgct	cgctatttct	agtgtctgct	cttcctttcc	ttcttcaaat	18720
gggtttgggt	taattctagt	tgctactggt	ccatcagagg	aattgcagag	aactgggtctt	18780
caaaacagtg	cagtatatatac	tttaggtgaa	gatacttcta	aaaacctttg	tatttttgagg	18840
taattctaga	gtcccaagaa	tttgcaaaaa	gagtacattg	tcagcaatat	ttttcccaat	18900
ggtgacatct	taatataact	gtagcacagt	agcagaatca	ggaaattgtc	attgggtaag	18960
gtacttttta	attctccaaa	taattcagcc	ctccaaaaaa	atcccaactc	ttatgttttc	19020
aaacctgtag	ctacttttga	tgcgtaacttc	ctaaattgca	tttttattac	tttaaaaaat	19080
ataataccta	gaagctcaaa	gctggaaaca	gcctgatcaa	tatagtactc	ttaaagctaaa	19140
aacaacctga	tcaatatagt	actcttaggg	aaatcactta	tgctgtggc	tttttttaaa	19200
ttttcttcct	gtcagctgtc	tcttcatgat	tttgtgggtt	ttattactgc	ttataccata	19260
gatgagggtat	agaaagtaaa	agaagttaaa	atgcattttt	ctcaatttag	tgaattaatg	19320
attacattca	gattttatagg	acaagggttg	aagctancaa	gggttgata	ggaatcttga	19380
tgtatctgag	tattttcccc	aactttatta	catgactggg	tcagactatt	ttatctaat	19440
acatttctact	cttggcagaa	atagcaaaac	agtcaaccaa	tgggtcaatgc	tgctgagaa	19500

tctggcctgt	gcagacatat	tggctgtttt	acttctaata	ccattctgct	tttcctgtcc	19560
tgctgctgat	ggatgtttct	tccagggttt	aaatatcaaa	caaaagggat	ctgtgggccc	19620
agtacaggga	atggctcttg	atagatttga	tttctctgca	tttcctttat	tttgatccag	19680
tgtaatttc	atgtagagtt	gtctgtttaa	caggattctc	ttaaaattcc	ttcttcagtt	19740
tacctgccag	cttttctttg	tccagggttt	agtatgaact	ccactcgatt	aatagagctc	19800
tctagttagt	acttgtggag	tgggttctct	gaacatttct	ggaagtgttg	ctgatagtga	19860
taatattgat	cactagtact	gttaatttgt	gtgcttacta	catgttggct	tttatatgta	19920
ttccttcaga	ttaaggactt	ctagaaaaca	tccatgaaaa	aacagattaa	aaaaaacaat	19980
tctgcatgta	tttgggacta	gaaggctacta	tgggaaggat	aatcttcata	ctcagaccat	20040
actgacctga	atttcattta	tcagttttaga	gaaccacttc	cccttccctt	caccctacct	20100
ccgagtgcct	gtgactttgt	atcacccgtc	tggcaccaca	tcctcatccc	agcaggattt	20160
gggaaggctg	ctttttgaaa	gcctttttaa	attctgtaag	ttgagaaaat	actaggggaa	20220
tgattttaaa	tttctttaga	attacaggct	ttagtcagta	tatgacagag	ccttttcccta	20280
gaaaaatgtg	catataaaaa	tttgcattgta	gttttagggg	ttcagagacc	cctaaagcct	20340
atccatagac	gtgggttcatt	gtctgattgt	gttttaggtac	ccttctaaaa	cccttttgag	20400
atggttaggaa	tcacaacaga	gtatctctga	aaatgtaatt	agcggaaaga	acatttcaaa	20460
gactgttgtt	ctgttagac	tttctagttt	gtcttctgcc	aggcttgccg	gaataaatga	20520
gtttcctggc	ctgatactca	aaagaattga	catttaaatt	agtctctctc	ttcccttggt	20580
ttcgcttgac	acatccttgt	ctctacattc	tgtctctgtc	tctgttagct	tatttctctc	20640
tcgagtcagc	aggatatagt	ggctgttatt	tcttccctct	atccttcaac	gatctacttt	20700
tgacaacact	ttgccttttt	ttttttgaga	tggagtttca	ctcttggtgc	ccaggctggg	20760
tgtaatgggtg	caatctcagc	tcactgcaac	ctttgcctcc	cgggttcaag	ccattttcct	20820
gcctcagcct	cccgagtagc	tgggattaca	gacatgcacc	accacgcctg	gctaattttg	20880
tattttcagt	agagatgggg	tttcaccatg	ttggtcaggc	tggctctgaa	ctcctgacct	20940
cagggtgatct	gcctgcctcg	gcctcccaaa	gtgcagggat	tacaggcgtg	agccactgtg	21000
ccctgcctgc	tatttgcctt	tttaattctca	tgaatgttc	tcttttcttg	gctgaagtgt	21060
cacttttctt	gttgaacagc	atgctgtgtg	agtagaatgt	tataaaaagg	gatggacttt	21120
ggagttagag	agaccagggt	tcctgttcgg	cattgcagaa	atgctgttct	gcaataggct	21180
gtgtgtcagt	gggcaaatta	cttatctctc	agagccttat	tggtaagggtg	tgagtgatag	21240
ctcctttcag	gcacettaca	gaggctgtct	cctaactctg	gtagcgtacc	tggctcatag	21300
atggcattta	aaagtgttg	tgatgacagt	catagctcac	cattagcata	gcgctggatc	21360
catggcaggg	aagcgtgca	catgcagtat	ctcttggtact	acacaggggc	ctcatgaatt	21420
aggaactgct	gtttcatgag	gatagggatg	aggaaattag	acttgctgcc	cctcactgcc	21480
ttccactcct	ctctccaag	ttaattggga	ctatgactct	gctttggctt	gattgccatg	21540
gaagattctc	acacagccaa	atttattgct	atcttagtta	aattatgcca	gaacacaaaa	21600
tatgaagtta	ttgtcaaagt	aatataatct	cagctgtaac	tgagatagtc	agaaactgtc	21660
tgtaatctga	tgtcctatct	gaaaggtagc	tgagaataaa	caagaaataa	agagaattca	21720
gtagcaaata	ttggtgacac	aaagctttta	tattttgact	agttaaagcta	gttcttaaat	21780
gtttccacta	aatatttcaa	gttttaagggc	atagcccagg	gcagcttatt	atgaacatga	21840
tgtatttttg	aatctttaca	ctttctctta	aaagttcttg	ggaggggcat	gtgaggccat	21900
aatataacca	taaaaccatt	tgttttaaaa	taaaaccctt	ttttaaaatt	cttccaaaata	21960
aaaaaattat	tgcaggaaaa	aatgctaaac	ctggttttta	actttgtacg	ccaactatat	22020
ttccaagatg	tgctgtagcc	tggtaaccat	acagaacctt	acagaattag	ttctcagaat	22080
ttattgtctg	cttacttttg	catttggtac	aggataaaca	gggtcgatta	tatggtttct	22140
aagacatgac	tagaaagaaa	tatgtttatc	agttattatt	tcttccatct	aaattagaag	22200
gggctaggga	gagggcttca	acaggaattt	atatacttta	gagaaaagtg	atcattgata	22260
gccaatagat	atagatatct	caaccaata	acacagggtg	tgtctgtctc	tgggatcata	22320
cactgtaggg	gagaatcttt	gcaagcaaca	ttctacttat	agggagccat	aacaaaagtt	22380
tcatatgtat	aataattata	agtcttaagt	catcaagaaa	aagttaactt	gtgaatgata	22440
atccctgatt	aaaaagagag	atgtataata	atggataaga	gatttttctt	ggttaatttt	22500
tagtattaaa	atggctaaat	cttctttggg	atattctgac	tagtatgggtg	cattgtctaa	22560
tagatttccc	atagctgaga	gctaactcat	ttgtaatctg	tggaaaactg	tcctcttttg	22620
ctaaaacttt	attgtaattc	ctctaaatcc	tcagctttta	ttttctacag	actttttttt	22680
ttttttaaca	tttcttctct	ctgactcact	ccttttggtc	tcatttttcat	ggcctgagaa	22740
catgggtgat	gatagaatta	ttcttttcac	agattaacag	ttttcttttc	gagtatcggt	22800
gagctcatgt	gtgtattaac	tagagaagtc	tcctttacat	ttcattttta	tgttttcttt	22860
ctcatcagga	gatagtttgt	agccatttac	tttcaaatcc	aagtttctgc	ggttcttaag	22920
acctgtatca	tttgtctcct	gaatttcact	tcatttctct	tttaaaccat	gtcctctggt	22980
tcccatcttc	tgcacccact	ttgccacttc	ctgtttgttt	aattggcaag	ggccactctc	23040
tgtgttggaa	atttttcttt	tttgaagct	caactaaca	cttctaggaa	gttttttatt	23100
gctactgtta	tcaattcata	ccatcttacc	cttgtttttg	caaccctttg	ttaataacat	23160

atttatttaa	ctatagttat	tagcagttctg	agatcatttt	acttggttac	ataaggagca	23220
catatatcta	cccagcatca	ttgtaaggca	tgtgagacct	ttgtttgatt	gctgtcctaa	23280
cctagtaccg	agtcctaaaa	actcatttagt	agaagatgaa	gtgtccttgc	cttttgctga	23340
acatatatat	acacactgaa	tatttagtgg	caattcatag	ttgcatttgg	ccattttttg	23400
tttataattt	cccttttctc	attaaaaaaa	ctttgttttc	tagacttttag	gatttagaga	23460
agctcatttt	gttccataca	catgctgctg	ttggattatt	taggtatttt	gtgactgtat	23520
tttatctttg	aaataaaaaa	cctttcaaga	aatgcaaaaa	aaaaaagctc	aaaaaacaga	23580
aaatgtatat	tttttaataa	tctcagatag	atttaaagaa	attttaaaca	tcctaatacat	23640
agtacttttg	aagcccattc	atagtacaac	ctgtgaagag	cctcatgtac	gcgctaactg	23700
ggtoctgtct	ctgcagttga	ctggattggt	gctgacatct	tggccatcag	gcagaatgcg	23760
ctaggacatg	tgggctacgt	gctgaaagaa	gggttaaaat	ggctgccatt	gtatgggtgt	23820
tactttgctc	aggtaacttg	tttccatgct	tttctctcta	tatatgtagt	ttataaattt	23880
tttttttttt	ttttggagac	agtctcactt	tattgctcag	gctgagtga	gtgggtgtgaa	23940
cacagctcac	tgcagccttg	acctctgggg	ctcaagtga	cctcctgcct	ctgcctccca	24000
agtagttggg	acogtagtgc	ccaccatcat	gccccgctaa	attttctatt	ttttgtagag	24060
atgggggtct	cgtgtgtgtg	cccaggctgg	tcttggactc	aagcaatctg	cctgtctcag	24120
cctacccaaa	tgctggatta	taggtgtgaa	ctgccatacc	caaccctata	aaaatgttat	24180
attttaaaaa	ttaacaatat	acttcatgtg	aatgtatggt	ttttaaaatg	ggtttaatat	24240
tttattctca	gttgaagtaa	ttttgttttg	catttttagt	gggtgtgtatt	tatatacgtc	24300
tgattatcca	tatgcggttt	tccttcagca	tctgtgggga	ttgggttttag	aaccaccaca	24360
gataccaaaa	tctaaggtgt	tcaagaccct	catatagaat	gggatagtat	ttgcatataa	24420
cctgtgcact	actttaaatc	atctctagat	tactataat	atctaataca	ttataaatgc	24480
catgtaaatg	gttggttatac	tttatttttt	atttgtatta	ttttaattgt	tataattatt	24540
ttaattttta	tttggttcaca	tattttttgat	ctgtgatttg	ttgaatctgc	agatgtggaa	24600
ctcatggatg	tgaagggcca	gctgcagtaa	aatgaaagag	caaaaatgca	aatgtacaaa	24660
gttcaaacaa	ataggaaatt	taaaggcata	gaatttgata	ggcaattaca	ttaaactgtt	24720
gataacagta	attagtgtac	tgtatgatat	taaaaaaaa	aagcaaactg	tatatataaa	24780
acttactttc	tcagtttctg	gaggctagac	atccaagatc	aagggtgtga	cagggttagt	24840
ttctcccaag	gcctctctcc	caggcttgca	gacagcatcc	ttcttctgtg	gtcctcaggt	24900
ggtttttttc	ccgtgtccca	agcaccctcg	gcactgcttc	ctcttcttag	aaggactagt	24960
tacactggat	gaftaatcct	tctacagaga	ctgtcaaggt	cccactctga	ggcccttttt	25020
taaccttaat	tapcacctct	aagtcctctc	ctctgaatac	agtcacagtg	ggaactatta	25080
gggcttttagt	agactgattt	gggggaacac	acttctgtcc	gtaacagtgc	cacataaata	25140
tcttttagcag	gattgatttt	ttaaaatccc	taaagatcgt	gagttattgac	atgttaagga	25200
cgcttttttag	tgactctgta	ataagtgggt	ggaagaattg	ggagttaaat	ccatctgatg	25260
gatcagggtt	tttattttta	aaaatgtgta	tttaagaaag	aaagcatttt	cattttaact	25320
gccaacaaaa	ctaaacttca	tgtgttttcc	aatacagtg	cacatgcagt	ttttttgaat	25380
tatgttgaga	caaggcaatt	tgttcttttag	aagctaattg	ttgaagatat	25440	
taaataataga	ttaaattctg	aaatgtagtt	ttcattctgt	acttttttga	agagaagttg	25500
ccttttttgat	gactctggcc	aattgttatt	ttaaaagtaa	atgctctttc	tcccgaatttg	25560
attgtggcag	catggaggaa	tctatgtaaa	gcgagtgcc	aaatttaacg	agaaagagat	25620
gcgaacaag	ttgcagagct	acgtggacgc	aggaactcca	gtaagagcct	acccgttttt	25680
attttttctta	ccagctctca	gtttctaaat	ttagaatta	aattaaaatc	taagaattgt	25740
tttgacaatg	tattttccca	tgtgtaatta	ctaattcagg	gttatgctga	ggtaacagaa	25800
accctctatg	tacaggtagg	caggtttttc	agccatcaga	aagattgctg	taaacacta	25860
ggctcctttg	tggtcagtgg	accttaaaaga	ggaataaaaa	gagcatttgg	tgtcgttcag	25920
agtcctataaa	tagaactaac	tgcattttta	cctgacattt	aagctagtgt	acaagctcat	25980
cttacttctt	gtcttcttta	gtatcagatt	tggttttaga	agcagcaact	gttttctgtt	26040
agtgcataat	ttgaatgtct	tacatgtaca	gaaaaaccaa	aaaaggatga	atctctacaa	26100
atgttaaatc	attcagtgtg	aataatattt	tataaaactt	tattccacaa	aagtggggag	26160
agttcaatct	gctttgtata	gaatgctgat	tgtctccaaa	ggcttttccc	ctggttccct	26220
ccggagacaa	agcaccatga	tcaccggggc	gacttgggct	ttctctttca	gtacatgaca	26280
tgtgctcaga	agcttagctc	gtgtgcacag	gctttccctt	tcctttctgg	ctccctccct	26340
ctgtcttccc	tcctctcctc	ttgccctccc	ctcaccagg	gtcctgggca	gcagctggag	26400
ctcatggtga	aggaagaatt	cttcatggtc	agctggcgaa	gtgcctgggt	tgagcattgt	26460
ttattcacat	gcctcttcta	gggtgtttta	cattagaaca	ttgcatctgt	tttggggcatg	26520
tgttgggtga	cagaagcaga	atgggaatgag	atgaacagtg	accctttatc	ctgttatagc	26580
taacccttga	gaaccaagct	tgggtgtcttc	aaaggggtctg	tttagtctga	aacagtgtgg	26640
tgaatttggg	cagaattgtg	gtcattgcat	gtaggtctcc	aaaagacaga	ataagttggg	26700
aatatgggtt	atcgactttt	tacaaaaaaa	atttaaaaat	catgaattta	taccttaaaa	26760
tgtccatccc	acttctctcc	cagctgtcca	gtcaccctag	caatggatga	ctgctgtgga	26820

gttccttctg	tgtcctgctg	tgggcattgt	atatatgaag	caaatgaaga	tagctgcctt	26880
ttgggtgatg	ttggcatcct	atgcacagt	gtcccttgct	tttttgcccc	catgaatata	26940
gctgccagtg	gcgctagggc	tgaaaaaatc	agctctttac	acttgatcatg	tgtcttgttt	27000
atgtggctgc	cttcgtgagt	ttcttcttgt	ttttggtttg	cagcagttta	agtatcatat	27060
atctgagtg	caattaaaaa	tttttacctg	gattggctct	ctgagcttgg	atctatgatt	27120
tgggtgtctg	tatttaatttt	ggaaatttct	ttgctcttat	ttccttaaat	attattccta	27180
ccccagtcct	tccttctccag	ttatgtttgt	gttgggttcat	ttctcgtctg	tcttttagttc	27240
ttagatgcat	tattcgtttt	ttgttggttt	ttttttaaat	tttttttttt	acgccccctc	27300
ccttttttct	ttttgtgtta	catttttgat	aatttctgtt	gacccacctt	tgagttcatg	27360
gattcttctt	ttggctgtgt	tgagctact	ggtagaccag	tttaaggcac	tcttcacctc	27420
tgctactcgc	tgtttcattc	ctcacatttc	cctttgacct	tgtttcatag	tttccatctc	27480
tgtgctagtg	tatctatctg	atcataaagc	ttagtcacgt	ttccagttg	aacctttatc	27540
attttattat	acttgacagt	ctcttaaat	ccctgcttga	taattccaac	atctgggcca	27600
tatctgagtc	tgcaaatttt	gattacttta	tctcttcaga	ttgtgcttta	tcttgctttt	27660
gtcatacttc	ctaagatttt	gcctaacgct	gggccttttt	tgtaagacag	gagaaatgga	27720
ggcaagtgtg	cttgatacct	ggaaatggat	agacttgtct	ttctgcttgg	cttttagtgt	27780
tgaggagtgg	agtcagtcga	ctgaggaggt	gcactgcatt	tgggttttgc	tcatgtgctt	27840
tttctcacag	cttcagggtt	ctgtagaact	cattactttg	ttttaggtt	gggatgtctc	27900
tcccgcctaga	gcttttcttc	agtgtctatt	tcacactcag	cgttttcaca	tagacctttg	27960
gagtggtctt	cttcttttatg	cctttcccca	ctatacttct	tggatacttg	ttactgaact	28020
ctcgctagtt	tgggtggtaga	aggagagggg	agggaggtgt	cttttcattc	ttagggagaa	28080
tctcagggtt	ggagccttct	ctgacccctg	cctgtctctg	gctgtaagtc	tgtgccaggt	28140
atgtattcct	gcctttacta	agagtttttc	ctgttctctc	tcaccagcc	tcacgagta	28200
ttcatccgtg	ccpcatgggt	agcagggttt	tgttgcctct	gttcacaggt	ttcaggctgc	28260
tgttccatag	gaaaggtaga	aagaaggatg	tgggctgggc	cctgagccct	tcccacaggg	28320
ctgcttttcc	ctcccacaag	cctacatcca	gtcttccctg	accgcaggtg	gttttctttt	28380
ttctttgtct	tgtgagtaca	caggaggtct	gtgggtcgag	cctgtgaaat	gtgctgcatt	28440
ctccttgtgt	ctgtagccca	gggttctgct	tgttccactg	gctcatactt	ggctttctgc	28500
aaaattgata	aaatttttag	ctaaattctt	tttactggtg	tctgttacat	tggccccaac	28560
ctaaacaacc	acttgcatct	tgtttctcct	ttgagttttc	catctttcct	tagacttttg	28620
ggtagtttgg	ttgccttgca	accttgagc	tctctgaagg	gtctaagaaa	agtcatgaat	28680
ctacagcttg	tcagtgttgt	tgttgttgta	gggttggcag	tagtatctct	tcagcattct	28740
acatacttaa	tggaaagccg	ctcccatttt	tggtaataaa	atttcaaaac	ttggaacaat	28800
gttagattta	caaaaacgtc	agaaagaaca	gagtgttctt	gtttattctt	tatatagctt	28860
tttttttttt	tttttttttt	gagttggagt	ctcggtctgt	caccaggtct	ggagtgcagt	28920
ggcacgatct	tggctcactg	caacctctgc	ctcacgggtt	caagcaatct	cctgcctcag	28980
cctcctgagt	agctgggatt	acaggcgtgc	accgccatgc	ccggctaatt	tttgtatttt	29040
tagtagagac	agggtttcac	catgttggcc	aggctggctt	cgaactcctg	acctcttgat	29100
ccgcccgcct	cgccccccca	cagtgcctgg	attataggtg	tgagccacca	cgcccagcct	29160
tcttcatcta	gctttaacat	ctaattgttg	catcttacat	aacatggtat	atatttgtca	29220
aaactaagaa	ataaacattg	gtaccacact	attaattgta	ctacagattt	ttattcagac	29280
tttaccaggt	tttccactaa	tgtccttttt	ctgttctaaa	atacaatcca	gaatagatac	29340
aaatccattc	aacttcagtg	ttttaaatta	ttgtttttca	ttatatgaag	tgtgtgtgtg	29400
tttttgtcaa	atctgttatt	ttggttttta	tcttcaagct	tgtcttttgt	tctttaagtg	29460
ataaaggcat	aattttaaag	gtgtgttggg	ttatttcagt	gcctaaagtc	ttgtctgagt	29520
cacttggttt	ctgctgttct	tgtcttatgg	actttctttc	cctgttttgt	ttgttatctt	29580
cctttgctgc	tggctgtgtt	tgggttaagtt	atttgtggaa	atcagttgaa	gcctcaggtg	29640
ggagtgtctt	tctccggaga	acatttctac	ctgttttagc	tgggccccct	aaggctcctc	29700
tagcgtgggc	ccpacccaaa	cgagattctg	agttgaaggt	gaactgagcc	attcaggcag	29760
tgcagccagg	gttgcagatg	cacgtgagac	ctgctcacct	ctcatttact	ttcacctga	29820
gagtagagcc	tttgggtgtt	cgttcacttg	tctgattctc	tcttcacagt	tctattagaa	29880
ggtecatggg	ttttgggttt	tgtgcccttc	atcttatgag	tcttgtaaata	caaagttctg	29940
ttttatgctt	acttctgctt	tactgtgttt	gcttaatttc	agtcttaaca	tcttgccaac	30000
tcttgggtac	ttttaaaata	atgttatatc	cagcttttta	agttgttttc	agtaggaagg	30060
ttgattcaaa	taacctagtc	tggttatggg	ctacgagaat	agcctccctg	ttttttgttg	30120
gcaaaattcc	agccttttat	gttcctagcg	cagtgtggat	aacagactgg	caggttcaag	30180
aggccgtgct	gagcagcttt	cactgtaagg	tcactgtccc	aggtcgggtt	tctaagaate	30240
tggatggttg	tttcatttct	taatatgtac	gccctgtgag	agcggatata	tcttgctcag	30300
gttcttatga	ttcttttgtt	tctgaaggtg	aattaagtaa	gtgacatggt	agaatatggt	30360
aagtcaactt	tcgtgtggct	tactagttct	catgaatcta	ttccatgatt	gtatcagttc	30420
ttattcagta	ttagtatatta	agaaatgcag	aattttgttt	caaaaaatat	atttgtatta	30480

taagttgtga	agaaatacat	ctccataaatt	attgctggga	caatacagta	ttttcttaag	30540
gaacttattg	gttgtggatg	caaatgaagc	atatttgtga	taaaaataac	taatagaagt	30600
cattttgtta	gactatgagc	tagtaaaact	tatggcacia	acatggagac	ttaacacttt	30660
ttcttccagc	tttcaactaa	gttccttttc	agataggagg	cagcctgggtg	gataagagta	30720
ttggttttga	aatttagattc	aggtttaaat	cccagatctt	ctgtttaatc	tttattttat	30780
ttcaggtaga	ttttctggat	aacttgctat	agcttatacg	tcagtacttg	ccacttcaat	30840
tttatgttat	ggagagacgg	cttctttcct	taaacctcac	gaaccaacct	ctgctagctt	30900
ctaagttttt	tcctgccact	tctttacctc	tctcagcctt	cagagaatta	aagggaagta	30960
gggccttgct	ctggattagg	atttgcttta	agggagtggt	gtggctgggt	tgatgtttta	31020
tctagagcac	tcaaactttc	tccatatcag	caataaggct	gttttgcttt	ctaactatc	31080
atgtgttcag	tgaagtagca	cttttaattc	tctttaagaa	cttttcttt	gcatccgcaa	31140
cttggtctgt	tagtggaag	gacctagctt	ttgacctacc	ttggctttca	acataccttc	31200
ctcactaagc	catttctagc	tattgatgta	aagtggagaga	catgcaactc	ttcctttcac	31260
tggaaacgct	agcagccatt	gtaggggtat	taattggcct	aatttcaata	ttgtgtgtc	31320
tcagggaata	gggaaaccca	aggggaggta	gagagaaaga	gagacaggag	aacaggccat	31380
cattggagca	gtcagaacac	acacgacatt	tatcaattaa	atttgtcctc	ttatatgggt	31440
gcaattcatg	gcaccccaa	acaattacaa	tagtaacatc	agagatcaca	gatcacata	31500
acagatataa	taatatgaaa	tattgtgaga	ttaccgaaat	atgacacaga	gacgtgaggt	31560
gagcacatac	tggttgaaaa	atggcaccaa	tagacttgct	cgatgcaggg	ttgtcataaa	31620
ccttcaatgg	gaaaaaaatg	caatttccgt	gaagctcagt	aaagcgaagc	atgataaaat	31680
gagatgagcc	tgctcactct	aagaatgttc	ctgtacaagt	tttttgcctc	tggtacttac	31740
cttttcttat	ttgtgaatag	tatctttttt	tagtaactgt	gtttttttat	ttttatacat	31800
ttatatgtat	cttttgaaga	acatactttt	aagcttaatt	tattgatttt	ttttctctca	31860
taatttccac	tttttgtatc	ctattttaaga	agtccttgcc	aaacttaagg	ttgctaagat	31920
tttctccttt	gttttcttct	ggaaatttta	gagttttgct	ttacatttta	gttctaggat	31980
ttatttataa	ttaatgtttt	catatgggtg	aagatcgaag	ttcatatttt	tttaatatag	32040
gtaaccatca	ctatagaaaa	gattatttcc	cccgaatgtt	tgaataaagt	agactgaata	32100
tagatgggtc	tggtatccct	agatcaatgg	cctgacttgg	ctgttatatt	gatctatata	32160
tatatatcct	tatgccaata	ccatactgtc	ttataaatgc	ttgctttgca	gtaagttttt	32220
aaatagtgtg	gttgtcttct	aaatttgttc	tttcttttca	aagttgtttt	ggctattttt	32280
gggtttttgc	atttctgtgt	gaattataga	attagctcga	caatttctac	ccaaagtttg	32340
tgggcttttc	attttgattg	tattgaagat	atagatgaat	ttgggaagaa	ttgatataac	32400
aggattgaat	ctttggattc	atgaacgtag	cctgcatttg	ttacttagg	tcttctttat	32460
ttatctcagt	gtgttttgta	gtttaatgta	cagatttgca	catcttttgc	cagatatatc	32520
cctaagaatt	tcagtttttg	atactattgt	agatgacatt	taaaaaaatt	tcaagttttt	32580
gtttgttgac	ctaggcatat	atttgacttt	ttaatatact	aaccttgcta	aacttattta	32640
tcattctagta	acttacaaaa	tatatctcct	aggatttcc	acataaacia	tcagtgcatt	32700
gttttagaaa	taacagtttt	actttgtcct	ttttaatctt	gatggctttt	atttcttttt	32760
cttgctaaat	tttctggcta	gacctcctag	tacagccttg	actagaactg	gtgtgaggga	32820
aatcctttcc	atattcctca	tcttttaggga	aaagcactca	ttcttttatc	cattcttttag	32880
ttcctagccc	cattgcccct	cctaaatttt	ttctcatcat	tttcttcat	cacaccttgt	32940
tctttttctt	tgcaatcata	tcatgatatg	taacgacatg	tttttattta	tctgtttaat	33000
gtattttctt	tcptcacttg	tccatgaagg	gaaggaccat	atgtgttggt	atccttttgt	33060
cagttcctgg	aapataataa	gtatataaga	aatagtttct	gaattagctg	tgaatgaatt	33120
catgccttcc	tggtgtctgt	caatgttctt	ttaaattaaa	catctaagac	agcaaataat	33180
accacatgag	ttattaacct	gagaaataat	cgttttat	ataaatgact	gagttgaaag	33240
ctgatagccc	acagtaattg	ctttcatggc	tttgaatata	aaccttactg	ttacaaaaca	33300
cattttcatg	aaaatgaatg	tgtgggtgtt	ggaactagct	ttaatgtttg	tcttctctgt	33360
tttcttctta	gttgctataa	tataataagg	aattttgtat	gtttttccta	attgtaccca	33420
cttttctaca	ttttcttaac	agatctgggtg	aatcttcatt	attaaatata	attatacata	33480
taaattattg	tttaataata	atattaatta	ttaaaaataa	tataaattat	taaatataaa	33540
gatacatata	atattatctg	ttaatttcta	agttaggtgt	gggttctgaa	gactattata	33600
tgaatgaaca	aaaagcttgc	atatttgcgt	ggaagctgaa	agtacgaaat	tttttagatac	33660
cattatacca	gtatctaaag	aaaaaattca	gttaccataa	gggttttaag	taggagctgt	33720
atgatcatag	gtcatccaga	tgaaggaagg	cttctgtacc	agacgtacag	aggtagacag	33780
tggtgtctga	gtactgtctg	agatctggca	agaatgaatc	caataaacgt	agttttctcc	33840
catgagctcc	tgtcttgttt	cctgtattct	gtttgtat	gaaaagattt	gggtgtgcata	33900
acttattttt	gtcttttggc	tgtcaatcaa	agttattagt	gtagtttttg	taactcagtt	33960
ctcaagctag	gagtttttgc	tgataattt	taattgatta	gtttttactt	tcctaagcag	34020
ataagcgtaa	aaacttagac	taattgatta	cttattaaac	gtccagcttg	atattctctt	34080
ttatattatt	ttagtttcag	tttatataac	aaatgaggtt	tcttataaat	aaaaattaaa	34140

atgcactaaa	ggagctgtgt	gaaataggaa	ttctgtgtga	agcttttgaa	tgtgaacatt	34200
tagaacgttt	cacatgggtg	gaatttacta	tatgattttc	atcaaatgag	gtacttttta	34260
gtgttggtac	ttaacgatac	tgattttctaa	aatttgtatt	tctaaaaatg	acgtattaca	34320
ggatctgaaa	gggcaaaaac	tcattgaggg	ttgtatgag	tcagcgtttc	atggcctatt	34380
tttaattagt	gaattattag	catataatta	gaaatgtttt	tagattcttc	atggctgacc	34440
taccaatgaa	tgtagcactg	cattttaaatt	atagttcacg	ttatgttcat	acttaattgt	34500
tgcattttgt	ttgcccctct	tgaacgaag	gtcacatgta	aataaatata	cattttctcc	34560
tactgttagga	aatactctgt	tagcattagt	aggtttagct	tttttaggtt	aacaataaca	34620
aaaacaaagc	tcacacaaaa	taaaccaaat	ttgctctatg	tcccacagat	gtatcttgtg	34680
atttttccag	aaggtacaag	gtataatcca	gagcaaacaa	aagtcctttc	agctagtcag	34740
gcatttgctg	cccaacgtgg	taagtaaaaa	tttgagtgtt	tgaacaaata	attttcaaag	34800
ataataacat	tttttagttt	tcttcctgga	aaagatactt	ttgttttaca	gttgaaggaa	34860
tgaatgtatt	catctcctga	attagtgtac	atattatctc	ttaggaaatg	aagtttcttc	34920
tccttaattc	actttcatgc	tattattaca	tatatctgag	aaattaagtt	gaagtgtctg	34980
ttacgatata	taftcttgtg	ccatggattt	atttaaaatc	tatctaagta	catgattatg	35040
tagatggaag	ctttttctac	agtgtatggg	ttatatgtaa	tggagcttct	gttttgtaag	35100
atgacagacc	taagtgtggg	tccaaactcg	tacttttatt	agctgtatgg	ttgcaacttg	35160
gaagtgtgtg	aatgttgctg	agcttgcttc	ttcatctctt	aaaagaacat	atgccttata	35220
agtagatcta	aatctgtgtg	aggattagat	tagaaaatat	gtcaagtttc	tattggagaa	35280
gttacacaaa	gttgggtccac	agtgttgga	agctgttaat	gtcttcaaca	atggtaatgt	35340
tcttaatatc	catattttag	aaaattgaat	aattgggtaca	ccaataagct	atgcaattta	35400
accaaattgg	gaagtataca	gaaaacagtg	gctatgctat	gttcttagag	gtgtctttga	35460
agcttgactg	tgatttagtg	tgtgatctcc	atatgttgat	agtcactcac	tgagcaataa	35520
ccttggtggg	gacattacag	cagggcctat	gacagtgtcg	tctaattgga	ctttctgcaa	35580
taatggtaaa	gttcttcatc	tgttctgtcc	agtgtgtctg	ctcctaccaa	tgtgggtttt	35640
gagcattcaa	catgtgacta	gtgcatgaaa	ctaattttta	attttattta	attttagttt	35700
aataaaaaat	aagggggagt	ttttacaagg	tgcttacaag	agcagatatg	tcataggtat	35760
atgacatcat	ttgtaacagt	acttttaaaa	aatgccagtt	tgtttttaaa	cacatgtcct	35820
attaagtaag	gagtgtttca	gaataggagg	gttcagttgg	tctcccatc	tgccagctct	35880
cttttgactt	tcattgtctc	ctctgtctaa	tagacatgac	gttctgtcat	ttcagttgct	35940
cttttgcaat	gccattgtct	cttttgccct	tttcacattt	attaacaga	acaaaacaaa	36000
aaccactctc	gaatctgtag	tctacttttg	ttgtaagcac	ttttccagt	actcactctg	36060
ccctcaattt	gttttggtct	gatttgaaat	tctctcccta	gacttctgtg	gggctgttct	36120
ccattatcct	cccaactctc	tggcgattac	ttcctagcct	cctttccagc	ctctttctgc	36180
ttcattttct	cctgtctacat	gtgttatatt	cagtgtcagg	ttttgggtgt	tgattaaatt	36240
cactttttgt	ttctcatggg	ggccttctct	taaatccatg	gcttttagcca	tcgtttcctt	36300
gactgtctgat	gactcgcaaa	agcttctctc	cctccatgtc	tctctgccta	actctggacc	36360
catttgtaca	attgtccatt	agagagcttc	gcttgactgg	cccaaaaagga	tgtctcaaac	36420
tcagcatatt	gaagatagaa	tttatccttc	catgcataca	ctcatatttc	ttgtcttggt	36480
aactccatca	ttcagttttt	ttgcctaagt	tttattcaca	aaaagaacaa	attgatagca	36540
gttgcatacc	tcttatagga	aacttagaca	tggaggaaga	agctgttcag	atggggctct	36600
gcagaagtgc	agggcactgtg	gtaatatatta	aacttttctc	agctgttcga	agggttttgt	36660
tttaactaat	tttcccttaga	cttgttttag	gtatttggct	ttctaattgt	tataagggat	36720
gtggaattaa	atgtatctta	atctgccacc	tggaccattt	aaagtaagcc	cctatgggtg	36780
tttttttttt	taattgccat	ggttaaaacc	atagttgcta	gcgaagggtga	catacttaag	36840
ctttttgaac	tctcttaaaa	gaaaacagaa	atttaatgat	gtgtctataa	tggcaaacca	36900
gatacctaga	atttccatgt	tattcatagg	gtgaataaca	ctggcgattg	tagagatttg	36960
agagttcttt	caaaacagga	gaacaaaggg	aataagctac	aaagcaattt	ttttctttgt	37020
agacttaact	gaataaaaaat	tatttttatg	tctcaaacat	catatgaaca	aatttagttg	37080
gcaaatggca	agctaataat	attttataat	ataggatatt	aataacttta	atattacaaa	37140
agtgtttcat	aattagaaaa	gacataaact	agaaaaatgg	gaaaagggca	tgaataagaa	37200
attcaagaga	tacaaatgac	ccacacactt	gaacaaatgt	ttattctttc	tcataatcaa	37260
agaagttaga	atataatgaa	tactttgaag	ccaacttctg	agaaagcata	gcaaacaaga	37320
aagctagtgc	tcagctttgt	gtggtaacgg	cactctcgct	cttaagaagg	tgtgtttgct	37380
ccctgtggct	gctctcaggc	agggccacaa	acttgggtgg	ttaaaacacc	acagatttct	37440
tctcttacat	ttgagaagtc	tgaatgggt	cttactcagc	tgaatcaag	gtgttggtg	37500
ggctgcagtc	ctttgtggag	gcttgggggg	atcttgttct	cctgtacggg	gtcctgtgct	37560
tggttcgggg	tcctgtgctt	ggctggggat	ccgtgtgctg	gttcgaggtc	ctgtgtggg	37620
tccagtgtct	tgcttttacc	accttgaagt	tcatctggaa	atggcactgg	ctcgccaca	37680
ccatatagct	gactctggtt	ctccctctct	ctcactcgct	ctaaacctgt	gtttttggct	37740
gattttcta	ctctctttcc	ttggcccttc	tgcagcttgc	agggccttct	gcagctcttg	37800



tctgccccag	ccccgggggc	tgcccatccc	agtgtctgggc	tggtctgttc	ctgcccctgcc	37860
tttccctcagc	ccttgggcaac	cctgtttgtt	ttctcccttc	cttagcagtg	gagaacatcg	37920
taagatcaat	gctgactgcc	ttctgcagcc	aagccaggcc	atttcatttc	agccgagcca	37980
agtctgtgtg	gagcagttct	tttatttttc	tccttttgac	tacctcatgg	ttttcacgga	38040
tttttgttct	cttcacattc	aaggattttt	tgctttcaga	aagttatatt	tctctggaaa	38100
gagtgcaccc	aatatccctt	ttgatttcaa	aatcttaatg	tggagtctct	tgacttggat	38160
ttctttggaa	gaactgtctg	aagctgccat	gtctaagaag	aaaacttttg	agaaaaattt	38220
tcttcttaga	catggcaacg	tcaacagttt	ctaagctctt	gattccgtct	accctgtctc	38280
catcgttgcc	tcagtcatct	gccttacttc	ttctgcaggg	tttctcccag	cttgcaaatg	38340
tactccaatt	ctgaaataac	taagtctata	gctgtgcaaa	gagaagtctg	ggccccctgc	38400
tttcttgtgt	ttgactccat	ccactctcca	gaaatgaatc	ccacttctca	cttaaccact	38460
gacctccaaa	gcacgtatc	atltgtgtca	gttgtcatat	ttgttaactt	tcacataact	38520
tttgacatta	ttataacctt	tataaccagg	aaataatttt	aactttattg	tagaaataaa	38580
caatggagta	taatttttct	tggtgaagat	aaatatcacc	tcctcttctt	ttaaaccatct	38640
cttccctttg	tttttgtatt	acattggttt	cccccttttt	tttatttcct	gggttgctgt	38700
attccctgtt	attattttta	cctttttttt	tttaagtggg	atgtttccgg	agtctgtatt	38760
tcttgccctt	tcactctctg	ccctttatta	ttctcagcca	ctgccattac	ttcagttatc	38820
cattcccatg	gtttccacat	gcttagcttc	gggttgattct	tgccatttta	cagaccatat	38880
ttccaactac	ttctagaatg	ttttgttctt	tcagcctcag	tatgcccatt	ttgaactcat	38940
gttctctctc	cccccttctt	cttcccttct	tccttctgct	tctctccctt	ccttcttttc	39000
tttccctccc	tccttctctt	ccttccctca	cttcttctct	cttgcttgct	tgctttctct	39060
cctctctctc	tttcttttct	gcnnnnnnnn	nnnattcttc	tcctcccttc	tcttcttctt	39120
ctccccact	ccccaaactt	caggctaaag	cagtcctcct	gagtagttag	gactacagac	39180
atacacgtgc	caccgcgccc	agctccgtgt	tctctttgtt	tccttgcttc	ctgctcttcc	39240
acttatcttt	gcattggcagg	tggtgtgcag	caggcatgct	ctgcatgtct	tcctcttggt	39300
cattcccttt	ctagttagtg	tgtggcttta	tctacgcgtt	ctggagcaga	agcctagtca	39360
caaagctatt	tttttaaaac	attcatgata	attcatttcc	ttttatgttt	taaaaatact	39420
agctttctgt	ctttatttcc	ttactaactt	acttggatgc	cagtaattag	ttgttttagt	39480
gaacaccaca	gagtgatatt	ttgaaacttt	ggacttcata	aagttggatg	agctccagta	39540
gcaaagaagg	aagtgttaac	tagtttaact	gacaaataaa	tgcttcccag	cttggtgtgc	39600
gattgagatt	ttgttgcaa	gtttgtgaat	caatttaact	gcccctgccc	tggggactaa	39660
agtcagatac	gtgcttggtg	gaatctttgt	ctttcccaca	ccaccctgca	ttttaaaacc	39720
tcttgtgtgg	gagagtccca	ccatgtaata	gctgttcttc	cttactcagc	tactttccct	39780
ccagagaggc	cagtagaaaa	tctagactag	ttttttatag	tctattttca	tgctacttat	39840
tgagagctac	tgttttctgt	taaattgtca	gtaaatattt	taatcaagga	aaagggaggc	39900
aataggaagg	agagaagaac	aaatccttaa	ccctagttagg	aacctaatga	atgggatttg	39960
ttctggataa	ttgcagtagt	ccccagcta	aagaaccttt	taaaaatatg	tcagatatac	40020
ccaagaggat	tgaaatcgta	tgttcataca	aaagcttggt	cacctgcagc	cttcatatgc	40080
aattcctatg	aatgttcata	gcagcattat	tcataatagc	caaagtatgg	atgcaaccca	40140
aatgtccatg	aagcaattaa	taggtaaaca	aaatgtgatc	tgttcacaca	gtggaatact	40200
aactattcag	ccataaaaag	gaatgaagca	ctgagtcctg	cagccacaca	gatgaacctc	40260
agatccatgc	tgagcgaaa	aagccagaaa	caggaggcca	tgtgctgtgt	gactgtatct	40320
ctaggaaatc	ttgagtcacc	atgggcaaga	tgctatcacc	tttgttcagt	ggccagaagc	40380
gagggcacta	atatttacct	ttgccggggt	ctactagatt	gaagcgtttc	cgctaggcca	40440
taaacttoca	acacggtgac	ttgtacatgt	agatatttga	tcaatatata	gcaaatgaat	40500
attgatttaa	acagaaaaag	gcaagtgaga	gtgctttcta	aacttagagc	cctaaatata	40560
tgaggttggt	gaattaatag	attctgttgt	gtgtgtttga	gggaatttaa	aaataattta	40620
gatgttaaac	agtatatgtt	ggaggtgttt	tgttaactaat	taatgacggc	actgaattga	40680
cttctaggcc	ttgcagtatt	aaaacatgtg	ctaaccaccac	gaataaaggc	aactcacgtt	40740
gcttttgatt	gcattgaagaa	ttattttagat	gcaattttatg	atgttacggt	ggtttatgaa	40800
gggaaagacg	atggagggca	gcgaagagag	tcaccgacca	tgacgggtaa	gtgtgttcac	40860
gcacctgaaa	tgccgttaca	cggtatatac	agtgacatg	tttatgtaga	attcagtttt	40920
acaaagtagg	ttaagtgtac	ttttttcctc	cattacattt	acccggtata	tttttcaaga	40980
tgttattaag	attgaacagt	ggagatttca	ttagtctctg	aaagtgtggt	atttcttggt	41040
tgctgtgtga	gtcctgtgga	ctcaccaatt	atcattaatc	cagcctcttt	ctactcaaag	41100
ttcacactta	aaaggaaagc	tctgtaaaag	ggaggaagac	gtgaagaagg	agcacgcctg	41160
gcagtactga	gtgcacgtta	ttagtcagtg	ctgccctttt	gctgtatttt	tcgtaaaata	41220
tttattaaat	ttgggtgtca	ttgtgacaag	aagaaatgca	gttaagtgtg	accttttttt	41280
ttccccaaac	atgttaggtt	ttaagaacct	ttagactatt	gtcagatata	accagaaaaa	41340
aatagaattt	taagttagca	ggataactta	gttaaaactaa	ccaaacatag	tgtagctgtt	41400
tagagaaatg	taaactatgga	aataggcaaa	cagggaagtg	tgtggagttt	ctgtttcctt	41460

ttcaaaatat	ctgttttgagc	tgggggttgag	agagaacact	aggcttcatg	gggtttttttt	41520
gttttttcgtt	ttttgtttttg	agacaagagt	ttcgcctctgt	cgcccaggct	ggagtgcagt	41580
ggcgcaatct	tggctcactg	caacctccgc	ctcccacgtt	cacacgattc	tcctgcctta	41640
gcctcctgag	tagctggaac	tacatgcgtg	tgccaccatg	catgactaat	atttgtattt	41700
ttagtagata	tgggatttca	ccttgttggc	caggctggtc	tcaaactcct	tacctcagg	41760
gatccacgca	ccctcgccctc	ccaaatgagc	tttgtgtttt	tacctcatca	gctgtttggg	41820
gttgagccac	taagtatgtc	agtgtgcttg	tatcagtagg	atctactgag	ggcagatggt	41880
caaaatatga	gcctccagca	cgttttacat	ggaaaccctc	acctgaagca	ttcgtctgaa	41940
gttgatgtgc	cttggaatt	ttatagagta	atatttttaa	ctacaacaaa	acattttataa	42000
aagtagacat	tattaaagca	ttcagaagtg	agcaaggata	gaaattattc	tgcccaacct	42060
tacacgtagg	ccttctagac	gtagtactgt	gcaccgttac	attatctaac	actgtctgtg	42120
tgtcatcttt	ggatgttagg	gattttttcca	aagttcagtg	agattatagt	tgtcaaatga	42180
ttagtctgtt	aaataatgat	aagatgaggg	tcactcaggt	tttaaaagaa	aagctctttg	42240
actgaaagag	agagcagctg	tctactgcag	aaagttaggg	agggaggctg	gaggagtgg	42300
gcccaggggc	tagctagtat	aaaaattggt	tatggctcgaa	ggaaaaaaa	atgtaacata	42360
tttatatctg	aaagatgatt	gttctcataa	ttgtatataa	cacagagtaa	ttgtaaagta	42420
gaaaactaag	gtgttttttca	tttttagatgt	aaatgttttag	aatatgtaat	gcatacagttt	42480
aaaaattaaa	actgtacgaa	atgcacagtg	aaacgtcttc	cttgcttttc	acctgtctac	42540
ctggccttcc	cttctccttc	ctagcgataa	ccagttttct	taatttggtg	tgcgttgat	42600
gtgcaaattt	aagtatatct	tcttattcta	ccatccctcc	cttcttacag	aaaagtggca	42660
tattaatatt	ttctctttt	aaactatcga	aggagtta	tacctatttt	tgcatttcaa	42720
aacagacagt	tcatcaagat	gttcgttggt	ttattaaaca	tagtttaaga	ttaacaagt	42780
gtttataacc	aatgaaaaac	agatagactc	cccataataa	ccttgtttaa	atgctgtctac	42840
ttttatcatg	tcccctcctg	tctaagaacc	ccttggttca	gcagagctca	tgggtaaggc	42900
cagcctctgt	tgcctgccat	cggaggaatg	cgttccagcc	gtgatctctg	ccttgccctc	42960
gcttctctct	gtgctgtgcc	gtgaagcctc	ggcctgtgtg	aagctggctg	actgagtcct	43020
cctgcacccc	atgcataattc	agtagttgaa	gctttgtgt	ggccaatcct	gctttccaca	43080
ggaaaccacc	ctctctttttg	ttgccctcat	ccaaggctac	tgttctccca	gagtgacagg	43140
cggcaccttt	ccpagcatag	cactgtgcct	tctcctgccc	ctgctcttgc	agtactgctg	43200
tggcactgat	ggcgtgtgtt	acagtgtctg	cacttagcac	agggctctgc	ctttctctct	43260
tcccagccgc	atcataagtg	ccttgaggaa	gccaaaacct	tctgtgagtt	gcattgcctg	43320
ggttccaacc	tcccactgcc	ctgcttatcc	tgtgtctacat	gtgagctgac	tgtggctttg	43380
gggtggtcac	tgcctatgtg	tattcattac	aaattgtctc	cttttgaaag	attgaccttt	43440
ctgacttacc	cagataccat	aaagaaaata	aaatcttatc	acttcagtca	aggataaagt	43500
atttctgaat	taaaggaaaa	atacaccaga	gtaaaatcaa	gactgaaaga	caaactggga	43560
aattatttgc	aacctagatc	atagaaaagg	ggtcatttcc	ttcttgcgta	aagtgcactt	43620
acaaattgat	aagaagatga	ctgataacta	gaaagaaaaa	tgggtaaaga	acaacaatag	43680
acatttcaca	tttaacctca	ttcatgataa	ggtaagtgca	aatgaaaact	acaggggata	43740
cctttttttt	tttttaatcc	attagattgg	caaacatccc	aagggttgat	cataggctca	43800
gtgggtgaga	tttaagtatt	atcaggcatt	tttatacttt	gctgttagga	atgcaatgta	43860
gtacaaacct	ttgtagaagt	tgttttgtaa	atgtctctca	gatgtacaaa	tgcattcaca	43920
ttttagattt	agcattcccg	ctttctgaga	cattattcaa	catgtatacg	tgtgcacata	43980
agatataata	ataacacgtt	tttcttcta	gtgtgttgct	tttaacctgt	agcttgaaaa	44040
aactctgctt	tcattgtttt	tttttgtttt	ctgtcactgg	ctcagccctg	ctttcaattg	44100
tttatatgaa	ttgatgggtg	ttctggtctg	gttataatct	actttagttt	aagagtcact	44160
ttaaattata	tgacatctga	tataagttgt	gttaggtaga	aaattctgta	acttggaata	44220
ctgtaagtac	tttgtggcca	catttccatta	gtattaaata	ttatctctat	atatagtagg	44280
ctattttaata	ttcatatttt	atgatgcaat	taagaaataa	tttttttctg	aagttggtag	44340
attgttgata	tgccatggcc	cagtgtttct	caaagcattc	tgggggatca	ctgtttgtca	44400
gaattagctg	cagtgtattg	tgaacatgca	gggcctctgc	tccactccac	gttgctacca	44460
ggacgctctg	caggtgagag	ctgggaagct	gtagaagctg	cagtgtctaac	aaatgctaca	44520
ggaattcttg	tagtcacctt	catgaggtct	tatgttgagg	agaggcagcc	agtagtgtcc	44580
cttgcctctc	ccgttttatg	gtgtaagttt	catttttaagg	gaggtataaa	tcaaagccca	44640
cctgggcatt	ctctcatggt	tcactgcttc	ttgtaatcat	ggaagatgtc	attgcggcag	44700
agacgaaaca	gtgtagtttg	attactattg	attttttttt	aattattttt	ctgaagtggc	44760
tgttgtaatg	taataaattg	tgtgcttaag	gacaaccttt	ggtattctat	ttgagtattg	44820
tgtatgatcc	tagttaagtt	ttttctacca	gtattttcat	attacaacat	atttactttc	44880
catttctatt	aatattttta	tattttaagt	atggaggccg	ggcacagtg	ctcacgcgtg	44940
taatcccagc	attttgggat	gctgagggcg	gtggatcaca	aggtcaggag	ttctagacca	45000
gcgtgaccaa	cacggtgaaa	tcccatctct	actaaaaata	caaaaattag	ccgggcacag	45060
tggtaggcac	ctgtaattcc	agctactcag	gaggctgagg	taggagaatc	acttgaatcc	45120

gggaggcagc	agttgcagtg	agctaagatc	gtgccactgg	actctagcct	ggctgacaga	45180
gcaagaatcc	gcctaaaaaa	aaagggatca	gggaagaggg	gattacagat	aacccaaaga	45240
agaaggaaaa	atctccacaa	gttcacctgt	ccagcggtaa	ccccaatgtg	gatatcttcc	45300
tttaacaatt	tggatatatt	cctttaaatc	ctctttttta	taatgtctat	atgttgagaga	45360
gagtatgtgc	ctttacgtat	tttttaaaga	tgagatttct	gtgtgtgtct	atatctcctg	45420
ttcttcata	tttcttgtgt	gttataaaca	gctgtacatg	tcagtatata	tacttccgta	45480
actttttttt	aaaggctata	tagtgttcat	tgatgtgatt	taacagcagt	tatctccccg	45540
gcttcatctt	gttggaatgt	gggtcctgtg	tggtgccttc	agagcaaagt	gggcttggtt	45600
ttgcagcaag	tagacctgtg	acctgtacga	atagttggaa	gactttctct	attaccaag	45660
cgtatcagta	tacttttagtg	cctactagaa	atztatgggt	agaaaaacaa	taatatctta	45720
gagtattttt	tcctagattc	cctaagggtg	tatagggtga	tttttactca	tgtaacatga	45780
actatgcttc	aactaagata	gtttttgcaa	atgtggatat	ataagtactt	tattaaacct	45840
ataggaagta	tttataccac	ttatttcctc	ccttcagtg	tagaacctcc	taaatggcat	45900
ttgacattga	actgctttcc	actttgtcgc	atgctcctct	cattgtccct	acctgggtcc	45960
tgaaccttag	ggacttggtc	gttatagccc	caccatggct	acgctgggcc	ttggctcgtc	46020
ctgagactta	gtttcttcat	cttacaagga	gataataaca	gcccctgcct	gcgtagaatt	46080
gcagagatca	aatgaaaata	ttaacatact	caaaagcatg	cagtaaacac	attctgagca	46140
catgtacgtt	ttaggaaaaa	caaaaggacc	catgcacatt	tcggagtgtc	tttgtctcag	46200
cagcactgcc	tcttcttcca	aagctgacgt	cttagtagag	gccctggcac	gtcctgagca	46260
ctgtactcca	cgaagcattc	tatttctgac	attcgaaatg	cagtctgttc	catcttccct	46320
acaatctgta	tgccagcact	tgaataaccg	ggatctcgca	gtgttgacca	gggtgattact	46380
taattatgga	aatgttgagg	tggagatcta	gataattcag	tgaaggcagg	aaaattggtg	46440
tcggaatctg	tctttttatg	tgtcagaaat	agaaataaga	tagggtgaga	agtaatttgt	46500
ggctaaaaaca	ctataatagc	taacacatag	tgcatactgt	gtgccaaagca	ctcctgtagg	46560
tgcttgaaat	cttctattat	tattatccct	actttataga	cttgccacct	taggcacaga	46620
gaggcggaca	gtgtgtccaa	gttaccaccag	agggtggagat	ccaggctacc	tgactccacc	46680
atgtgtgtct	ttccctaggg	cacagttgtg	ctgctaaaaa	tactttttta	gcagttcttt	46740
gattattcag	atgatagtac	tgtaggaaaa	ttaagacaaa	aataatgaaa	aattaaaatc	46800
tttatttttag	tgttttgcac	atgtattatt	aaagccagtt	tactcctgga	agtgtgtaag	46860
aatacaggg	atttttgatc	acctaaatgc	tgcatgttac	taagagctcg	acactgaagt	46920
caagaagagc	agttgcagag	agtaacttagc	aaaaacggga	agtgtgtggg	gttgaaggag	46980
caaagacaag	tcttcctcgg	acgggtggagt	gtagaattca	tcattttctca	gaacacgtct	47040
ttgaacgcag	tttcaatttg	aggccaaagg	tctcagcctc	ccactcggca	tacctcccta	47100
ccttagtcag	ctcttaaatc	ttagggaatat	ttctttgttc	ttcaaggaaac	ttaaaatagt	47160
taacattctt	acctgtccac	agggagcccc	ctacaaagaa	gggagtttct	agtctccgtt	47220
ctttcttgga	ataaataata	gcctcataacc	ttgtgcaatc	gaggctgaaa	aagactgtct	47280
ccttttttca	aataagcaag	tcttagaaac	tacagttggt	tacagggctc	atggctattc	47340
cacagtaata	attttggttc	ttttaccaat	tatataatat	gttaaaatat	ggcaagtatc	47400
aggaaagcaa	ggagtggcaa	tgattagaaa	ccaatggcca	agttagagag	gaggggcaat	47460
tgctccccca	agtttgttgt	ggctgtgtag	cagtcagtga	cgagaagctg	tgtgtcaggc	47520
gacaagcaaa	gttgaggatt	atcaggcgcc	tgtgagtgcc	cagctgtgtg	ccaggctcagg	47580
aggtgccatc	gtgagccaga	ccagcttccct	ctcggccctc	gtggagctcg	cagctcgggtg	47640
gggaggcagc	agtcaccatg	gtgacagggtg	acacactagg	atggggctgg	tggtggtagg	47700
catttgccgg	tcpccttcaga	gagggtgagta	tggacttaga	ggaggctcca	gcttccctatt	47760
cctgggctgt	ctatagcact	aaaagttgtc	acatgaaaaa	taacatttgg	tactattgat	47820
ttaacttaat	gacttatgta	attgtagtgt	acttagaaat	tataacatgc	tcttctactt	47880
cagcttgaaa	ccccaacca	ccagtttata	atcctttttt	tttaactttt	gtttattttt	47940
cctaaggaa	ctgtactttt	tcttcatttt	acaacttttt	ttgtcctgtt	acctattttt	48000
cattttttact	ttatatgacc	atgagttcta	aaatagtaaa	aaaaaagaat	tattttttgtt	48060
ctttgttaga	atctctctgc	aaagaatgtc	caaaaattca	tattcacatt	gatcgatcgc	48120
acaaaaaaga	tgtcccagaa	gaacaagaac	atatgagaag	atggctgcat	gaacgtttcg	48180
aatcaaaaga	taagttagta	acaacagttc	cagcacttcc	ggaacttcgg	ttcaactaga	48240
tttcagtata	gtcaacaatt	tgaaccaaat	gtaaatgggt	atattgtctc	agaatacat	48300
tttataaaat	caaatcaaat	tttatgcatg	tctgatcgtg	ttttaaactt	tacttgtaca	48360
aatcagctta	aaagaacttg	ttacagtggg	ccatctact	tgcattgata	gtatttcttg	48420
gacaatacta	cgtgataaca	tagcaaatta	aattaaaaac	aacaacaaaac	acacaaaaaa	48480
actttccagt	gtcagatgcc	cggacctacc	tgtcagggtca	cataaagtgg	tggtactgtg	48540
tgaggctctg	ctgttgggcc	agtgtgcgca	gaaaagcaag	ggaggggtag	aggactatgc	48600
ggacgtgcag	gtggacatga	tgtgtttata	tttgttggaa	atagaagggg	gcagttgaca	48660
gcgttatatc	caaagtgtct	tctgtgttta	attattatca	gaaatttttag	ccaattgttt	48720
tattctctaa	atatgtactt	tctgtctcaag	aaactatcat	tggtcttctt	ttccttgttt	48780

tacagtacag	tgtttttaat	taaccctcct	gggttaactt	taccaggtga	aaatgattaa	48840
aagtgttaata	gggttaacaat	gaaactttta	gcttctattt	ttcattgact	cttaactgta	48900
catgatgtaa	tgtattcagc	gagccattca	ggaccacttt	ggcccatgga	agaaatttaa	48960
aagtaagatc	tacatgtatt	gacatgaaaa	tatgttctca	gaaaaaagac	taatgtattt	49020
aatgtcctac	ttattttata	agtatttaga	atacctctgg	acatttttaa	acaatgatta	49080
ttgctagggg	gtgtgattta	taaagcaata	gaagcgcttt	ccctttctgt	ttgtgtttta	49140
gattattata	tcgggtatgt	tctgctatca	taactttaca	aatcttatgt	aatatgggaa	49200
aatgagttaa	ctatgctgtt	ttccttcttt	tacctgcctt	tctaattctg	tgggaataaa	49260
ggcgtttttg	agacagccca	gggtgtagtga	gcagtcacata	tccatggatt	ccacattcat	49320
ggattccacc	aagcacagac	caaaaatact	cagaaaaaaa	gggggctggc	tgtggtggct	49380
catgcatgta	atcccagcac	tttgggaggc	taaggcaggc	aaattgcttg	agcccagaag	49440
ttcaagacag	cctgggcaac	atggcaaaac	cctgtctcta	cagaaaaatac	aaaaatttag	49500
caggcgtgca	cctgtagtcc	cagctactca	ggaggccgag	gtgcgaggat	cacctgagcc	49560
tggaaggttg	agactgcagt	gagctatcat	tgtgccaaact	ccagcctggt	aacagagtgc	49620
cttttttcaa	aaaaaaaaaa	aaaaaaggat	ttgggaggat	atgcatatgt	tatattcaaa	49680
tacatgccat	tttattcata	tatcagggag	ttgagcatcc	tttgatcttg	gtctctgccg	49740
ggtatcctgg	gaccagcccc	ctgtcgtatc	agagggaccg	ctgtctaaga	accgctggct	49800
ctatctttga	cttctggcgg	aataggagct	ccatgtaaaa	aggaggagaa	gctgcagcgg	49860
gttattagcc	atttgtgagt	caggctactg	taaaacttta	tcaaaagttt	aaaagacaaa	49920
aagcatcctc	ataaaatgcc	ttaaaaccac	ctgttgaaat	attacatata	caattcatgt	49980
atactaatac	tagagcatat	taaagatatt	ttagaagact	agaaacttct	attaaaccaa	50040
gtttctggat	gtttccgtat	ttcatccttat	tttccaggga	cctgcataac	ttttccagcg	50100
tgtaatagct	acctgattga	tattttttga	attgaaatac	tgaagtgact	aaaatctaaa	50160
ctttttccat	tctggccata	ggatgcttat	agaattttat	gagtcaccag	atccagaaag	50220
aagaaaaaga	tttctctggga	aaagtgttaa	ttccaaatta	agtatcaaga	agactttacc	50280
atcaatgttg	atcttaagtg	gtttgactgc	aggcatgctt	atgaccgatg	ctggaaggaa	50340
gctgtatgtg	aacacctgga	tatatggaac	cctacttggc	tgctgtggg	ttactattaa	50400
agcatagaca	agtagctgtc	tccagacagt	ggatgtgtgt	acattgtcta	tttttggcgg	50460
ctgcacatga	catcaaattg	tttcttgaat	ttattaagga	gtgtaataaa	agccttggtt	50520
attgaagatt	ggataataga	atttgtgacg	aaagctgata	tgcaatggct	ttgggcaaac	50580
atacctgggt	gtacaacttt	agcatcgggg	ctgctggaag	ggtaaaagct	aaatggagtt	50640
tctcctgctc	gttccatttc	ctatgaacta	atgacaactt	gagaaggctg	ggaggattgt	50700
gtattttgca	agtcagatgg	ctgcattttt	agcatttaat	ttgcagcgta	tttcaacttt	50760
tctgttattt	tcaatttatt	acaacttgac	agctccaagc	tcttattact	aaagtattta	50820
gtatcttgca	gctagttaat	atttcatctt	ttgcttattt	ctacaagtca	gtgaaataaa	50880
ttgtatttag	gaagtgtcag	gatgttcaaa	ggaaaagggt	aaaagtgttc	atggggaaaa	50940
agctctgttt	agcacatgat	tttattgtat	tgcgttatta	gctgatttta	ctcattttat	51000
atttgcaaaa	taaaatttcta	aatattgctta	aatattgctta	atttgcacac	cctgtacaca	51060
cagaaaaatg	tataaaatat	gagaacgaag	tttaaaattg	tgactctgat	tcattatagc	51120
agaactttta	atttcccagc	tttttgaaga	tttaagctac	gctattagta	cttccctttg	51180
tctgtgccat	aagtgccttg	aaacgttaag	gttttctgtt	ttgttttggt	tttttaatat	51240
caaaagagtc	gggtgtgaacc	ttgggtggac	cccaagttca	caagattttt	aaggtgatga	51300
gagcctgcag	acattctgcc	tagattttact	agcgtgtgcc	ttttgcctgc	ttctctttga	51360
tttcacagaa	tattcattca	gaagtcgcgt	ttctgtagt	tggtggattc	ccactgggct	51420
ctggctcctt	ccttggatcc	cgctcagtggt	gctgctcagc	ggcttgacag	tagacttgct	51480
aggaagaaat	gcagagccag	cctgtgctgc	ccactttcag	agttgaactc	tttaagccct	51540
tgtgagtggt	cttcaccagc	tactgcagag	gcattttgca	tttgtctgtg	tcaagaagtt	51600
cacctttctc	agccagtga	atacagactt	aattcgtcat	gactgaacga	atttgtttat	51660
ttcccattag	gtttagtga	gctacacatt	aatatgtatc	gccttagagc	aagagctgtg	51720
ttccaggaac	cagatcacga	tttttagcca	tggaacaata	tatcccagtg	gagaagacct	51780
ttcagtggtg	actgttctat	ttttgtgtta	taattttaaac	ttcgattttc	tcatagtcct	51840
ttaagttgac	atttctgctt	actgctactg	gatttttggct	gcagaaatat	atcagtggcc	51900
cacattaaac	ataccagttg	gatcatgata	agcaaaatga	aagaaataat	gattaagggg	51960
aaattaagtg	actgtgttac	actgcttctc	ccatgccaga	gaataaactc	tttcaagcat	52020
catctttgaa	gagtcgtgtg	gtgtgaattg	gtttgtgtac	attagaatgt	atgcacacat	52080
ccatggacac	tcaggatata	gttggcctaa	taatcggggc	atgggtaaaa	cttatgaaaa	52140
tttctctcat	ctgaattgta	attttctctt	acctgtaaag	taaaatttag	atcaattcca	52200
tgtctttgtt	aagtacaggg	atttaatata	ttttgaatat	aatgggtatg	ttctaaattt	52260
gaacttttag	aggcaatact	gttggaaatta	tgtggattct	aactcatttt	aacaaggtag	52320
cctgacctgc	ataagatcac	ttgaatgtta	ggtttctatg	aactatacta	atcttctcac	52380
aaaaggtcta	taaaatacac	tcgttgaaaa	aaattttgta	tcaaaatggt	tggaaaaatta	52440

gaagcttctc	cttaacctgt	attgatactg	acttgaatta	ttttctaaaa	ttaagagccg	52500
tatacctacc	tgtaagtctt	ttcacatatc	atttaaactt	ttgtttgtat	tattactgat	52560
ttacagctta	ggtattaatt	tttctttata	agaatgccgt	cgatgtgcat	gcttttatgt	52620
ttttcagaaa	aggggtgtgt	tggtatgaaag	taaaaaaaaa	aataaaatct	ttcactgtct	52680
ctaattggctg	tgctgtttta	cattttttga	ccctaaaatt	caccaacagt	ctcccagtac	52740
ataaaatagg	cttaatgact	ggccctgcat	tcttcacaat	atttttccct	aagctttgag	52800
caaagtttta	aaaaaataca	ctaaaataat	caaaactggt	aagcagtata	ttagtttggt	52860
tatataaatt	catctgcaat	ttataagatg	catggccgat	gttaatttgc	ttggcaattc	52920
tgtaatcatt	aagtgatctc	agtgaacat	gtcaaatgcc	ttaaattaac	taagttgggt	52980
aataaaagtg	ccgatctggc	taactcttac	accatacata	ctgatagtgt	ttcatatgtt	53040
tcattttccat	gtgattttta	aaatttagag	tggcaacaat	tttgcttaat	atgggttaca	53100
taagctttat	tttttccttt	gttcataatt	atattctttg	aataggtctg	tgtcaatcaa	53160
gtgatctaac	tagactgac	atagatagaa	ggaaataagg	ccaagttcaa	gaccagcctg	53220
ggcaacatat	cgagaacctg	tctacaaaaa	aattaaaaaa	aattagccag	gcatggtggc	53280
gtacactgag	tagtttgtcc	cagctactcg	ggagggtgag	gtgggaggat	cgcttcagcc	53340
caggaggttg	agattgcagt	gagccatgga	cataccactg	cactacagcc	taggtaacag	53400
cacgagaccc	caactcttag	aaaatgaaaa	ggaaatatag	aaatataaaa	tttgcttatt	53460
atagacacac	agtaactccc	agatatgtac	cacaaaaaat	gtgaaaagag	agagaaatgt	53520
ctaccaaagc	agtattttgt	gtgtataatt	gcaagcgcat	agtaaaataa	ttttaacctt	53580
aatttgtttt	tagtagtggt	taagttagaa	gttgagttaa	atattttctt	ggcagatatt	53640
ccgtatctgg	tggaaagcta	caatgcaatg	tcgttgtagt	tttgcatggc	ttgctttata	53700
aacaagattt	tttctccctc	cttttggggc	agttttcatt	acgagtaact	cacacttttt	53760
gattaaagaa	cttgaaatta	cgttatcact	tagtataatt	gacattatat	agagactatg	53820
taacatgcaa	tcattagaat	caaaattagt	actttgggtc	aaatatttac	aacattcaca	53880
tacttgtcaa	atattcatgt	aattaaactga	attttaaacc	ttcaactatt	atgaagtgtc	53940
cgtctgtaca	atcgctaatt	tactcagttt	agagtagcta	caactcttcg	atactatcat	54000
caatatttga	catcttttcc	aatttgtgta	tgaaaagtaa	atctattcct	gtagcaactg	54060
gggagtcata	tatgaggtca	aagacatata	ccttgttatt	ataatatgta	tactataata	54120
atagctgggt	atcctgagca	ggggaaaagg	ttatttttag	gaaaaccact	tcaaatagaa	54180
agctgaagta	cttctaatat	actgagggaa	gtataatatg	tggaaacaa	tctcaacaaa	54240
atgtttattg	atgttgatga	aacagatcag	tttttccatc	cggattatta	ttggttcatg	54300
attttatatg	tgaatatgta	agatatgttc	tgcaatttta	taaatgttca	tgtctttttt	54360
taaaaaagg	gctattgaaa	ttctgtgtct	ccagcaggca	agaatacttg	actaactcct	54420
tttgtctctt	tafggtattt	tcagaataaa	gtctgacttg	tgtttttgag	attatttggt	54480
cctcattaat	tcagcaataa	aggaaaatat	gcatctcaaa	aattggtgat	aaaaagttat	54540
ttcttgata	tgtgataaag	tttacatggt	gtgtatatat	gttgatttgc	caaatacggc	54600
tattaaatac	tapgtcatat	tttaaagggt	cagttttag	tgatagtaaa	caagcagtg	54660
actaagcctc	ttgcgggcat	catctcatct	cactgtcatc	acaaacccca	tgccacagcg	54720
tagcttgacc	actaaaagta	atgcatctgc	aagcatactg	ccagggtttg	gatagtttgt	54780
accaacagtt	accttatcaa	ggtaaatccc	agactctaaa	agagttgggt	ctgtgtcact	54840
acatgcataa	ctttaaataa	atttccctgc	ggcgccggtg	gctcacgcct	gtaatcccag	54900
cagttttggga	ggccgaggga	agtggatcac	ttgaggtcag	gagtttgaga	ccagcctggc	54960
caacgtgggt	aaacctgtgc	tctactaaaa	atacaaaaat	tagccaggcg	tgtggtggca	55020
ggcacctgta	atccagctga	cttgggagga	tgaggcagga	gaatcatttg	aatcctgcag	55080
gcgaggttg	cagtgaagca	agatggcgtc	attgcactcc	agcctgggcg	acaagagcga	55140
gactccgtat	taaaaaaaaa	aaaaaaaaaa	aaaaaaaaatt	cctctcctgt	ttgagctttc	55200
ccttacctgt	aaagagggga	gaatatgtat	ttacttcaaa	gagttcaggg	aaatgactct	55260
cactagtttg	agattctagg	tataaaaata	cattcttata	taattttaac	accaatgtga	55320
gagattatta	ttcttgctaa	accaattcag	ttttatttgc	tgtctaaaa	gtgtgaataa	55380
gtaattgtcc	attattttct	gaagtgtttt	ggaactcaac	acatgattgt	gaggaggatt	55440
tgttgctaaa	catctttctg	gttattcaag	ctcgtgtata	ctgtgctctg	ttgagacatg	55500
cagagtact	ttctgtctgg	gtcacaggtc	agttcttgat	agttttcgga	caattaacca	55560
gttttcattt	gcccagacc	acctttatct	tttttcttca	actgcaccca	tcttttataa	55620
ggtctttcag	tttattgcag	agaagatggt	ggagaaaagc	cggaattccc	accaccgct	55680
gccatcccca	tgttttatca	ttggctagag	tggaaaatag	cagtaactac	tgtgagagat	55740
catttgttta	tataatggaa	acaaagatga	ggaaagaacc	tggcttagat	cagagaactg	55800
atgtatttag	attctttttt	tttttttttt	taagacggag	tgttgctctg	ttgccagac	55860
tggagtacag	tggtcctatc	tcggctcact	gcaacctcca	tttccctggt	tcaagcaatt	55920
atcctgcctc	agcctcccaa	gtatttggtg	ttacaggcgt	gttccaccac	acctggctaa	55980
ttttttgtat	ttttagtaga	gacggggttt	cgccatgttg	gccaggctgg	tctcgaaatc	56040
ctgacctcag	atgatccacc	cgccttggtc	tcccaaagtg	ctgggattac	aggcgcgagc	56100

caccgcgcct ggcccaatgt atttggattc ttaaagaaca ctttcaaatt aaatatcagt 56160  
 tgaagagaac tagaactaaa gaatttctgt gtcaaactgt ttagcaaatg taagtagaag 56220  
 ctgggagatg tgccttgaa tgaatgaata catcagtaaa ataccatacg tatgttatga 56280  
 tgttattgtt tctttgcctt gggtgatttg gttttactgt gaaataattt tcaatataga 56340  
 attgtgatcg ttggaatttg gtcacttact agaaaatgag aaagaagtta atagctatct 56400  
 tccttaaaga ttcttgaggt tgggattaag gtagtggtcc caaggtgttc taaaacggca 56460  
 gcgagagctg tgcactcact tcacaaattt gaattcctgc tctgtgttag gcgctg 56516  
 <210> 2  
 <211> 23  
 <212> DNA  
 <213> Homo sapiens  
 <400> 2  
 ggctgtccag cgcttggtag aag 23  
 <210> 3  
 <211> 5227  
 <212> DNA  
 <213> Homo sapiens  
 <220>  
 <221> polyA\_signal  
 <222> 5180..5186  
 <223> AATAAA  
 <400> 3  
 ctgctgtccc tgggtgtcca cacgtactcc atg cgc tac ctg ctg ccc agc gtc 54  
 Met Arg Tyr Leu Leu Pro Ser Val  
 1 5  
 gtg ctc ctg ggc acg gcg ccc acc tac gtg ttg gcc tgg ggg gtc tgg 102  
 Val Leu Leu Gly Thr Ala Pro Thr Tyr Val Leu Ala Trp Gly Val Trp  
 10 15 20  
 cgg ctg ctc tcc gcc ttc ctg ccc gcc cgc ttc tac caa gcg ctg gac 150  
 Arg Leu Leu Ser Ala Phe Leu Pro Ala Arg Phe Tyr Gln Ala Leu Asp  
 25 30 35 40  
 gac cgg ctc tac tgc gtc tac cag agc atg gtg ctc ttc ttc ttc gag 198  
 Asp Arg Leu Tyr Cys Val Tyr Gln Ser Met Val Leu Phe Phe Phe Glu  
 45 50 55  
 aat tac acc ggg gtc cag ata ttg cta tat gga gat ttg cca aaa aat 246  
 Asn Tyr Thr Gly Val Gln Ile Leu Leu Tyr Gly Asp Leu Pro Lys Asn  
 60 65 70  
 aaa gaa aat ata ata tat tta gca aat cat caa agc aca gtt gac tgg 294  
 Lys Glu Asn Ile Ile Tyr Leu Ala Asn His Gln Ser Thr Val Asp Trp  
 75 80 85  
 att gtt gct gac atc ttg gcc atc agg cag aat gcg cta gga cat gtg 342  
 Ile Val Ala Asp Ile Leu Ala Ile Arg Gln Asn Ala Leu Gly His Val  
 90 95 100  
 cgc tac gtg ctg aaa gaa ggg tta aaa tgg ctg cca ttg tat ggg tgt 390  
 Arg Tyr Val Leu Lys Glu Gly Leu Lys Trp Leu Pro Leu Tyr Gly Cys  
 105 110 115 120  
 tac ttt gct cag cat gga gga atc tat gta aag cgc agt gcc aaa ttt 438  
 Tyr Phe Ala Gln His Gly Gly Ile Tyr Val Lys Arg Ser Ala Lys Phe  
 125 130 135  
 aac gag aaa gag atg cga aac aag ttg cag agc tac gtg gac gca gga 486  
 Asn Glu Lys Glu Met Arg Asn Lys Leu Gln Ser Tyr Val Asp Ala Gly  
 140 145 150  
 act cca atg tat ctt gtg att ttt cca gaa ggt aca agg tat aat cca 534  
 Thr Pro Met Tyr Leu Val Ile Phe Pro Glu Gly Thr Arg Tyr Asn Pro  
 155 160 165  
 gag caa aca aaa gtc ctt tca gct agt cag gca ttt gct gcc caa cgt 582  
 Glu Gln Thr Lys Val Leu Ser Ala Ser Gln Ala Phe Ala Ala Gln Arg  
 170 175 180  
 ggc ctt gca gta tta aaa cat gtg cta aca cca cga ata aag gca act 630  
 Gly Leu Ala Val Leu Lys His Val Leu Thr Pro Arg Ile Lys Ala Thr  
 185 190 195 200

cac gtt gct ttt gat tgc atg aag aat tat tta gat gca att tat gat	678
His Val Ala Phe Asp Cys Met Lys Asn Tyr Leu Asp Ala Ile Tyr Asp	
205 210 215	
gtt acg gtg gtt tat gaa ggg aaa gac gat gga ggg cag cga aga gag	726
Val Thr Val Val Tyr Glu Gly Lys Asp Asp Gly Gly Gln Arg Arg Glu	
220 225 230	
tca ccg acc atg acg gaa ttt ctc tgc aaa gaa tgt cca aaa att cat	774
Ser Pro Thr Met Thr Glu Phe Leu Cys Lys Glu Cys Pro Lys Ile His	
235 240 245	
att cac att gat cgt atc gac aaa aaa gat gtc cca gaa gaa caa gaa	822
Ile His Ile Asp Arg Ile Asp Lys Lys Asp Val Pro Glu Glu Gln Glu	
250 255 260	
cat atg aga aga tgg ctg cat gaa cgt ttc gaa atc aaa gat aag atg	870
His Met Arg Arg Trp Leu His Glu Arg Phe Glu Ile Lys Asp Lys Met	
265 270 275 280	
ctt ata gaa ttt tat gag tca cca gat cca gaa aga aga aaa aga ttt	918
Leu Ile Glu Phe Tyr Glu Ser Pro Asp Pro Glu Arg Arg Lys Arg Phe	
285 290 295	
cct ggg aaa agt gtt aat tcc aaa tta agt atc aag aag act tta cca	966
Pro Gly Lys Ser Val Asn Ser Lys Leu Ser Ile Lys Lys Thr Leu Pro	
300 305 310	
tca atg ttg atc tta agt ggt ttg act gca ggc atg ctt atg acc gat	1014
Ser Met Leu Ile Leu Ser Gly Leu Thr Ala Gly Met Leu Met Thr Asp	
315 320 325	
gct gga agg aag ctg tat gtg aac acc tgg ata tat gga acc cta ctt	1062
Ala Gly Arg Lys Leu Tyr Val Asn Thr Trp Ile Tyr Gly Thr Leu Leu	
330 335 340	
ggc tgc ctg tgg gtt act att aaa gca tag acaagtagct gtctccagac	1112
Gly Cys Leu Trp Val Thr Ile Lys Ala *	
345 350	
agtgggatgt gctacattgt ctatttttgg cggctgcaca tgacatcaaa ttgtttcctg	1172
aatttattaa ggagtgtaaa taaagccttg ttgattgaag attggataat agaatttgtg	1232
acgaaagctg atatgcaatg gtcttgggca aacatacctg gttgtacaac ttttagcatcg	1292
gggctgctgg aagggtaaaa gctaaatgga gtttctcctg ctctgtccat ttcctatgaa	1352
ctaatacaca cttgagaagg ctgggaggat tgtgtatttt gcaagtcaga tggctgcatt	1412
tttgagcatt aatttgcagc gtatttcact ttttctgtta ttttcaattt attacaactt	1472
gacagctcca agctcttatt actaaagtat ttagtatctt gcagctagtt aatatttcat	1532
cttttgccta tttctacaag tcagtgaat aaattgtatt taggaagtgt caggatgttc	1592
aaaggaaagg gtaaaaagtg ttcattggga aaaagctctg ttttagcacat gattttattg	1652
tattgcgtta ttagctgatt ttactcattt tatatttgca aaataaattt ctaataattt	1712
ttgaaattgc ttaatttgca caccctgtac acacagaaaa tgggtataaaa tatgagaacg	1772
aagtttaaaa ttgtgactct gattcattat agcagaactt taaatttccc agctttttga	1832
agatttaagc tacgctatta gtacttccct ttgtctgtgc cataagtgtc tgaaaacgtt	1892
aaggttttct gttttgtttt gtttttttaa tatcaaaaga gtcgggtgtga accttggttg	1952
gaccccaagt tcacaagatt ttaagggtga tgagagcctg cagacattct gcctagattt	2012
actagcgtgt gccttttggc tgcttctctt tgatttcaca gaatattcat tcagaagtcg	2072
cgtttctgta gtgtgttgga ttccactgg gctctggtcc ttcccttgga tcccgtcagt	2132
ggtgctgctc agcggcttgc acgtagactt gctaggaaga aatgcagagc cagcctgtgc	2192
tgccacttt cagagttgaa ctctttaaag cccttggtgag tgggcttcac cagctactgc	2252
agaggcattt tgcatattgtc tgtgtcaaga agttcacctt ctcaagccag tgaaatacag	2312
acttaattcg tcatgactga acgaatttgt ttatttccca ttaggttttag tggagctaca	2372
cattaatatg tatcgctta gagcaagagc tgtgttccag gaaccagatc acgattttta	2432
gccatggaac aatatatccc atgggagaag accttccagt gtgaactgtt ctatttttgt	2492
gttataattt aaacttcgat ttctcatag tcctttaagt tgacatttct gcttactgct	2552
actggatttt tgctgcagaa atatatcagt ggcccacatt aaacatacca gttggatcat	2612
gataagcaaa atgaagaaa taatgattaa gggaaaatta agtgactgtg ttacactgct	2672
tctcccatgc cagagaataa actctttcaa gcatcatctt tgaagagtgc tgtggtgtga	2732
attggtttgt gtacattaga atgtatgcac acatccatgg acactcagga tatagttggc	2792
ctaataatcg gggcatgggt aaaacttatg aaaatttccct catgctgaat tgtaattttc	2852
tcctacctgt aaagtaaaat ttagatcaat tccatgtctt tgttaagtac agggatttaa	2912
tatattttga atataatggg tatgttctaa atttgaactt tgagaggcaa tactgttgga	2972

```

attatgtgga ttctaactca ttttaacaag gtagcctgac ctgcataaga tcacttgaat 3032
gttaggtttc atagaactat actaatcttc tcacaaaagg tctataaaat acagtcggtg 3092
aaaaaaattt tgtatcaaaa tgtttggaag attagaagct tctccttaac ctgtattgat 3152
actgacttga attattttct aaaattaaga gccgtatacc tacctgtaag tcttttcaca 3212
tatcatttaa acttttgttt gtattattac tgatttacag cttagttatt aatttttctt 3272
tataagaatg ccgtcgatgt gcatgctttt atgtttttca gaaaagggtg tgtttggtatg 3332
aaagtaaaaa aaaaaataaa atcttttact gtctctaatg gctgtgctgt ttaacatttt 3392
ttgaccctaa aattcaccaa cagtctccca gtacataaaa taggcttaat gactggccct 3452
gcattcttca caatattttt ccctaagctt tgagcaaaagt tttaaaaaaa tacactaaaa 3512
taatcaaaac tgtaagcag tatattagtt tggttatata aattcatctg caatttataa 3572
gatgcatggc cgtatgtaat ttgcttgga attctgtaat cattaagtga tctcagtga 3632
acatgtcaaa tgccttaaat taactaagtt ggtgaataaa agtgccgatc tggctaactc 3692
ttacaccata catactgata gtttttcata tgtttcattt ccatgtgatt tttaaaattt 3752
agagtggcaa caattttgct taatatgggt tacataagct ttattttttc ctttgttcat 3812
aattatattc tttgaatagg tctgtgtcaa tcaagtgatc taactagact gatcatagat 3872
agaaggaaat aaggccaagt tcaagaccag cctgggc'aac atatcgagaa cctgtct'aca 3932
aaaaaattaa aaaaaattag ccaggcatgg tggcgtacac tgagtagttt gtcccagcta 3992
ctcgggaggg tgaggtggga ggatcgcttc agcccaggag gttgagattg cagtgagcca 4052
tggacatacc actgcactac agcctaggta acagcacgag accccaactc ttagaaaatg 4112
aaaaggaaat atagaaatat aaaatttgct tattatagac acacagtaac tcccagatat 4172
gtaccacaaa aaatgtgaaa agagagagaa atgtctacca aagcagtatt ttgtgtgtat 4232
aattgcaagc gcatagtaaa ataattttta ccttaatttg ttttagtag tgtttagatt 4292
gaagattgag tgaaatattt tcttggcaga tattccgtat ctggtggaaa gctacaatgc 4352
aatgtcgttg tagttttgca tggcttgctt tataaacaag attttttctc cctccttttg 4412
ggccagtttt cattacgagt aactcacact ttttgattaa agaacttgaa attacgttat 4472
cacttagtat aattgacatt atatagagac tatgtaacat gcaatcatta gaatcaaaa 4532
tagtactttg gtcaaaatat ttacaacatt cacatactg tcaaataattc atgtaattaa 4592
ctgaatttaa aaccttcaac tattatgaag tgctcgctcg tacaatcgct aatttactca 4652
gttttagagta gctacaactc ttcgatacta tcatcaatat ttgacatctt ttccaatttg 4712
tgtatgaaaa gtaaattctat tcctgtagca actggggagt catatatgag gtcaaagaca 4772
tataccttgt tattataata tgtatactat aataatagct ggttatcctg agcaggggaa 4832
aaggttattt ttaggaaaac cacttcaaat agaaagctga agtacttcta atatactgag 4892
ggaagtataa tatgtggaac aaactctcaa caaaatgttt attgatgttg atgaaacaga 4952
tcagtttttc catccggatt attattgggt catgatttta tatgtgaata tgtaagatat 5012
gttctgcaat ttataaaatg ttcatgtctt tttttaaaaa aggtgctatc gaaattctgt 5072
gtctccagca ggcaagaata cttgactaac tctttttgtc tctttatggt attttcagaa 5132
taaagctcga cttgtgtttt tgagattatt ggtgcctcat taattcagca ataaaggaaa 5192
atatgcattt caaaaanaaa aaaaaaaaaa aaaaa 5227

```

<210> 4

<211> 353

<212> PRT

<213> Homo sapiens

<220>

<221> HELIX

<222> 1..33

<223> Rao and Argos identification method, potential helix

<221> HELIX

<222> 4..20

<223> Klein, Kanehisa and DeLisi identification method, potential helix

<221> HELIX

<222> 4..24

<223> Eisenberg, Schwarz, Komarony, Wall identification method, potential helix

<221> MYRISTATE

<222> 12..16

<223> Prosite match

<221> HELIX

<222> 50..70

<223> Eisenberg, Schwarz, Komarony, Wall identification method, potential helix

<221> CARBOHYD



<222> 57..59  
<223> Prosite match  
<221> HELIX  
<222> 76..96  
<223> Eisenberg, Schwarz, Komarony, Wall identification method,  
potential helix  
<221> PHOSPHORYLATION  
<222> 78  
<223> potential Tyrosine kinase site, Prosite match  
<221> PHOSPHORYLATION  
<222> 84  
<223> potential caseine kinase II site, Prosite match  
<221> SITE  
<222> 94..115  
<223> potential Leucine zipper site, Prosite match  
<221> MYRISTATE  
<222> 119..123  
<223> potential site, Prosite match  
<221> PHOSPHORYLATION  
<222> 133  
<223> potential protein kinase C, Prosite match  
<221> PHOSPHORYLATION  
<222> 147  
<223> potential caseine kinase II site, Prosite match  
<221> PHOSPHORYLATION  
<222> 194  
<223> potential protein kinase C, Prosite match  
<221> PHOSPHORYLATION  
<222> 215  
<223> potential Tyrosine kinase site, Prosite match  
<221> SULFATATION  
<222> 221  
<223> Prosite match  
<221> PHOSPHORYLATION  
<222> 233  
<223> potential cAMP and cGMP dependant protein kinase site, Prosite  
match  
<221> PHOSPHORYLATION  
<222> 235  
<223> potential caseine kinase II site, Prosite match  
<221> PHOSPHORYLATION  
<222> 306  
<223> potential protein kinase C, Prosite match  
<221> HELIX  
<222> 310..330  
<223> Eisenberg, Schwarz, Komarony, Wall identification method,  
potential helix  
<221> MYRISTATE  
<222> 319..323  
<223> Prosite match  
<221> MYRISTATE  
<222> 323..327  
<223> Prosite match  
<221> AMIDATION  
<222> 329  
<223> Prosite match  
<221> HELIX  
<222> 333..353  
<223> Eisenberg, Schwarz, Komarony, Wall identification method,  
potential helix  
<221> MYRISTATE

&lt;222&gt; 341..345

&lt;223&gt; Prosite match

&lt;221&gt; PHOSPHORYLATION

&lt;222&gt; 350

&lt;223&gt; potential protein kinase C, Prosite match

&lt;400&gt; 4

```

Met Arg Tyr Leu Leu Pro Ser Val Val Leu Leu Gly Thr Ala Pro Thr
1      5      10      15
Tyr Val Leu Ala Trp Gly Val Trp Arg Leu Leu Ser Ala Phe Leu Pro
20     25     30
Ala Arg Phe Tyr Gln Ala Leu Asp Asp Arg Leu Tyr Cys Val Tyr Gln
35     40     45
Ser Met Val Leu Phe Phe Phe Glu Asn Tyr Thr Gly Val Gln Ile Leu
50     55     60
Leu Tyr Gly Asp Leu Pro Lys Asn Lys Glu Asn Ile Ile Tyr Leu Ala
65     70     75     80
Asn His Gln Ser Thr Val Asp Trp Ile Val Ala Asp Ile Leu Ala Ile
85     90     95
Arg Gln Asn Ala Leu Gly His Val Arg Tyr Val Leu Lys Glu Gly Leu
100    105    110
Lys Trp Leu Pro Leu Tyr Gly Cys Tyr Phe Ala Gln His Gly Gly Ile
115    120    125
Tyr Val Lys Arg Ser Ala Lys Phe Asn Glu Lys Glu Met Arg Asn Lys
130    135    140
Leu Gln Ser Tyr Val Asp Ala Gly Thr Pro Met Tyr Leu Val Ile Phe
145    150    155    160
Pro Glu Gly Thr Arg Tyr Asn Pro Glu Gln Thr Lys Val Leu Ser Ala
165    170    175
Ser Gln Ala Phe Ala Ala Gln Arg Gly Leu Ala Val Leu Lys His Val
180    185    190
Leu Thr Pro Arg Ile Lys Ala Thr His Val Ala Phe Asp Cys Met Lys
195    200    205
Asn Tyr Leu Asp Ala Ile Tyr Asp Val Thr Val Val Tyr Glu Gly Lys
210    215    220
Asp Asp Gly Gly Gln Arg Arg Glu Ser Pro Thr Met Thr Glu Phe Leu
225    230    235    240
Cys Lys Glu Cys Pro Lys Ile His Ile His Ile Asp Arg Ile Asp Lys
245    250    255
Lys Asp Val Pro Glu Glu Gln Glu His Met Arg Arg Trp Leu His Glu
260    265    270
Arg Phe Glu Ile Lys Asp Lys Met Leu Ile Glu Phe Tyr Glu Ser Pro
275    280    285
Asp Pro Glu Arg Arg Lys Arg Phe Pro Gly Lys Ser Val Asn Ser Lys
290    295    300
Leu Ser Ile Lys Lys Thr Leu Pro Ser Met Leu Ile Leu Ser Gly Leu
305    310    315    320
Thr Ala Gly Met Leu Met Thr Asp Ala Gly Arg Lys Leu Tyr Val Asn
325    330    335
Thr Trp Ile Tyr Gly Thr Leu Leu Gly Cys Leu Trp Val Thr Ile Lys
340    345    350
Ala

```

Ala

&lt;210&gt; 5

&lt;211&gt; 364

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 5

```

Met Leu Leu Ser Leu Val Leu His Thr Tyr Ser Met Arg Tyr Leu Leu
1      5      10      15
Pro Ser Val Val Leu Leu Gly Thr Ala Pro Thr Tyr Val Leu Ala Trp
20     25     30
Gly Val Trp Arg Leu Leu Ser Ala Phe Leu Pro Ala Arg Phe Tyr Gln

```

35 40 45  
 Ala Leu Asp Asp Arg Leu Tyr Cys Val Tyr Gln Ser Met Val Leu Phe  
 50 55 60  
 Phe Phe Glu Asn Tyr Thr Gly Val Gln Ile Leu Leu Tyr Gly Asp Leu  
 65 70 75 80  
 Pro Lys Asn Lys Glu Asn Ile Ile Tyr Leu Ala Asn His Gln Ser Thr  
 85 90 95  
 Val Asp Trp Ile Val Ala Asp Ile Leu Ala Ile Arg Gln Asn Ala Leu  
 100 105 110  
 Gly His Val Arg Tyr Val Leu Lys Glu Gly Leu Lys Trp Leu Pro Leu  
 115 120 125  
 Tyr Gly Cys Tyr Phe Ala Gln His Gly Gly Ile Tyr Val Lys Arg Ser  
 130 135 140  
 Ala Lys Phe Asn Glu Lys Glu Met Arg Asn Lys Leu Gln Ser Tyr Val  
 145 150 155 160  
 Asp Ala Gly Thr Pro Met Tyr Leu Val Ile Phe Pro Glu Gly Thr Arg  
 165 170 175  
 Tyr Asn Pro Glu Gln Thr Lys Val Leu Ser Ala Ser Gln Ala Phe Ala  
 180 185 190  
 Ala Gln Arg Gly Leu Ala Val Leu Lys His Val Leu Thr Pro Arg Ile  
 195 200 205  
 Lys Ala Thr His Val Ala Phe Asp Cys Met Lys Asn Tyr Leu Asp Ala  
 210 215 220  
 Ile Tyr Asp Val Thr Val Val Tyr Glu Gly Lys Asp Asp Gly Gly Gln  
 225 230 235 240  
 Arg Arg Glu Ser Pro Thr Met Thr Glu Phe Leu Cys Lys Glu Cys Pro  
 245 250 255  
 Lys Ile His Ile His Ile Asp Arg Ile Asp Lys Lys Asp Val Pro Glu  
 260 265 270  
 Glu Gln Glu His Met Arg Arg Trp Leu His Glu Arg Phe Glu Ile Lys  
 275 280 285  
 Asp Lys Met Leu Ile Glu Phe Tyr Glu Ser Pro Asp Pro Glu Arg Arg  
 290 295 300  
 Lys Arg Phe Pro Gly Lys Ser Val Asn Ser Lys Leu Ser Ile Lys Lys  
 305 310 315 320  
 Thr Leu Pro Ser Met Leu Ile Leu Ser Gly Leu Thr Ala Gly Met Leu  
 325 330 335  
 Met Thr Asp Ala Gly Arg Lys Leu Tyr Val Asn Thr Trp Ile Tyr Gly  
 340 345 350  
 Thr Leu Leu Gly Cys Leu Trp Val Thr Ile Lys Ala  
 355 360

<210> 6  
 <211> 26  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> misc\_binding  
 <222> 1..26  
 <223> primer oligonucleotide GC1.5p.1  
 <400> 6  
 ctgtccctgg tgctccacac gtactc  
 <210> 7  
 <211> 26  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> misc\_binding  
 <222> 1..26  
 <223> primer oligonucleotide GC1.5p.2  
 <400> 7  
 tgggtgctcca cacgtactcc atgcgc

26

26

```

<210> 8
<211> 27
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..27
<223> primer oligonucleotide pg15RACE196
<400> 8
caatatctgg accccggtgt aattctc
<210> 9
<211> 34
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..34
<223> primer oligonucleotide GC1.3p
<400> 9
cttgccctgct ggagacacag aatttcgata gcac
<210> 10
<211> 24
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..24
<223> primer oligonucleotide PGRT32
<400> 10
tttttttttt tttttttttg aaat
<210> 11
<211> 6
<212> PRT
<213> Homo sapiens
<220>
<221> SITE
<222> 160..165
<223> box2 from SEQID4, present in AF003136, P33333, P26647, U89336,
U56417, AB005623.
<400> 11
Phe Pro Glu Gly Thr Arg
1 5
<210> 12
<211> 6
<212> PRT
<213> Homo sapiens
<220>
<221> SITE
<222> 129..134
<223> box2 from Z72511
<400> 12
Phe Pro Glu Gly Thr Asp
1 5
<210> 13
<211> 6
<212> PRT
<213> Homo sapiens
<220>
<221> SITE
<222> 223..228
<223> box2 from P38226, Z49770

```

<400> 13  
Phe Pro Glu Gly Thr Asn  
1 5  
<210> 14  
<211> 6  
<212> PRT  
<213> Homo sapiens  
<220>  
<221> SITE  
<222> 90..95  
<223> box2 from Z49860 and Z29518  
<400> 14  
Phe Val Glu Gly Thr Arg  
1 5  
<210> 15  
<211> 9  
<212> PRT  
<213> Homo sapiens  
<220>  
<221> SITE  
<222> 211..219  
<223> box3 from SEQID4, present in AF003136  
<400> 15  
Leu Asp Ala Ile Tyr Asp Val Thr Val  
1 5  
<210> 16  
<211> 9  
<212> PRT  
<213> Homo sapiens  
<220>  
<221> SITE  
<222> 204..212  
<223> box3 from Z72511  
<400> 16  
Val Glu Tyr Ile Tyr Asp Ile Thr Ile  
1 5  
<210> 17  
<211> 9  
<212> PRT  
<213> Homo sapiens  
<220>  
<221> SITE  
<222> 271..279  
<223> box3 from P38226  
<400> 17  
Ile Glu Ser Leu Tyr Asp Ile Thr Ile  
1 5  
<210> 18  
<211> 9  
<212> PRT  
<213> Homo sapiens  
<220>  
<221> SITE  
<222> 265..273  
<223> box3 from Z49770  
<400> 18  
Leu Asp Ala Ile Tyr Asp Val Thr Ile  
1 5  
<210> 19  
<211> 9  
<212> PRT

```

<213> Homo sapiens
<220>
<221> SITE
<222> 138..146
<223> box3 fromZ49860
<400> 19
Val Pro Ala Ile Tyr Asp Met Thr Val
1      5
<210> 20
<211> 9
<212> PRT
<213> Homo sapiens
<220>
<221> SITE
<222> 218..225
<223> box3 from Z29518
<400> 20
Val Pro Ala Ile Tyr Asp Thr Thr Val
1      5
<210> 21
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-123
<221> allele
<222> 24
<223> polymorphic base C
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-123.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-123.mis2
<400> 21
tttctcatcc tcacacctca ctgcgccct cctgaaccca ctccttt
<210> 22
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-26
<221> allele
<222> 24
<223> polymorphic base G
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-26.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-26.mis2
<400> 22
ccctgtgtnaga cacgtcctgt atcgttggtg agatgggaaa gtgcatc
<210> 23
<211> 47
<212> DNA
<213> Homo Sapiens

```

47

47

```

<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-14
<221> allele
<222> 24
<223> polymorphic base T
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-14.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-14.mis2
<400> 23
gcaggagca gaccagacat gatttgttct agtctagctg attcata
<210> 24
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-77, extracted from SEQ ID1 12057 12103
<221> allele
<222> 24
<223> polymorphic base C
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-77.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-77.mis2
<400> 24
gctgttcaga cttaaacttgg agactacagt cagtcagaga acttgct
<210> 25
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-217, extracted from SEQ ID1 34469 34515
<221> allele
<222> 24
<223> polymorphic base C
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-217.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-217.mis2
<400> 25
atatagttca cgttatgttc atacttaatt gttgcatttt gtttgcc
<210> 26
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-67, extracted from SEQ ID1 51612 51658

```

<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-67.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-67.mis2  
<400> 26  
gccagtga aa tacagactta attcgtcatg actgaacgaa tttgttt 47  
<210> 27  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-213  
<221> allele  
<222> 24  
<223> polymorphic base T  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-213.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-213.mis2  
<400> 27  
ccttagcatt caagcccctg agctctggtg ttgtccaccc ctggggg 47  
<210> 28  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-221  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-221.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-221.mis2  
<400> 28  
agcttgagaa accagaaaag ccaaaaggag gctcctacca catgggt 47  
<210> 29  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-135  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind



```

<222> 1..23
<223> potential microsequencing oligo 99-135.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-135.mis2
<400> 29
agtcactata tctatgttta atgaagatag aaagagatgc agaaatg
<210> 30
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-123, variant version of SEQ ID21
<221> allele
<222> 24
<223> base T ; C in SEQ ID21
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-123.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-123.mis2
<400> 30
tttctcatcc tcacacctca ctgtgccct cctgaacca ctccttt
<210> 31
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-26, variant version of SEQ ID22
<221> allele
<222> 24
<223> base A ; G in SEQ ID22
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-26.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-26.mis2
<400> 31
ccctgtnaga cacgtcctgt atcattgttg agatgggaaa gtgcatc
<210> 32
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-14, variant version of SEQ ID23
<221> allele
<222> 24
<223> base C ; T in SEQ ID23
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-14.mis1
<221> primer_bind
<222> 25..47

```

```

<223> complement potential microsequencing oligo 4-14.mis2
<400> 32
gcagggagca gaccagacat gatctgttct agtctagctg attcata
<210> 33
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-77, variant version of SEQ ID24
<221> allele
<222> 24
<223> base G ; C in SEQ ID24
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-77.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-77.mis2
<400> 33
gctgttcaga ctaaacttgg agagtacagt cagtcagaga acttgct
<210> 34
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-217, variant version of SEQ ID25
<221> allele
<222> 24
<223> base T ; C in SEQ ID25
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-217.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-217.mis2
<400> 34
atatagtcca cgttatgttc atatttaatt gttgcatttt gtttgcc
<210> 35
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-67, variant version of SEQ ID26
<221> allele
<222> 24
<223> base T ; C in SEQ ID26
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-67.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-67.mis2
<400> 35
gccagtgaag taagactta atttgcatg actgaacgaa tttgttt
<210> 36

```

<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-213, variant version of SEQ ID27  
<221> allele  
<222> 24  
<223> base C ; T in SEQ ID27  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-213.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-213.mis2  
<400> 36  
ccttagcatt caagcccctg agccctggtg ttgtccaccc ctggggg  
<210> 37  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-221, variant version of SEQ ID28  
<221> allele  
<222> 24  
<223> base C ; A in SEQ ID28  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-221.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-221.mis2  
<400> 37  
agcttgagaa accagaaaag ccacaaggag gctcctacca catgggt  
<210> 38  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-135, variant version of SEQ ID29  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID29  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-135.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-135.mis2  
<400> 38  
agtactata tctatgttta atggagatag aaagagatgc agaaatg  
<210> 39  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>

47

47

47

<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer 99-123-PU  
<400> 39  
aaagccagga ctagaagg 18  
<210> 40  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer 4-26-PU  
<400> 40  
tacagccctg taagacac 18  
<210> 41  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer 4-14-PU  
<400> 41  
tctaacctct catccaac 18  
<210> 42  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer 4-77-PU, extracted from SEQ ID1 11930  
11947  
<400> 42  
tgttgattta caggcggc 18  
<210> 43  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> upstream amplification primer 99-217-PU, extracted from SEQ ID1 34216  
34234  
<400> 43  
ggtgggaatt tactatatg 19  
<210> 44  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer 4-67-PU, extracted from SEQ ID1 51596  
51613  
<400> 44  
aagttcacct tctcaagc 18  
<210> 45  
<211> 20  
<212> DNA

<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..20  
<223> upstream amplification primer 99-213-PU  
<400> 45  
atactggcag cgtgtgcttc  
<210> 46  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> upstream amplification primer 99-221-PU  
<400> 46  
ccctttttct tcactgttc  
<210> 47  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer 99-135-PU  
<400> 47  
tggaagttgt tattgccc  
<210> 48  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> downstream amplification primer 99-123-RP  
<400> 48  
tattcagaaa ggagtggg  
<210> 49  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> downstream amplification primer 4-26-RP  
<400> 49  
tgaggactgc taggaaag  
<210> 50  
<211> 20  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..20  
<223> downstream amplification primer 4-14-RP  
<400> 50  
gactgtatcc tttgatgcac  
<210> 51  
<211> 20  
<212> DNA  
<213> Homo Sapiens

```

<220>
<221> primer_bind
<222> 1..20
<223> downstream amplification primer 4-77-RP, extracted from SEQ ID1 12339
123
58 complement
<400> 51
ggaaaggtag tcattcatag                20
<210> 52
<211> 21
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..21
<223> downstream amplification primer 99-217-RP, extracted from SEQ ID1 34625
34645 complement
<400> 52
gtttattttg tgtgagcttt g              21
<210> 53
<211> 20
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..20
<223> downstream amplification primer 4-67-RP, extracted from SEQ ID1 51996
520
15 complement
<400> 53
tgaaagagtt tattctctgg                20
<210> 54
<211> 21
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..21
<223> downstream amplification primer 99-213-RP
<400> 54
ttattgcccc acatgcttga g              21
<210> 55
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> downstream amplification primer 99-221-RP
<400> 55
tcattcgtct ggctagggtc                19
<210> 56
<211> 18
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..18
<223> downstream amplification primer 99-135-RP
<400> 56
aaacacctcc cattgtgc                  18

```

<210> 57  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1482  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-1482.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-1482.mis2  
<400> 57  
agtgaagtct gagggggaaa aatcaaccct atagaggga g gatctg 47  
<210> 58  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-73, extracted from SED ID1 13657 13703  
<221> allele  
<222> 24  
<223> polymorphic base C in PG1 (13680) SEQ ID1  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-73.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-73.mis2  
<400> 58  
gttttcctta tgatgttaca tggcttattt ttaaaggtaa tgaaaac 47  
<210> 59  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-65, extracted from SEQ ID1 51448 51494  
<221> allele  
<222> 24  
<223> polymorphic base T in PG1 (51471) SEQ ID1  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-65.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-65.mis2  
<400> 59  
ggtgctgctc agcggcttgc acgtagactt gctaggaaga aatgcag 47  
<210> 60  
<211> 47  
<212> DNA  
<213> Homo Sapiens

```

<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-1482, variant version of SEQ ID57
<221> allele
<222> 24
<223> base A ; C in SEQ ID57
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-1482.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-1482.mis2
<400> 60
agtgaagtct gagggggaaa aataaacct atagaggga ggatctg
<210> 61
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-73, variant version of SEQ ID58
<221> allele
<222> 24
<223> base G ; C in SEQ ID58
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-73.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-73.mis2
<400> 61
gttttcctta tgatgttaca tgggttattt ttaaaggtaa tgaaaac
<210> 62
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-65, variant version of SEQ ID59
<221> allele
<222> 24
<223> base C ; T in SEQ ID59
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-65.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-65.mis2
<400> 62
ggtgctgctc agcggcttgc acgcagactt gctaggaaga aatgcag
<210> 63
<211> 21
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..21
<223> upstream amplification primer 99-1482-PU

```

47

47

47



<400> 63  
atcaaatacag tgaagtctga g 21  
<210> 64  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer 4-73-PU, extracted from SEQ ID1 13547  
13564  
<400> 64  
atcgctggaa cattctgg 18  
<210> 65  
<211> 20  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..20  
<223> upstream amplification primer 4-65-PU, extracted from SEQ ID1 51149  
51168  
<400> 65  
gatttaagct acgctattag 20  
<210> 66  
<211> 20  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..20  
<223> downstream amplification primer 99-1482-RP  
<400> 66  
acaaatctat ataaggctgg 20  
<210> 67  
<211> 20  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..20  
<223> downstream amplification primer 4-73-RP, extracted from SEQ ID1 13962  
13981 complement  
<400> 67  
ctcttggtta aacagcagtg 20  
<210> 68  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> downstream amplification primer 4-65-RP, extracted from SEQ ID1 51482  
51499 complement  
<400> 68  
tggctctgca tttcttcc 18  
<210> 69  
<211> 5226  
<212> DNA  
<213> Homo sapiens  
<400> 69

ctgctgtccc	tggtgctcca	cacgtactcc	atg	cgc	tac	ctg	ctg	ccc	agc	gtc	54
			Met	Arg	Tyr	Leu	Leu	Pro	Ser	Val	
			1				5				
gtg	ctc	ctg	ggc	acg	gcg	ccc	acc	tac	gtg	ttg	102
Val	Leu	Leu	Gly	Thr	Ala	Pro	Thr	Tyr	Val	Leu	
10						15				20	
cgg	ctg	ctc	tcc	gcc	ttc	ctg	ccc	gcc	cgc	ttc	150
Arg	Leu	Leu	Ser	Ala	Phe	Leu	Pro	Ala	Arg	Phe	
25					30					35	
gac	cgg	ctc	tac	tgc	gtc	tac	cag	agc	atg	gtg	198
Asp	Arg	Leu	Tyr	Cys	Val	Tyr	Gln	Ser	Met	Val	
				45					50		
aat	tac	acc	ggg	gtc	cag	ata	ttg	cta	tat	gga	246
Asn	Tyr	Thr	Gly	Val	Gln	Ile	Leu	Tyr	Gly	Asp	
			60				65			70	
aaa	gaa	aat	ata	ata	tat	tta	gca	aat	cat	caa	294
Lys	Glu	Asn	Ile	Ile	Tyr	Leu	Ala	Asn	His	Gln	
75						80				85	
att	gtt	gct	gac	atc	ttg	gcc	atc	agg	cag	aat	342
Ile	Val	Ala	Asp	Ile	Leu	Ala	Ile	Arg	Gln	Asn	
90						95				100	
cgc	tac	gtg	ctg	aaa	gaa	ggg	tta	aaa	tggt	ctg	390
Arg	Tyr	Val	Leu	Lys	Glu	Gly	Leu	Lys	Trp	Leu	
105					110				115		
tac	ttt	gct	cag	cat	gga	atc	tat	gta	aag	cgc	438
Tyr	Phe	Ala	Gln	His	Gly	Gly	Ile	Tyr	Val	Lys	
				125					130		
aac	gag	aaa	gag	atg	cga	aac	aag	ttg	cag	agc	486
Asn	Glu	Lys	Glu	Met	Arg	Asn	Lys	Leu	Gln	Ser	
				140					145		
act	cca	atg	tat	ctt	gtg	att	ttt	cca	gaa	ggt	534
Thr	Pro	Met	Tyr	Leu	Val	Ile	Phe	Pro	Glu	Gly	
				155			160			165	
gag	caa	aca	aaa	gtc	ctt	tca	gct	agt	cag	gca	582
Glu	Gln	Thr	Lys	Val	Leu	Ser	Ala	Ser	Gln	Ala	
170						175				180	
ggc	ctt	gca	gta	tta	aaa	cat	gtg	cta	aca	cca	630
Gly	Leu	Ala	Val	Leu	Lys	His	Val	Leu	Thr	Pro	
185					190				195		
cac	gtt	gct	ttt	gat	tgc	atg	aag	aat	tat	tta	678
His	Val	Ala	Phe	Asp	Cys	Met	Lys	Asn	Tyr	Leu	
				205					210		
gtt	acg	gtg	gtt	tat	gaa	ggg	aaa	gac	gat	gga	727
Val	Thr	Val	Val	Tyr	Glu	Gly	Lys	Asp	Asp	Gly	
				220					225		
caccgacat	gacggaattt	ctctgcaaag	aatgtccaaa	aattcatatt	cacattgatc	787					
gtatcgacaa	aaaagatgtc	ccagaagaac	aagaacatat	gagaagatgg	ctgcatgaac	847					
gtttcgaaat	caaagataag	atgcttatag	aattttatga	gtcaccagat	ccagaaagaa	907					
gaaaaagatt	tcttgggaaa	agtgttaatt	ccaaattaag	tatcaagaag	actttaccat	967					
caatgttgat	cttaagtggg	ttgactgcag	gcatgcttat	gaccgatgct	ggaaggaagc	1027					
tgtatgtgaa	cacctggata	tatggaaccc	tacttggtg	cctgtgggtt	actattaaag	1087					
catagacaag	tagctgtctc	cagacagtgg	gatgtgctac	attgtctatt	tttgccggct	1147					
gcacatgaca	tcaaattggt	tctgaatttt	attaaggagt	gtaaaataaag	cctgtgtgat	1207					
tgaagattgg	ataatagaat	ttgtgacgaa	agctgatatg	caatgggtctt	gggcaaacat	1267					
acctggttgt	acaacttttag	catcggggct	gctggaaggg	taaaagctaa	atggagtttc	1327					
tctgtctctg	tccatttcct	atgaactaat	gacaacttga	gaaggctggg	aggattgtgt	1387					
attttgcaag	tcagatggct	gcatttttga	gcattaattt	gcagcgtatt	tcactttttc	1447					
tgattatttc	aattttattac	aacttgacag	ctccaagctc	ttattactaa	agtatttagt	1507					
atcttgcagc	tagttaatat	ttcatctttt	gcttattttc	acaagtcagt	gaaataaatt	1567					
gtatttagga	aggtgcagga	tgttcaaagg	aaagggtaaa	aagtgttcat	ggggaaaaag	1627					
ctctgttttag	capatgattt	tattgtattg	cgttatttagc	tgatttttact	catttttatat	1687					

ttgcaaaata	aattttcta	attttattgaa	attgctta	ttgcacaccc	tgtacacaca	1747
gaaaatggta	taaaatatga	gaacgaagtt	taaaattgtg	actctgattc	attatagcag	1807
aacttttaaat	ttcccagctt	tttgaagatt	taagctacgc	tatttagtact	tccctttgtc	1867
tgtgccataa	gtgcttgaaa	acgttaaggt	tttctgtttt	gttttgtttt	tttaatatca	1927
aaagagtcgg	tgfgaacctt	ggttggaccc	caagttcaca	agatttttaa	ggtgatgaga	1987
gcctgcagac	attctgccta	gatttactag	cgtgtgcctt	ttgcctgctt	ctctttgatt	2047
tcacagaata	ttcattcaga	agtcgcgttt	ctgtagtgtg	gtggattccc	actgggctct	2107
ggtccttccc	ttggatcccc	tcagtgtgtc	tgctcagcgg	cttgacagta	gacttgctag	2167
gaagaaatgc	agagccagcc	tgtgctgccc	actttcagag	ttgaactcct	taagcccttg	2227
tgagtgggct	tcaccagcta	ctgcagaggc	attttgcatt	tgtctgtgtc	aagaagttca	2287
cctttctcaag	ccagtggaaat	acagacttaa	ttcgtcatga	ctgaacgaat	ttgtttattt	2347
cccattaggt	ttagtggagc	tacacattaa	tatgtatcgc	cttagagcaa	gagctgtgtt	2407
ccaggaacca	gatcacgatt	tttagccatg	gaacaatata	tcccatggga	gaagaccttt	2467
cagtgtgaac	tgttctattt	ttgtgttata	attttaaactt	cgatttcctc	atagtccttt	2527
aagttgacat	ttctgcttac	tgctactgga	tttttgcctg	agaaatata	cagtggccca	2587
cattaaacat	accagtggga	tcattgataag	caaaaatgaaa	gaaataatga	ttaggggaaa	2647
attaagtgc	tgtgttacac	tgcttctccc	atgccagaga	ataaactcct	tcaagcatca	2707
tctttgaaga	gtcgtgtggt	gtgaattggt	ttgtgtacat	tagaatgtat	gcacacatcc	2767
atggacactc	aggatatagt	tggtcctaata	atcggggcat	gggtaaaact	tatgaaaatt	2827
tcctcatgct	gaattgtaat	tttctcttac	ctgtaaagta	aaatttagat	caattccatg	2887
tctttgttaa	gtacagggat	ttaatatatt	ttgaatataa	tggttatgtt	ctaaatttga	2947
actttgagag	gcaataactgt	tggaattatg	tggtattctaa	ctcatttttaa	caaggtagcc	3007
tgacctgcat	aagatcactt	gaatgttagg	tttcatagaa	ctataactaat	cttctcacia	3067
aaggtctata	aaatacagtc	gttgaaaaaa	attttgtatc	aaaatgtttg	gaaaattaga	3127
agcttctcct	taacctgtat	tgatactgac	ttgaattatt	ttctaaaatt	aagagccgta	3187
tacctacctg	taagtctttt	cacatatcat	ttaaactttt	gtttgtatta	ttactgattt	3247
acagcttagt	tatttaatttt	tctttataag	aatgccgtcg	atgtgcatgc	ttttatgttt	3307
ttcagaaaag	ggtgtgtttg	gatgaaagta	aaaaaaaaaa	taaaatcttt	cactgtctct	3367
aatggctgtg	ctgtttaaca	ttttttgacc	ctaaaattca	ccaacagtct	cccagtacat	3427
aaaataggct	taatgactgg	ccctgcattc	ttcacaaat	ttttccctaa	gctttgagca	3487
aagtttttaa	aaaatacact	aaaataatca	aaactgttaa	gcagtatatt	agtttgggta	3547
tataaattca	tctgcaattt	ataagatgca	tggtccgatg	taatttgcct	ggcaattctg	3607
taatcattaa	gtgatctcag	tgaacatgt	caaatgcctt	aaatttaacta	agttgggtga	3667
taaaagtgcc	gatctggcta	actcttacac	catacatact	gatagttttt	catatgtttc	3727
atttccatgt	gattttttaa	atttagagtg	gcaacaattt	tgcttaatat	gggttacata	3787
agctttattt	tttcccttgt	tcataattat	attcctttgaa	taggtctgtg	tcaatcaagt	3847
gatctaacta	gactgatcat	agatagaagg	aaataaggcc	aagttaaga	ccagcctggg	3907
caacatatcg	agaacctgtc	tacaaaaaaa	ttaaaaaaa	ttagccaggc	atgggtgcgt	3967
acactgagta	gtttgtccca	gctactcggg	aggggtgaggt	gggaggatcg	cttcagccca	4027
ggaggttgag	attgcagtga	gccatggaca	taccactgca	ctacagccta	ggtaacagca	4087
cgagacccca	actcttagaa	aatgaaaagg	aaatatagaa	atataaaatt	tgcttattat	4147
agacacacag	taactcccag	atatgtacca	caaaaaatgt	gaaaagagag	agaaatgtct	4207
accaaagcag	tattttgtgt	gtataattgc	aagcgcatag	taaaataaatt	ttaaccttaa	4267
tttgttttta	gtagtgttta	gattgaagat	tgagtgaat	attttcttgg	cagatattcc	4327
gtatctgggtg	gaaagctaca	atgcaatgtc	gttgtagtgt	tgcatggcct	gctttataaa	4387
caagattttt	tctccctcct	tttgggcccag	ttttcattac	gagtaactca	cactttttga	4447
ttaaagaact	tgaattacg	ttatcactta	gtataattga	cattatatag	agactatgta	4507
acatgcaatc	attagaatca	aaattagtac	tttggtaaaa	atatttacaa	cattcacata	4567
cttgtcaaat	attcatgtaa	ttactgaat	ttaaaacctt	caactattat	gaagtgcctg	4627
tctgtacaat	cgctaattta	ctcagtttag	agtagctaca	actcttcgat	actatcatca	4687
atatttgaca	tctttttcaa	tttgtgtatg	aaaagtaaat	ctattcctgt	agcaactggg	4747
gagtcatata	tgaggtcaaa	gacatatacc	ttgttattat	aatatgtata	ctataataat	4807
agctgggttat	cctgagcagg	ggaaaaggtt	atttttagga	aaaccacttc	aaatagaaag	4867
ctgaagtact	tctaataatc	tgagggaagt	ataatatgtg	gaacaaactc	tcaacaaaat	4927
gtttattgat	gttgatgaaa	cagatcagtt	tttccatccg	gattattatt	ggttcatgat	4987
tttatatgtg	aatatgtaag	atatgttctg	caattttata	aatgttcatg	tcttttttta	5047
aaaaaggtgc	tatcgaaatt	ctgtgtctcc	agcaggcaag	aatacttgac	taactctttt	5107
tgtctcttta	tggtattttc	agaataaagt	ctgacttgtg	tttttgagat	tattgggtgcc	5167
tcattaaatc	agcaataaag	gaaaatatgc	atttcaaaaa	naaaaaaaaa	aaaaaaaaa	5226

&lt;210&gt; 70

&lt;211&gt; 228

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 70

```

Met Arg Tyr Leu Leu Pro Ser Val Val Leu Leu Gly Thr Ala Pro Thr
1      5      10      15
Tyr Val Leu Ala Trp Gly Val Trp Arg Leu Leu Ser Ala Phe Leu Pro
20      25      30
Ala Arg Phe Tyr Gln Ala Leu Asp Asp Arg Leu Tyr Cys Val Tyr Gln
35      40      45
Ser Met Val Leu Phe Phe Phe Glu Asn Tyr Thr Gly Val Gln Ile Leu
50      55      60
Leu Tyr Gly Asp Leu Pro Lys Asn Lys Glu Asn Ile Ile Tyr Leu Ala
65      70      75      80
Asn His Gln Ser Thr Val Asp Trp Ile Val Ala Asp Ile Leu Ala Ile
85      90      95
Arg Gln Asn Ala Leu Gly His Val Arg Tyr Val Leu Lys Glu Gly Leu
100     105     110
Lys Trp Leu Pro Leu Tyr Gly Cys Tyr Phe Ala Gln His Gly Gly Ile
115     120     125
Tyr Val Lys Arg Ser Ala Lys Phe Asn Glu Lys Glu Met Arg Asn Lys
130     135     140
Leu Gln Ser Tyr Val Asp Ala Gly Thr Pro Met Tyr Leu Val Ile Phe
145     150     155     160
Pro Glu Gly Thr Arg Tyr Asn Pro Glu Gln Thr Lys Val Leu Ser Ala
165     170     175
Ser Gln Ala Phe Ala Ala Gln Arg Gly Leu Ala Val Leu Lys His Val
180     185     190
Leu Thr Pro Arg Ile Lys Ala Thr His Val Ala Phe Asp Cys Met Lys
195     200     205
Asn Tyr Leu Asp Ala Ile Tyr Asp Val Thr Val Val Tyr Glu Gly Lys
210     215     220
Asp Asp Gly Gly
225

```

&lt;210&gt; 71

&lt;211&gt; 158

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 71

```

gccttgacagt attaaaacat gtgctaacac cacgaataaa ggcaactcac gttgcttttg      60
attgcatgaa gaattattta gatgcaattt atgatgttac ggtggtttat gaagggaag      120
acgatggagg gtagcgaaga gagtcaccga ccatgacg      158

```

&lt;210&gt; 72

&lt;211&gt; 1381

&lt;212&gt; DNA

&lt;213&gt; Mus musculus

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; 608..629

&lt;223&gt; amplification primer g34292.pu

&lt;221&gt; misc\_binding

&lt;222&gt; 740..758

&lt;223&gt; amplification primer g34292.rp

&lt;400&gt; 72

```

gagccgagag gatgctgctg tccctggtgc tccacacgta ctct atg cgc tac ctg      56
                                     Met Arg Tyr Leu
                                     1
ctc ccc agc gtc ctg ttg ctg ggc tgc gcg ccc acc tac ctg ctg gcc      104
Leu Pro Ser Val Leu Leu Leu Gly Ser Ala Pro Thr Tyr Leu Leu Ala
5      10      15      20
tgg acg ctg tgg cgg gtg ctc tcc gcg ctg atg ccc gcc cgc ctg tac      152
Trp Thr Leu Trp Arg Val Leu Ser Ala Leu Met Pro Ala Arg Leu Tyr

```

	25	30	35	
cag cgc gtg gac gac cgg ctt tac tgc gtc tac cag aac atg gtg ctc				200
Gln Arg Val Asp Arg Leu Tyr Cys Val Tyr Gln Asn Met Val Leu				
	40	45	50	
ttc ttc ttc gag aac tac acc ggg gtc cag ata ttg cta tat gga gat				248
Phe Phe Phe Glu Asn Tyr Thr Gly Val Gln Ile Leu Leu Tyr Gly Asp				
	55	60	65	
ttg cca aaa aat aaa gaa aat gta ata tat cta gcg aat cat caa agc				296
Leu Pro Lys Asn Lys Glu Asn Val Ile Tyr Leu Ala Asn His Gln Ser				
	70	75	80	
aca gtt gac tgg att gtt gcg gac atg ctg gct gcc aga cag gat gcc				344
Thr Val Asp Trp Ile Val Ala Asp Met Leu Ala Ala Arg Gln Asp Ala				
	85	90	95	100
cta gga cat gtg cgc tac gta ctg aaa gac aag tta aaa tgg ctt ccg				392
Leu Gly His Val Arg Tyr Val Leu Lys Asp Lys Leu Lys Trp Leu Pro				
	105	110	115	
ctg tat ggg ttc tac ttt gct cag cat gga gga att tat gta aaa cga				440
Leu Tyr Gly Phe Tyr Phe Ala Gln His Gly Gly Ile Tyr Val Lys Arg				
	120	125	130	
agt gcc aaa ttt aat gat aaa gaa atg aga agc aag ctg cag agc tat				488
Ser Ala Lys Phe Asn Asp Lys Glu Met Arg Ser Lys Leu Gln Ser Tyr				
	135	140	145	
gtg aac gca gga aca ccg atg tat ctt gtg att ttc cca gag gga aca				536
Val Asn Ala Gly Thr Pro Met Tyr Leu Val Ile Phe Pro Glu Gly Thr				
	150	155	160	
agg tat aat gca aca tac aca aaa ctc ctt tca gcc agt cag gca ttt				584
Arg Tyr Asn Ala Thr Tyr Thr Lys Leu Leu Ser Ala Ser Gln Ala Phe				
	165	170	175	180
gct gct cag cgg ggc ctt gca gta tta aaa cac gta ctg aca cca aga				632
Ala Ala Gln Arg Gly Leu Ala Val Leu Lys His Val Leu Thr Pro Arg				
	185	190	195	
ata aag gcc act cac gtt gct ttt gat tct atg aag agt cat tta gat				680
Ile Lys Ala Thr His Val Ala Phe Asp Ser Met Lys Ser His Leu Asp				
	200	205	210	
gca att tat gat gtc aca gtg gtt tat gaa ggg aat gag aaa ggt tca				728
Ala Ile Tyr Asp Val Thr Val Tyr Glu Gly Asn Glu Lys Gly Ser				
	215	220	225	
gga aaa tac tca aat cca cca tcc atg act gag ttt ctc tgc aaa cag				776
Gly Lys Tyr Ser Asn Pro Pro Ser Met Thr Glu Phe Leu Cys Lys Gln				
	230	235	240	
tgc cca aaa ctt cat att cac ttt gat cgt ata gac aga aat gaa gtt				824
Cys Pro Lys Leu His Ile His Phe Asp Arg Ile Asp Arg Asn Glu Val				
	245	250	255	260
cca gag gaa caa gaa cac atg aaa aag tgg ctt cat gag cgc ttt gag				872
Pro Glu Glu Gln Glu His Met Lys Lys Trp Leu His Glu Arg Phe Glu				
	265	270	275	
ata aaa gat agg ttg ctc ata gag ttc tat gat tca cca gat cca gaa				920
Ile Lys Asp Arg Leu Leu Ile Glu Phe Tyr Asp Ser Pro Asp Pro Glu				
	280	285	290	
aga aga aac aaa ttt cct ggg aaa agt gtt cat tcc aga cta agt gtg				968
Arg Arg Asn Lys Phe Pro Gly Lys Ser Val His Ser Arg Leu Ser Val				
	295	300	305	
aag aag act tta cct tca gtg ttg atc ttg ggg agt ttg act gcg gtc				1016
Lys Lys Thr Leu Pro Ser Val Leu Ile Leu Gly Ser Leu Thr Ala Val				
	310	315	320	
atg ctg atg acg gag tcc gga agg aaa ctg tac atg ggc acc tgg ttg				1064
Met Leu Met Thr Glu Ser Gly Arg Lys Leu Tyr Met Gly Thr Trp Leu				
	325	330	335	340
tat gga acc ctc ctt ggc tgc ctg tgg ttt gtt att aaa gca taa				1109
Tyr Gly Thr Leu Leu Gly Cys Leu Trp Phe Val Ile Lys Ala *				
	345	350	355	

```

gcaagtagca ggctgcagtc acagtctctt attgatggct acacattgta tcacattggt 1169
tcctgaatta aataaggagt tttcttggtg ttgttttttt tgttttggtt tgttctggtt 1229
taagccttga tgatngnnn cnnnnnnnnn ncnantcng ngaccacagc caacatgcat 1289
ttgatttggg gcaaacacat gtggcttttc aggtgctggg gttgctggag acatggaagc 1349
taagtggagt ttatgctgtt tttttttttt tt 1381

```

&lt;210&gt; 73

&lt;211&gt; 15766

&lt;212&gt; DNA

&lt;213&gt; Mus musculus

&lt;220&gt;

&lt;221&gt; exon

&lt;222&gt; 52..121

&lt;223&gt; exon2

&lt;221&gt; exon

&lt;222&gt; 682..797

&lt;223&gt; exon3

&lt;221&gt; exon

&lt;222&gt; 2628..2717

&lt;223&gt; exon4

&lt;221&gt; exon

&lt;222&gt; 7834..7924

&lt;223&gt; exon5

&lt;221&gt; exon

&lt;222&gt; 9804..9965

&lt;223&gt; exon6

&lt;221&gt; exon

&lt;222&gt; 11404..11527

&lt;223&gt; exon7

&lt;221&gt; exon

&lt;222&gt; 13539..14035

&lt;223&gt; exon8

&lt;221&gt; misc\_feature

&lt;222&gt; 13762..13764

&lt;223&gt; stop CDS

&lt;221&gt; polyA\_signal

&lt;222&gt; 13835..13839

&lt;223&gt; AATAAA potential

&lt;400&gt; 73

```

ttttttttt ttaattgtca aagtcattgat tctttttggt ttctctttta gatattgcta 60
tatggagatt tgccaaaaa taaagaaaat gtaatatatc tagcgaatca tcaaagcaca 120
ggtttgatt tcatttgatg aaatttgggt ttttctagaa atggtaaatg agcattaata 180
tgtacacaca catacacaca aacacacata tgtacacaca catatgtttt aaagacagga 240
tttcatgtga ccgagaatgg cctcactatc tctgagtagc tgagaatgat ttttaagcttg 300
tgacacacct gccttcactc ccaaggtaca ggaattgcag gtgctttctt tgnnnnnnnn 360
nntttttttt tttgagtttt ggggaggggg tatatttttt aatgtgtctg tagttggctt 420
tgttttaagc attttaatca tactttattt ttaaaaaaac taaaagcttt ttaaggcta 480
ggctctgcta tgtggcccta gtgttcctgg gacttgctct gtacaccggg ttgactctga 540
gcctgtgcgc cttctgcctc tgccctcata gttagattct caggacatgt tacaagact 600
gtgctgtgaa gatgagtttt tgttcctggg aggggaaggt ggagctgact tgtgaggtac 660
tgacttgggt ctgccttaca gttgactgga ttgttgcgga catgctggct gccagacagg 720
atgccctagg acatgtgcgc tacgtactga aagacaagtt aaaatggctt ccgctgtatg 780
ggttctactt tgctcaggta aactttgtct ttgccctttt atttcaaact taacaccatt 840
taatgaaact atatctgatt tttttgttta tgtgtttggt ttatggtacc cgtgattgaa 900
catggggtca tatgtgtgct actgagtgac agccttagtt cagacatttt ttaaagcgac 960
ttttactagt atttttattt agaattctat atgtgtgcac atgcatatgt gtgcttgtgt 1020
gcacacgtgg atgcatgtga ggtcgaagga caattttcag tacaagtgtg agtgtcactt 1080
tttaggcacc ttccactcct attttgagac agtctcctag acctttgctg agttgccag 1140
gctagccggc cagtgagccc tgggcatcta ccggtctctg cctccttacc tttacttagg 1200
ttacaagtgt gtgctgctac gccagctgt ttactagatt ctagggatcc aaatgtgggt 1260
cctcgtaact tgtgagacaa gtactttcca aactgagcca cctccctagc tcttcttcac 1320
ggttcctgat ggtgtgtgtc tagatggctg gttgtccgta tatttaagtc cagtagcaga 1380

```

aatacaaaata	cctaggagtc	caatagaaag	ctacaagtgc	agaattgaca	atcggtaatg	1440
ttcggaaatt	gattcaaaaag	tagttagtga	gtgacagaca	ggagctaaaa	gcagactctg	1500
agctcagagt	gtgaagtgtg	gagaaatgtg	ttttctcaca	gttctgaagg	ctgaaagtct	1560
ccccagggtc	aggatgtggg	tggtagtgc	gtctcccaan	cacccacctc	tttggattat	1620
agactgcagc	cttctccctg	tgttctgagc	cggcctttcc	cacatgtgga	catccttggt	1680
gggtgttcca	ccagcagggc	ctcagctagt	gcccttattt	cacttaactg	taatgatttt	1740
cttaaagacc	ctgtctccat	acacagtcac	tgtggaagct	gaagcttcaa	tgtaagagtt	1800
aaggggggag	ggggaaattt	agtccataat	ggtgtcacac	caatctctgt	agctgagtc	1860
atgattcagt	tctttaaagg	ctctgagtg	agacattatc	ttaattattt	tgccatttta	1920
tgtattatct	tttaatttatt	ttatgtaact	gaatgcctgt	gtatatatgt	ttctggttcc	1980
tagtccatat	tttaatttctt	ttaaaggatg	aggtgtagac	ttttgtcttt	tttaatttctt	2040
atccttccct	cctggcctcc	tgtggcctct	tttactgtatt	tattattttt	aatttatttt	2100
atgtgtttga	gtgtttttgta	tctatgcatt	cctggggccc	atggaggtca	gaaaaacaca	2160
ttaggtgggc	tacaactgag	ttatgggtgg	tttgtgacca	tggggtgctg	ggacttgatc	2220
accagtctct	gagaagactc	gtgtctgctg	agccttctct	ccagtccctg	gagtggtgat	2280
attttaagga	tactttttaa	tgacttggfg	aatgacagta	gaaaatcaaf	gagttaggat	2340
ccatcgga	aagcttttga	actaaatctt	ttaaagagaa	aataatttta	gtgctaacaa	2400
aattaaatgt	gtattttcca	tgatgcagtt	ttacttgggc	tctgtagaaa	taggattttc	2460
aggtacatat	tgtatatata	gttggaata	tttaataact	aactgtcgct	tgagtctga	2520
aatgtagttt	tatgtttttt	actcattagg	agtcagttg	ccttaataac	tacggagatt	2580
agttattaaa	gaataattgc	tcttcttttt	tctttctgt	gtaccagcat	ggaggaattt	2640
atgtaaaacg	aagtgcctaa	tttaatgata	aagaaatgag	aagcaagctg	cagagctatg	2700
tgaacgcagg	aacaccggta	agtgccccc	cttttattcc	tcaaggcagg	ttaagaagtt	2760
aagtctctaa	gtcattttga	aaatatatta	ccccatgtgg	agcaatggaa	ctgggtccgg	2820
gtttgtttga	gataagctgt	cctctggccg	tgaggttaaga	ttgctgcagg	tgattgttaag	2880
gtttctcctg	agtaacagtc	agcatgggct	cgggacgggc	aagggcaggc	cttagtgtgc	2940
agaggatgga	gctcactgaa	gccccaaaga	gttagtcttc	acatgagatt	cagttctaga	3000
agaagttaa	ttgctttctt	tctgtgtaaa	tttggatttt	tattgtagaa	attaaagttt	3060
gtttctttt	aaaacaaaca	caaaccaga	gcaaagagtc	tcttagtgaa	gagtcattcc	3120
gtgtcagtat	tttacacaac	tgtttttctg	ttaaaggggg	aaaagaattc	aaatcttctc	3180
tttcaagaat	gctgactgct	gccaactgcc	tctcccgtg	gccccctctc	gtatagacag	3240
gcatagctat	ggtagaggct	tgggcccgtc	ttgtctttct	cctctctctg	cttctctacc	3300
ctttctctcg	tgccctccac	ttaccaggcc	ctgggaagct	acacaccagg	caacagtgac	3360
cagggcctcg	gcctgggctt	cgaccaatta	ctagagcaga	aacagcagca	gctgcagtg	3420
tgttttgtgc	tgtgcactgt	attaggttgt	gttttcatca	cctttgggtt	ttgtgatgtt	3480
ttgatgaagt	cctggtagca	ttctagtgtt	tacattctgg	gtagatagag	tttattcaag	3540
gtctcaaggc	atatgaatgg	aagagctcct	ctttacagcc	attcgtgtag	catgcataac	3600
tgtcttctct	tattctctct	agtgctcttt	tttttgtgtg	tgaatctgat	gtcttgttat	3660
tcacctacaa	tgtggagtga	tgggtcataaa	catataaagt	acttatgcct	ttatctgcca	3720
aattgtattt	aacttttctag	cttttaatat	aactttttat	ataataatta	atttatttta	3780
aaaaaaattg	aataccagcc	tgtttatagt	gcataatgc	gtgttcttag	cactcaggag	3840
acaaaggcag	aagtgtgaga	acttcagact	catactcagc	tatatacaag	accccaaatt	3900
tgtgctagat	tctgcagtac	agccatgagt	gtcccatct	tagagggaga	tcgctcatcc	3960
ttgtgctgtt	ctttaagtct	tacctgcaa	cccactgtaa	gtacactctt	gctcacagtc	4020
ctttagaatc	tcacactctt	tctctttaca	gacaccatgt	cattgcccac	tttattattt	4080
atctgatgtc	tacaaagatt	atgaaagaga	aacttgtatg	cattctgtgt	aaagtacttg	4140
acacaaataa	tagtattcaa	gaatgacttc	ttaaatgaac	actgaatgaa	tagtttggtc	4200
taattttttt	gatcaacaaa	tcaaaaaata	tttagattaa	atatctaaga	tacaaagcat	4260
aataccacat	gaatcattaa	agtgagtaat	caatcttata	agtgactgac	cctaaaactc	4320
atagacatta	ataattgctt	tcatgtctta	gatataaact	ttattgatta	atacgttctc	4380
atgaaagtgg	ttcttgggaag	gttctggaaa	cgaaaatatt	tttcttactg	cttttttctt	4440
ctagtaactg	attgaatttt	tctgcagttc	cataaagcat	ctgggtcaatt	gctattatcc	4500
aatatgagga	tatataacaa	agtattgatt	tttaaatttg	gcggtgataa	gacaagactg	4560
ggcgtgtgaa	tgaggggggc	tctgtttctt	gtcccttctc	ttgggttctt	ttccttttgt	4620
tggtttgctt	tctccagctg	ctatgtgatg	ggttctgatt	atcttattat	atcttatttt	4680
gttatttttc	attgttatct	cttagaagcc	aacatgttat	atcacctcca	ctcccaccat	4740
taggtgttct	cacaaatacc	caaagctaaa	caaccacatc	atgtcatgtt	nctgtatact	4800
tccataagtg	ttgttaactc	tactgactct	tgtgagcagg	cccaattggc	tttatcccta	4860
gctgggtgac	ctgggttctt	ccccaacacc	ataccgtcca	tcaaactgag	tccttttcca	4920
agcacacacc	agatactgct	catctgagga	ctcttctcat	ccacctaaag	actgctgtct	4980
cctcggcaga	aagggcctct	agtcccatcc	ccttaagccc	tcaccaatgc	cttaggaaca	5040

tgtgctcaat	gcccctgtgg	gtcatttccg	tttacagtag	ggaaatttgc	ctgataactt	5100
gcagcacacc	tataaagagg	ccttgcttgc	tctcatat	agctggagaa	gataatgtac	5160
tcaccaactc	cactctatgc	aaccagtcct	gctctgccc	tgccagtcag	acgtgaatct	5220
tacacctgga	ttcagattga	tgaatctaca	acatcaccca	ctccatgctt	ccttctaata	5280
cagcagttct	agcctgaatg	acagatgcta	cccaagtcct	atctagttag	ccctgtccgg	5340
agtaaccctg	accttgagga	ttagaccagg	atgcacatcc	tgccaccagt	ccctttgtcc	5400
acctgacttc	atcccacccg	ggccatagcc	catgctcagg	ctccaccctc	catgcacaaa	5460
gctggctttt	ccagcttcct	tcacctgtat	cagacacaaa	tagcaaaagg	ggtccacgtg	5520
cctaggtccc	atcacaagac	catgtgcggt	agtttgga	acagtctcca	cttgaggctc	5580
agatagnttg	gaatcttggc	tctcatgtag	ttgtactgat	tagatcagtt	taggaagtat	5640
gacctttntg	gaaagaacat	ataactggga	ggggctttga	gatgtaaagg	ccccacacaa	5700
ttcctagttc	aaactntact	gctgtctcaa	agcttgaggc	atgaactctc	actgttccctg	5760
atgtcatggt	tcctgtctgc	ttccacaatt	ccctatcatt	aggggtccct	ttccttccctg	5820
gaattttaag	tataaataaa	cncctctttn	taaaacaaca	agaacaacaa	atctgacnct	5880
gataatggat	tttaaggcgt	cttctctgga	taagaaaaaa	aaaagaatat	atttgcatag	5940
gtgctgtatt	acttttgtca	ttggtataac	ctgactggaa	gcaacttaaa	ggaagaagaa	6000
tgtatcttga	ttttagatt	aagagcacca	tgactaagaa	ggcatagcag	cacaggtgca	6060
ccagcaagaa	cataggctgc	tagctcagat	ctctgtagat	atgggaacag	ggcaggaagc	6120
tagtagtcta	taaacctcag	gacccatccc	atggagtccc	ttgtcttcca	gtgatgtcct	6180
gtgtcttaaa	gtttcacagt	ccccacagca	gcacctgccg	tctgggaacc	aacctgtggt	6240
ggatatttta	caacgtgata	ggcatatttt	gtctctagcc	ctgtagggtt	atagccatcc	6300
tatacttcag	tttatctagt	ccacctcagt	ctgatggctc	tatagttcca	acacttcaaa	6360
actacaaagt	cttaagggcc	atgggctcgg	gtttattaga	gcagtaacac	ctctactagc	6420
tttctgtgtt	accactcct	cttaaggctc	ggttgaaatc	ctaataaggaa	gcagcttgag	6480
aggaggggtt	attgtggccc	atactttgtt	ggtacattct	atcatgcaag	ggtggcactg	6540
tgatacagcc	gaggccatcc	gaggatggta	ctgttggctt	acatctgggt	gggacaggaa	6600
atgggtaattc	tcaaggccca	cctgcttggt	gacttcttcc	agttaagccc	catactctaa	6660
atcctctaca	acctccaac	ataatgccac	cagctgggga	tcagctgttg	acagtgcctg	6720
cccaggggag	cagttaaaat	ccagaccagg	ggacctgaaa	acagagaact	gcagaggggc	6780
tgtgggactt	tataccagct	ttgcagacaa	atcacggcat	ttctttgtga	gcttggttca	6840
taaaacaaata	tatatctcc	tataggctcc	tttagtgggt	gtttcatatc	cacaaatttg	6900
ttcagaaaaa	cactgtgttt	tatgctagct	gtgtaggaga	taataccgct	gggagtcact	6960
tgagcatgga	taagtacat	agttcgtcct	catgagctcc	tgctctgttt	ctgtattatg	7020
tttacttgat	gagtttagtt	tgctcagttg	ccaccaatta	aaaagtatca	ttttattttt	7080
tttacaatac	tcagttctca	agttaggagt	tttgttatta	tatggcttca	atattcacat	7140
tttaaccttt	ccaggagtta	agtataaaaa	ctgttatcaa	ctgttgactt	agtaaatatc	7200
tattacagat	actatattct	tcttagttta	tatcatgaat	atgaggttgc	ttaaagtaag	7260
tgatgtaaaa	tacactaggg	gatgcttata	aaatggaaatg	ttgtgagttt	tttgaaacac	7320
gagtactaaa	ttcataagtt	tttaaatagt	tacactgtta	gcttcagtac	tgctagatac	7380
atgtctataa	tggtgaaga	gtggagcttg	gatattataa	gtgtactctg	tatattcatg	7440
cagacatata	gcagattcca	ctagtatgtg	tggttaatat	gtgctaataa	aaatttataa	7500
caaaagtcat	gtttttattc	tggaagccag	aggggttggt	tggtgctgatt	tttaagtcagt	7560
gactattagc	atatttctaag	aaacagtttt	naggattttta	aagattggct	ttaccataaa	7620
tgtagagcta	tgttttacta	taatccatat	tatggctcggc	cttaattcaa	tctctgcagt	7680
ttggttactc	tgctcaaaat	gaaggtcatt	tataaatgat	acacattttc	tcaccatagg	7740
aaatactacc	tgcccaataa	cagagttaga	attgctaaat	tgatggtacc	aacaatggac	7800
tcaacacaaa	ctaaagttta	tttatgccca	cagatgtatc	ttgtgatttt	cccagaggga	7860
acaagggtata	atgcaacata	cacaaaactc	ctttcagcca	gtcaggcatt	tgctgctcag	7920
cggggtaagt	aaagatttta	ctgtatttcag	aaaaacactt	ttttaagaag	agtgatcttt	7980
gtttccttca	gagtcatact	aaagaatatg	cgtttcttgt	aagagctaag	tgagagaata	8040
tccgatcttc	tacagagtta	ggtatattct	tattagtctg	tgtctgagag	gttagagacg	8100
caggcttgct	atggcacatt	tcccatgctg	tgaattgagt	taaaaatgta	ggtaaatgat	8160
atccccaaaga	aagtatactt	ttnggagtga	ctcagataaa	agcctggtgt	tataacataa	8220
acacgcacgt	gcggatgtat	atgtagcaca	tatgtaaaca	caggatatatg	canttgtaat	8280
aagaaagtgg	aggtcggggc	ccactgcagg	caagtctttt	agtgatgctg	agctaattgct	8340
gagaggtaga	aagaccaaga	aggctggagt	tgctcattcg	gcaaagggtca	gagctcactg	8400
tgtgccataa	ctcgagtgtt	ctgtctccct	tttgatacag	ttttcttgtt	tttaattatt	8460
agtttttata	attatcccat	aaaatgtggg	ctcattgttg	tcatogtttt	cataaagtc	8520
ttcaagtata	caccagcaa	gtatctaaat	acactgggaa	gaatcagtca	gctgatggct	8580
tgaagtttca	ggacatctag	tgccacatca	tgcttcagaa	ccgacctgca	cttagtcagg	8640
gtcatattca	tgccacgtga	agacgagagg	aggccatgcc	gtctgactta	ggatggaaat	8700



ttccttcgag	caaacacgaa	cgggctaggt	cttagttata	ggcatagtgt	ctgtgggttat	8760
actaggcaga	cattagtggg	ctgggtgtta	gaaggtacag	acaggcaaga	atttgctgta	8820
gatttgtttc	cctcatgtgt	tgacaccaca	tctaacctgc	tttttgagct	tctagtccca	8880
ataatctcat	aaaaatactg	gttgaccag	aaatgggtgt	gcaaagctat	gatcccagct	8940
cctgggatct	aggggtgggag	gatcataaat	ttgaggccag	cttgggtctg	tcttagagaa	9000
aaaagaaaaa	taaaaagtct	ggtcaaggta	acatggagcc	tggaagtttc	acagggtgat	9060
tctgtaaagg	tcttgagaca	agatggcctc	tagtggcgaa	tgacttagct	gacaangaaa	9120
actttcccag	cttgggtgac	ttttcagact	tcatacaagt	ttgtgaataa	attacactcc	9180
ttctgccctt	gggactgaac	tcagatatgt	ggttgtggga	atggccttct	ttcccacacc	9240
accctgcatt	ttaaaaatct	ttctgtagac	agtcaccaca	tcctgtagct	gttcttccct	9300
atgtcgccac	tttccctgga	gagaggcagt	gcagacttca	acccgcttct	ccctagtcgc	9360
tgttcatagc	acatcgaaag	acctagtgtc	tcctgtgaaa	ttgtaagtac	atcctggagt	9420
ccaggagagg	aggaagccga	acagagtggg	gggaatgctg	agttctgtcc	taagaaagac	9480
tgctgtctta	gcaagatgct	gctgctctcc	tgctgtgtct	ttctgttcag	aactttcaa	9540
agagaaggct	cgcagtgggt	cataatcttc	ccaaggacca	gccttcccag	cttctcgag	9600
catatctcat	tcatgtagat	gtttaatgga	tatgtgtcaa	tggtgtgac	ctaagtgaga	9660
tggaatgta	tgtagcatt	ctaggtgtga	ggttatggca	ttaaacttta	atttccgtct	9720
atttggtgta	gttgataagt	aatttagatg	ttgactttca	tgtattccta	attatgacca	9780
cattgaatct	acctgctttc	taggccttgc	agtattaaaa	cagtgactga	caccaagaat	9840
aaaggccact	cagttgtgct	ttgattctat	gaagatcat	ttagatgcaa	tttatgtgt	9900
cacagtgggt	tatgaaggga	atgagaaagg	ttcaggaaaa	tactcaaatc	caccatccat	9960
gactggtaag	tcctgtattc	catagaagct	gaatagtaca	tggtacaggt	aagataaact	10020
cttggttggt	cgctttgctt	agcttggttc	agtttggttt	tcagtagagg	gttccactat	10080
gaagctctgg	ctggccggga	actcactatg	tagaccaagc	tggtcctggg	ctccactaca	10140
cccagcacca	atcaccact	cttatncttt	tatgcttttt	tggttttgct	ttgagctttc	10200
tttataacat	gtttgggaag	gacattgtca	ttattttcaa	gaagaaatat	ggtcttttcc	10260
caacatgcta	gaattttaag	actcagaact	cttgcctttg	tcagtgacaa	agtgagaatg	10320
gctgtgaagt	gacgtggctt	tgagtggaga	tagttcaggt	aactatagcc	acagactcaa	10380
catttgaaca	tggaacagg	tgagaacgga	gtgatggaag	attctggccc	ctttcagaga	10440
attcatttta	gagagagatg	agagtagtaa	ggaagagaga	agagagagac	gtggatattt	10500
gctgcagact	aaagagatct	cttataatcg	cagtactaag	gaggaagaag	cagaagatga	10560
tgactacagg	gcagggtga	acaatctagt	aaaatcctaa	gtcaggaagt	cagggctgag	10620
gtgcagctca	gtaggagagt	tggtgtctgc	cctacacaag	gcctggattt	agctcccagt	10680
agcaacgaag	ggaggcgagg	gtgggcaaaa	tcgaacactt	actcctggag	actcccttta	10740
tgaatattac	cacactccag	taaatactct	ccagagattt	cagatgagat	tctgcttctt	10800
ggtaaacagg	aggccaagaa	tattatgtca	cactgaacat	gggatggaag	acatgttctg	10860
aggaatgtct	gcactccagt	gtgatgaaga	cttgaagttt	agggacattt	tccttccctg	10920
gccccactca	ccccatctgt	attgagtatt	cccctagtgc	tcacttttat	ttgtatgtta	10980
actttcagga	aggggaagca	gattgatatt	caaaccacgc	cagttttctt	aaatactttg	11040
tggtatggat	tggttttgac	agtaaatgag	gaaatgtaaa	atgtaaaaga	ttctaatttt	11100
taatatttta	aaggtgaggt	tttctgttag	tacgcagagt	gagagggttc	ttactgatgt	11160
ctgcttacct	agaggaagga	tggtacttcc	tccaaggctt	gctgttagaa	gtcagtgaca	11220
tggtcttaac	aaagatatg	tgctaattgag	gttttaattt	cagcttaata	ctgcaaatca	11280
taagtgcata	gctttattgt	tttaaattct	tttagtctta	atgtttcatt	tttaccataa	11340
gttactttgt	ataatcacaa	attctaaact	agtaagacgt	gaaattttct	tcttctttgt	11400
tagagtttct	ctgcaaacag	tgcccaaaac	ttcatattca	ctttgatcgt	atagacagaa	11460
atgaagttcc	agaggaacaa	gaacacatga	aaaagtggct	tcattgagcgc	tttgagataa	11520
aagataggta	agtggtaaga	gctccagcat	ttagaaagtg	cagttcaacc	aaattttact	11580
ctcagatcct	gcttgaaagg	agtcttttta	tcttcattat	ttagtaaata	ctaatcatat	11640
ctgcatagac	aagaccacat	atacttaaat	gtagcatgtt	tcattggtgcg	ttacccttgt	11700
ttaacaatta	agttaaacat	cctacatcag	tttgctgtgt	gattttctgta	ccatgacaac	11760
tcaacacagc	gatgcgttta	ttccaaagtc	gatagcacag	caaaagtga	actaaagtct	11820
gtattgtttc	aagaaatgctt	cttgtaaac	cgggttaaat	cttattctat	cctttctgtg	11880
tcacattgta	cattttcatg	agtcactata	aaaatcatga	catggtggcc	tacctgcagt	11940
gtttgctgga	cagtaggctg	ctgtgtgata	agagcctttc	ctcttcagct	acacggggga	12000
cacgaggctt	tggtgttcaa	gactgaagca	cgggtgagca	caacaccttt	gtgtgtgtgg	12060
aaggaaggga	attgttcttt	tcataatgaa	attgtccctt	ttcttgagtt	agtagaaagt	12120
attacaagga	tagagagttg	aaatgaagct	tttatattaga	tttatgcctt	gtgtgtgtcac	12180
gtgtttctac	ctgacataac	ttttcaaccc	agccgctcag	gattattttg	atgatgggaa	12240
caatgtaaga	aggcctatgt	atcggttaact	cactgtgtga	gctctgtgga	ancgntcnn	12300
caggcagtag	ggacgcttct	gtgcttttgt	gcctgtcctg	ctgttagaat	cttacagagg	12360

```

aggatgaatg aatgaccctt tttattttctc ttgtctgctt ttctaatttt atgggaataa 12420
gaacttttgg taggtctctg tcaactggcct cttgttgtga agagacacct tgagcaaagc 12480
aactcttctg agagaaagca tttagttggg gaattcctta cagcttcaga ggttgagttc 12540
gttttcatca tgctgaggac agggaggcac tcaggcagga gaagtagttg agagccacat 12600
tctgatctac aggcagagag agacagactg agcctggcat gggttcttgg aacctcaaag 12660
cctctcatcc ctaccccatc tcccgacccc tatacacact tcctccaaca aggctacacc 12720
ttctaactct tcttaaagag tcaccacatc cagcgactaa gcattcagat atgtgaacct 12780
gtcggagcct ttcttactca gatcacctca ggaggaaaac tcctatgcta taagaatttc 12840
ttttctttcg catctttgaa agcttggttt tgtgtgatta gatcctggcc tcacacatgc 12900
tcggcaatca ttttactggt gagctccagc ctcagccgtt ttcattggct tatgggatgc 12960
gagccatggg agagaagcta gaaggccttt cgttttatga gtcgggttgg tggaaccact 13020
tacagatgga agatttataa acaaaaatga agctggggcc atcaaggctc agcactcgct 13080
gctcttccag agagtccagg ttcattttctc agtaaccaca tgggtggcttt gtaaagttaa 13140
cttcataattc aatgaccctg acacctctt ctggcctctg tgggcaccag acacaatcat 13200
gggtatacaga cacacacact agccaacacc catctacata aaagtatata anacatatct 13260
ttatcttaaa aatccccgaa gtcctcatta aatatcttag atccccgccg tgttttgatt 13320
tttgtttccc acgtggtgag gatataatat catgtccaaa ctgtaaggag tgaatgccct 13380
cccgtgcctc tcggacacct ctgcactcat ccaagttttc taaggagctg tacttgcctc 13440
gcaagtactc aatacctaataaatgggtttt tgtttgtttc aacaccaaaa atgtccaaaa 13500
ctgaaagatc cagattctgtt ttttcttctc ggccataggt tgctcataga gttctatgat 13560
tcaccagatc cagaaagaag aaacaaatct cctgggaaaa gtgttcattc cagactaagt 13620
gtgaagaaga ctttaccttc agtgttgatc ttggggagtt tgactgcggt catgctgatg 13680
acggagtccg gaaggaaact gtacatgggc acctgggtgt atggaacctc ccttggtctg 13740
ctgtgggtttg ttattaaagc ataagcaagt agcaggctgc agtcacagtc tcttattgat 13800
ggctacacat tgtatcacat tgtttcctga attaaataag gagttttctt gttgttgttt 13860
tttttgtttt gttttgttct gttttaagcc ttgatgattg aacactggat aaagtagagt 13920
ttgtgaccac agccaacatg catttgattt ggggcaaca catgtggctt ttcagggtgt 13980
ggggttgctg gagacatgga agctaagtgg agtttatgct gntttttttt tttttttnaa 14040
tgttttcatg aattaatgtc cacttgtaaa gattattgga tactttctgt aattcagaag 14100
gttgattttt aacactagtt tgcagtatgt ttcgctatat tggttatctt ccatttgact 14160
acttggcagc tcagactctt aataactaaag tattttacat tttgaagcta tgtgatactg 14220
gttttttgtt gttgttgttg ttgttaattt ctgaaagtca atgaaagaca ctgtaatgat 14280
gcgttaagat gttccaagaa aaagggtgaga attattcatg gcaaaaaaga tctgtctagt 14340
gtatattttt attatattgc tctatttagc taattttctt tatatttgca aaataatgaa 14400
catttttaat atttattaaa atgcttgatt tgcatacccc cgattctaca gagaataatg 14460
tgtaaaagtgt cagaatagac ttgaagctct ctgtgactc agtctcctt gtcagagctt 14520
ctagtagccc agctactgag ctgctttgtt agtacctcca gcacctgagc cgtaagtac 14580
ttataaatgc aagggacccg ttatcttcat atcggaatag acatgaacag agctctaagg 14640
cgatgaaagt ctgccagcat cctctctgtc ctgcacgtg ccttctgcct ggctccattt 14700
gctttggcac tgcgttcgat ctagagtgtg ggtgctcact gcttatttca gccctggctc 14760
tgtgtgtttg tgtcctccag tgggtgctgt cactgttggg gtgcagggtg tgcctgccctg 14820
actcagaggg gcagctccct ggctcctgag ggtgagcctt cttggctact acagaagtat 14880
tgtgctttg tgtatggcaa gaaccatcag gattggataa atgtgttatt tctctttgat 14940
ttccatggag ccacactgtt ggtacatgtc ccctgtgaac agagctacct ttcaggagca 15000
catcatactg tcgtgagtca cggcacgggt gtgcctgtga gaagaggctt tctaactgtg 15060
gatttgccgt gtttctatgt tgtgatttaa gcgtgattgc ctactagtca tccaaggtaa 15120
catttctgca aatttcatac agatttttgt cacaaaatta ctataccaat gatctagttg 15180
aaatagacca attgaatcac aataaataat tttttttaat tgagggaaaa tttgcttctt 15240
gttttttcaa agccagaaaa cgagccattt caaacatctt tgaagagtca tgtgctgtca 15300
cttggttttct atgtgttagt gtctatatct atgtatggat acacatgaac atgtatatct 15360
atacacacac gccaatagaa tataacagcc taaaaacaat ccagcttgtg tatcatgtta 15420
ctgtgctgaa ttgtaatggt ttttacttac aaagtggagg taaaaatgat ttcattgtct 15480
tggttaaatc gtttttttca gcaatcctat tagagcttat tttgaccaga tcaaaataag 15540
tacaagttca gagactttaa atatggctga ggtctagagc gatagctcag tagttaggaa 15600
cacatgccac tcttcaagg gcttcagttc ccagcactca tatggaggct cacagaaggc 15660
tggaattcca gcttcatgga attggacaca tctcttagct tccatggatc tgtctgtctg 15720
tctctccctt ctctctctct ctctctctct ctctctctct ctctct 15760

```

<210> 74  
 <211> 354  
 <212> PRT  
 <213> Mus musculus

<400> 74  
 Met Arg Tyr Leu Leu Pro Ser Val Leu Leu Leu Gly Ser Ala Pro Thr  
 1 5 10 15  
 Tyr Leu Leu Ala Trp Thr Leu Trp Arg Val Leu Ser Ala Leu Met Pro  
 20 25 30  
 Ala Arg Leu Tyr Gln Arg Val Asp Asp Arg Leu Tyr Cys Val Tyr Gln  
 35 40 45  
 Asn Met Val Leu Phe Phe Phe Glu Asn Tyr Thr Gly Val Gln Ile Leu  
 50 55 60  
 Leu Tyr Gly Asp Leu Pro Lys Asn Lys Glu Asn Val Ile Tyr Leu Ala  
 65 70 75 80  
 Asn His Gln Ser Thr Val Asp Trp Ile Val Ala Asp Met Leu Ala Ala  
 85 90 95  
 Arg Gln Asp Ala Leu Gly His Val Arg Tyr Val Leu Lys Asp Lys Leu  
 100 105 110  
 Lys Trp Leu Pro Leu Tyr Gly Phe Tyr Phe Ala Gln His Gly Gly Ile  
 115 120 125  
 Tyr Val Lys Arg Ser Ala Lys Phe Asn Asp Lys Glu Met Arg Ser Lys  
 130 135 140  
 Leu Gln Ser Tyr Val Asn Ala Gly Thr Pro Met Tyr Leu Val Ile Phe  
 145 150 155 160  
 Pro Glu Gly Thr Arg Tyr Asn Ala Thr Tyr Thr Lys Leu Leu Ser Ala  
 165 170 175  
 Ser Gln Ala Phe Ala Ala Gln Arg Gly Leu Ala Val Leu Lys His Val  
 180 185 190  
 Leu Thr Pro Arg Ile Lys Ala Thr His Val Ala Phe Asp Ser Met Lys  
 195 200 205  
 Ser His Leu Asp Ala Ile Tyr Asp Val Thr Val Val Tyr Glu Gly Asn  
 210 215 220  
 Glu Lys Gly Ser Gly Lys Tyr Ser Asn Pro Pro Ser Met Thr Glu Phe  
 225 230 235 240  
 Leu Cys Lys Gln Cys Pro Lys Leu His Ile His Phe Asp Arg Ile Asp  
 245 250 255  
 Arg Asn Glu Val Pro Glu Glu Gln Glu His Met Lys Lys Trp Leu His  
 260 265 270  
 Glu Arg Phe Glu Ile Lys Asp Arg Leu Leu Ile Glu Phe Tyr Asp Ser  
 275 280 285  
 Pro Asp Pro Glu Arg Arg Asn Lys Phe Pro Gly Lys Ser Val His Ser  
 290 295 300  
 Arg Leu Ser Val Lys Lys Thr Leu Pro Ser Val Leu Ile Leu Gly Ser  
 305 310 315 320  
 Leu Thr Ala Val Met Leu Met Thr Glu Ser Gly Arg Lys Leu Tyr Met  
 325 330 335  
 Gly Thr Trp Leu Tyr Gly Thr Leu Leu Gly Cys Leu Trp Phe Val Ile  
 340 345 350

Lys Ala

<210> 75

<211> 22

<212> DNA

<213> Mus Musculus

<220>

<221> misc\_binding

<222> 1..22

<223> amplification oligonucleotide g34292.pu

<400> 75

attaaaacac gtactgacac ca

<210> 76

<211> 19

<212> DNA

<213> Mus Musculus

<220>

<221> misc\_binding  
<222> 1..19  
<223> amplification oligonucleotide g34292.rp  
<400> 76  
agtcatggat ggtggattt 19  
<210> 77  
<211> 26  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..26  
<223> amplification oligonucleotide BOXIed  
<400> 77  
aatcatcaaa gcacagttga ctggat 26  
<210> 78  
<211> 33  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..33  
<223> amplification oligonucleotide BOXIIer  
<400> 78  
ataaaccacc gtaacatcat aaattgcatc taa 33  
<210> 79  
<211> 22  
<212> DNA  
<213> Mus Musculus  
<220>  
<221> misc\_binding  
<222> 1..22  
<223> sequencing oligonucleotide moPGrace3S473  
<400> 79  
gagataaaaag ataggttgct ca 22  
<210> 80  
<211> 19  
<212> DNA  
<213> Mus Musculus  
<220>  
<221> misc\_binding  
<222> 1..19  
<223> sequencing oligonucleotide moPGrace3S526  
<400> 80  
aagaaacaaa tttcctggg 19  
<210> 81  
<211> 18  
<212> DNA  
<213> Mus Musculus  
<220>  
<221> misc\_binding  
<222> 1..18  
<223> sequencing oligonucleotide moPGrace3S597  
<400> 81  
tcttggggag tttgactg 18  
<210> 82  
<211> 18  
<212> DNA  
<213> Mus Musculus  
<220>  
<221> misc\_binding

```

<222> 1..18
<223> sequencing oligonucleotide moPGrace5R323
<400> 82
gaccccggtg tagttctc
<210> 83
<211> 17
<212> DNA
<213> Mus Musculus
<220>
<221> misc_binding
<222> 1..17
<223> sequencing oligonucleotide moPGrace5R372
<400> 83
cagtaaagcc ggtagtc
<210> 84
<211> 17
<212> DNA
<213> Mus Musculus
<220>
<221> misc_binding
<222> 1..17
<223> sequencing oligonucleotide moPGrace5R444
<400> 84
caggccagca ggtaggt
<210> 85
<211> 19
<212> DNA
<213> Mus Musculus
<220>
<221> misc_binding
<222> 1..19
<223> sequencing oligonucleotide moPGrace5R492
<400> 85
agcaggtagc gcatagagt
<210> 86
<211> 27
<212> DNA
<213> Mus Musculus
<220>
<221> misc_binding
<222> 1..27
<223> amplification oligonucleotide moPG13LR2
<400> 86
ggaaacaatg tgatacaatg tgtagcc
<210> 87
<211> 18
<212> DNA
<213> Mus Musculus
<220>
<221> misc_binding
<222> 1..18
<223> amplification oligonucleotide moPG15
<400> 87
tggcgagccg agaggatg
<210> 88
<211> 36
<212> DNA
<213> Mus Musculus
<220>
<221> misc_binding
<222> 1..36

```

18

17

17

19

27

18

<223> amplification oligonucleotide moPG15Bam1  
 <400> 88  
 cgtggatccg gaaacaatgt gatacaatgt gtagcc 36  
 <210> 89  
 <211> 27  
 <212> DNA  
 <213> Mus Musculus  
 <220>  
 <221> misc\_binding  
 <222> 1..27  
 <223> amplification oligonucleotide moPG15Eco1  
 <400> 89  
 cgtgaattct ggcgagccga gaggatg 27  
 <210> 90  
 <211> 20  
 <212> DNA  
 <213> Mus Musculus  
 <220>  
 <221> misc\_binding  
 <222> 1..20  
 <223> amplification oligonucleotide moPG1RACE3.18  
 <400> 90  
 ctgccagaca ggatgcccta 20  
 <210> 91  
 <211> 23  
 <212> DNA  
 <213> Mus Musculus  
 <220>  
 <221> misc\_binding  
 <222> 1..23  
 <223> amplification oligonucleotide moPG1RACE3.63  
 <400> 91  
 acaagttaaa atggcttccg ctg 23  
 <210> 92  
 <211> 18  
 <212> DNA  
 <213> Mus Musculus  
 <220>  
 <221> misc\_binding  
 <222> 1..18  
 <223> sequencing oligonucleotide moPG1RACE3R94  
 <400> 92  
 caaatgcatg ttggctgt 18  
 <210> 93  
 <211> 20  
 <212> DNA  
 <213> Mus Musculus  
 <220>  
 <221> misc\_binding  
 <222> 1..20  
 <223> amplification oligonucleotide moPG1RACE5.276  
 <400> 93  
 gcaaatgcct gactggctga 20  
 <210> 94  
 <211> 22  
 <212> DNA  
 <213> Mus Musculus  
 <220>  
 <221> misc\_binding  
 <222> 1..22  
 <223> amplification oligonucleotide moPG1RACE5.350

<400> 94  
 aatcaaaaagc aacgtgagtg gc 22  
 <210> 95  
 <211> 20  
 <212> DNA  
 <213> Mus Musculus  
 <220>  
 <221> misc\_binding  
 <222> 1..20  
 <223> amplification oligonucleotide moPG3RACE2  
 <400> 95  
 tgggcacctg gttgtatgga 20  
 <210> 96  
 <211> 20  
 <212> DNA  
 <213> Mus Musculus  
 <220>  
 <221> misc\_binding  
 <222> 1..20  
 <223> amplification oligonucleotide moPG3RACE2n  
 <400> 96  
 tccttggctg cctgtgggtt 20  
 <210> 97  
 <211> 21  
 <212> DNA  
 <213> Mus Musculus  
 <220>  
 <221> misc\_binding  
 <222> 1..21  
 <223> sequencing oligonucleotide moPG3RACES20  
 <400> 97  
 gatggctaca cattgtatca c 21  
 <210> 98  
 <211> 24  
 <212> DNA  
 <213> Mus Musculus  
 <220>  
 <221> misc\_binding  
 <222> 1..24  
 <223> sequencing oligonucleotide moPG3RACES5  
 <400> 98  
 tcctgaatta aataaggagt ttcc 24  
 <210> 99  
 <211> 24  
 <212> DNA  
 <213> Mus Musculus  
 <220>  
 <221> misc\_binding  
 <222> 1..24  
 <223> sequencing oligonucleotide moPG3RACES90  
 <400> 99  
 gtttggtatt aaagcataag caag 24  
 <210> 100  
 <211> 216  
 <212> DNA  
 <213> Homo sapiens  
 <400> 100  
 ctgctgtccc tgggtgtcca cacgtactcc atgcgctacc tgctgtcccag cgtcgtgctc 60  
 ctgggcacgg cgccaccta cgtgttggcc tgggggtct ggcggctgct ctccgccttc 120  
 ctgcccggcc gcttctacca agcgtggac gaccggctgt actgcgtcta ccagagcatg 180  
 gtgctcttct tcttcgagaa ttacaccggg gtcacag 216

```

<210> 101
<211> 70
<212> DNA
<213> Homo sapiens
<400> 101
atattgctat atggagattt gccaaaaaat aaagaaaata taatatattt agcaaatcat    60
caaagcacag                                                                    70
<210> 102
<211> 116
<212> DNA
<213> Homo sapiens
<400> 102
ttgactggat tgttgctgac atcttggcca tcaggcagaa tgcgctagga catgtgctgt    60
acgtgctgaa agaagggtta aaatggctgc cattgtatgg gtgttacttt gctcag      116
<210> 103
<211> 90
<212> DNA
<213> Homo sapiens
<400> 103
catggaggaa tctatgtaaa gcgcagtgcc aaatttaacg agaaagagat gcgaaacaag    60
ttgcagagct acgtggacgc aggaactcca                                          90
<210> 104
<211> 91
<212> DNA
<213> Homo sapiens
<400> 104
atgtatcttg tgatttttcc agaaggtaca aggtataatc cagagcaaac aaaagtcctt    60
tcagctagtc aggcatttgc tgccaacgt g                                          91
<210> 105
<211> 159
<212> DNA
<213> Homo sapiens
<400> 105
gccttgacgt attaaaacat gtgctaacac cacgaataaa ggcaactcac gttgcttttg    60
attgcatgaa gaattattta gatgcaattt atgatgttac ggtggtttat gaagggaaag    120
acgatggagg gcagcgaaga gagtcaccga ccatgacgg                                159
<210> 106
<211> 124
<212> DNA
<213> Homo sapiens
<400> 106
aatttctctg caaagaatgt ccaaaaattc atattcacat tgatcgtatc gacaaaaaag    60
atgtcccaga agpacaagaa catatgagaa gatggctgca tgaacgtttc gaaatcaaag    120
ataa                                                                    124
<210> 107
<211> 4342
<212> DNA
<213> Homo sapiens
<220>
<221> polyA_signal
<222> 325..330
<223> AATAAA potential
<221> polyA_signal
<222> 694..699
<223> AATAAA potential
<221> polyA_signal
<222> 828..833
<223> AATAAA potential
<221> polyA_signal
<222> 1821..1826
<223> AATAAA potential

```



```

<221> polyA_signal
<222> 2480..2485
<223> AATAAA potential
<221> polyA_signal
<222> 2800..2805
<223> AATAAA potential
<221> polyA_signal
<222> 4264..4269
<223> AATAAA potential
<221> polyA_signal
<222> 4320..4315
<223> AATAAA
<400> 107
gatgcttata gaattttatg agtcaccaga tccagaaaga agaaaaagat ttcttgggaa 60
aagtgttaat tccaaattaa gtatcaagaa gactttacca tcaatgttga tcttaagtgg 120
tttgactgca ggcatgctta tgaccgatgc tggaaggaag ctgtatgtga acacctggat 180
atatggaacc ctacttggct gcctgtgggt tactattaaa gcatagacaa gtagctgtct 240
ccagacagtg ggatgtgcta cattgtctat ttttggcggc tgcacatgac atcaaattgt 300
ttcctgaatt tattaaggag tgtaaataaa gccttgttga ttgaagattg gataatagaa 360
tttgtgacga aagctgatat gcaatggtct tgggcaaaaca tacctggttg tacaacttta 420
gcatcggggc tgctggaagg gtaaaagcta aatggagttt ctctgtctct gtccatttcc 480
tatgaactaa tgacaacttg agaaggctgg gaggattgtg tattttgcaa gtcagatggc 540
tgcatTTTTT agcattaatt tgcagcgtat ttcacttttt ctgttatttt caatttatta 600
caacttgaca gtcceaagct cttattacta aagtatttag tatcttgca gtagttaata 660
tttcatcttt tgcttatttc tacaagtcag tgaaataaat tgtatttagg aagtgtcagg 720
atgttcaaag gaaagggtaa aaagtgttca tggggaaaaa gctctgttta gcacatgatt 780
ttattgtatt gcgttatttag ctgattttac tcatTTTata tttgcaaaat aaatttctaa 840
tatttattga aattgcttaa tttgcacacc ctgtacacac agaaaatggt ataaaatatg 900
agaacgaagt ttaaaattgt gactctgatt cattatagca gaactttaaa tttcccagct 960
ttttgaagat ttaagctacg ctattagtag ttcctttgt ctgtgccata agtgcttgaa 1020
aacgttaagg ttttctgttt tgttttgttt ttttaataac aaaagagtcg gtgtgaacct 1080
tggttggacc cpaagttcac aagattttta aggtgatgag agcctgcaga cattctgcct 1140
agatttacta gcgtgtgcct tttgcctgct tctctttgat ttcacagaat attcattcag 1200
aagtcgcgtt tctgtagtgt ggtggattcc cactgggctc tggctcctcc ctgggatccc 1260
gtcagtggtg ctgctcagcg gcttgcacgt agacttgcta ggaagaaatg cagagccagc 1320
ctgtgctgcc cactttcaga gttgaactct ttaagccctt gtgagtgagg ttcaccagct 1380
actgcagagg cattttgcat ttgtctgtgt caagaagttc accttctcaa gccagtgaag 1440
tacagactta attcgtcatg actgaacgaa tttgtttatt tcccattagg tttagtggag 1500
ctacacatta atatgtatcg ccttagagca agagctgtgt tccaggaacc agatcacgat 1560
tttttagccat ggaacaatat atcccattgg agaagacctt tcagtggtgaa ctgttctatt 1620
tttctgttat aatttaaac tcgatttccct catagtccct taagttgaca tttctgtcta 1680
ctgtactagg atttttgctg cagaaatata tcagtggtccc acattaaaca taccagttgg 1740
atcatgataa gcaaaatgaa agaaataatg attaagggaa aattaagtga ctgtgttaca 1800
ctgcttctcc catgccagag aataaaactct tccaagcatc atctttgaag agtcgtgtgg 1860
tgtgaattgg tttgtgtaca ttagaatgta tgcacacatc catggacact caggatatag 1920
ttggcctaata aatcggggca tgggtaaaac ttatgaaaat ttctcatgct tgaattgtaa 1980
ttttctctta cctgtaaaagt aaaatttaga tcaattccat gtctttgtta agtacaggga 2040
tttaatatat tttgaatata atgggtatgt tctaaatttg aactttgaga ggcaatactg 2100
ttggaattat gtggattcta actcatttta acaaggtagc ctgacctgca taagatcact 2160
tgaatgttag gtttcataga actatactaa tcttctcaca aaagggtctat aaaatacagt 2220
cgttgaaaaa aattttgtat caaaatggtt ggaaaattag aagcttctcc ttaacctgta 2280
ttgatactga cttgaattat tttctaaaaa taagagccgt atacctacct gtaagtcttt 2340
tcacatatca tttaaacttt tgtttgtatt attactgatt tacagcttag ttattaattt 2400
ttctttataa gaatgccgtc gatgtgcatg cttttatggt tttcagaaaa ggggtgtgtt 2460
ggatgaaagt aaaaaaaaaa ataaaatctt tcaactgtct taatggctgt gctgtttaac 2520
attttttgac cctaaaattc accaacagtc tcccagtaca taaaataggc ttaatgactg 2580
gcctgtcatt cttcacataa tttttcccta agctttgagc aaagttttaa aaaaatacac 2640
taaaataatc aaaactgtta agcagtatat tagtttggtt atataaatc atctgcaatt 2700
tataagatgc atggccgatg ttaatttgct tggcaattct gtaatcatta agtgatctca 2760
gtgaaacatg tcaaatgcct taaattaaact aagttggtga ataaaagtgc cgaactggct 2820
aactcttaca ccatacatat tgatagtttt tcatatgttt catttccatg tgatttttaa 2880

```

```

aatttagagt ggcaacaatt ttgcttaata tgggttacat aagctttatt ttttcctttg 2940
ttcataatta tatttctttga ataggtctgt gtcaatcaag tgatctaact agactgatca 3000
tagatagaag gaaataaggc caagttcaag accagcctgg gcaacatac gagaacctgt 3060
ctacaaaaaa attaaaaaaa attagccagg catggtggcg tacactgagt agtttgtccc 3120
agctactcgg gaggggtgagg tgggaggatc gcttcagccc aggaggttga gattgcagtg 3180
agccatggac ataccactgc actacagcct aggtaacagc acgagacccc aactcttaga 3240
aaatgaaaag gaaatataga aatataaaat ttgcttatta tagacacaca gtaactccca 3300
gatatgtacc acaaaaaatg tgaaaagaga gagaatgtc taccaaagca gtattttgtg 3360
tgtataattg caagcgcata gtaaaataat ttttaacctta atttgttttt agtagtgttt 3420
agattgaaga ttgagtgaat tattttcttg gcagatattc cgtatctggg ggaaagctac 3480
aatgcaatgt cgtttagtgg ttgcatggct tgctttataa acaagatttt ttctccctcc 3540
ttttgggcca gttttcatta cgagtaactc acactttttg attaaagAAC ttgaaattac 3600
gttatcactt agtataattg acattatata gagactatgt aacatgcaat cattagaatc 3660
aaaattagta ctttgggtcaa aatatttaca acattcacat acttggtcaa tattcatgta 3720
attaactgaa tttaaaacct tcaactatta tgaagtgtc gtctgtacaa tcgctaattt 3780
actcagttta gagtagctac aactcttcga tactatcatc aatatttgac atcttttcca 3840
atttggtgat gaaaagtaaa tctattcctg tagcaactgg ggagtcatat atgaggtcaa 3900
agacatatata cttgttatta taatatgtat actataataa tagctgggta tcctgagcag 3960
gggaaaagggt tatttttagg aaaaccactt caaatagaaa gctgaagtac ttctaataa 4020
ctgaggggaag tataatatgt ggaacaaact ctcaacaaaa tgtttattga tggtagtgaa 4080
acagatcagt ttttccatcc ggattattat tgggttcata ttttatatgt gaatatgtaa 4140
gatatgttct gcaattttat aaatgttcat gtcttttttt aaaaaagggt ctattgaaat 4200
tctgtgtctc cagcaggcaa gaatacttga ctaactcttt ttgtctcttt atgggtattt 4260
cagaataaag tctgacttgt gtttttgaga ttattgggtc ctcattaatt cagcaataaa 4320
ggaaaatatg catctcaaaa at 4342
<210> 108
<211> 62
<212> DNA
<213> Homo sapiens
<400> 108
agattggatt cgttagattaa acttgagaaa caaaccataa aagtgggaagg ccctctttaa 60
ca 62
<210> 109
<211> 86
<212> DNA
<213> Homo sapiens
<400> 109
gagatggggg tctcgctgtg ttgcccaggc tgggtcttga ctcaagcaat ctgcctgtct 60
cagcctacca aaatgctgga ttatag 86
<210> 110
<211> 116
<212> DNA
<213> Homo sapiens
<400> 110
gctaaagcag tcttcctgag tagttaggac tacagacata cacgtgccac cgcgccagc 60
tccgtgttct cttgttttcc ctgcctcctg ctcttccact tatctttgca tggcag 116
<210> 111
<211> 45
<212> DNA
<213> Homo sapiens
<400> 111
ggaaagacga tggagggcag cgaagagagt caccgaccat gacgg 45
<210> 112
<211> 5138
<212> DNA
<213> Homo sapiens
<220>
<221> misc_feature
<222> 31..33
<223> ATG
<221> misc_feature

```

<223> TAG

<221> polyA\_signal

<222> 5111..5116

<223> AATAAA

<400> 112

ctgctgtccc tgggtgtcca cacgtactcc atg cgc tac ctg ctg ccc agc gtc 54  
Met Arg Tyr Leu Leu Pro Ser Val  
1 5

gtg ctc ctg ggc acg gcg ccc acc tac gtg ttg gcc tgg ggg gtc tgg 102  
Val Leu Leu Gly Thr Ala Pro Thr Tyr Val Leu Ala Trp Gly Val Trp

10 15 20  
cgg ctg ctc tcc gcc ttc ctg ccc gcc cgc ttc tac caa gcg ctg gac 150

Arg Leu Leu Ser Ala Phe Leu Pro Ala Arg Phe Tyr Gln Ala Leu Asp  
25 30 35 40

gac cgg ctg tac tgc gtc tac cag<sup>c</sup> agc atg gtg ctc ttc ttc ttc gag 198  
Asp Arg Leu Tyr Cys Val Tyr Gln Ser Met Val Leu Phe Phe Phe Glu  
45 50 55

aat tac acc ggg gtc cag ttg act gga ttg ttg ctg aca tct tgg cca 246  
Asn Tyr Thr Gly Val Gln Leu Thr Gly Leu Leu Leu Thr Ser Trp Pro  
60 65 70

tca ggc aga atg cgc tag gacatgtgcg ctacgtgctg aaagaagggt 294  
Ser Gly Arg Met Arg \*  
75

taaaatggct	gccattgtat	gggtgttact	ttgctcagca	tggaggaatc	tatgtaaagc	354
gcagtgccaa	atttaacgag	aaagagatgc	gaaacaagtt	gcagagctac	gtggagcgcag	414
gaactccaat	gtatcttggt	atttttccag	aagggtacaag	gtataatcca	gagcaaacaa	474
aagtcccttc	agctagtcag	gcatttgctg	cccaacgttg	ccttgccagta	ttaaaacatg	534
tgctaacacc	acgaataaag	gcaactcacg	ttgcttttga	ttgcatgaag	aattatttag	594
atgcaattta	tgatgtttacg	ttggtttatg	aagggaaaga	cgatggagggg	cagcgaagag	654
agtcaccgac	catgacggaa	tttctctgca	aagaatgtcc	aaaaattcat	attcacattg	714
atcgtatcga	caaaaaagat	gtcccagaag	aacaagaaca	tatgagaaga	tggctgcattg	774
aacgtttcga	aatcaaagat	aagatgctta	tagaatttta	tgagtcacca	gatccagaaa	834
gaagaaaaag	atttcctggg	aaaagtgtta	attccaaatt	aagtatcaag	aagactttac	894
catcaatgtt	gatcttaagt	ggtttgactg	caggcatgct	tatgaccgat	ctgggaagga	954
agctgtatgt	gaacacctgg	atataatgaa	ccctacttgg	ctgcctgtgg	gttactatta	1014
aagcatagac	aagtagctgt	ctccagacag	tgggatgtgc	tacattgtct	atttttggcg	1074
gctgcacatg	acatcaaatt	gtttcctgaa	ttattaagg	agtgtaaata	aagccttggt	1134
gattgaagat	tggataaatt	aatttgtgac	gaaagctgat	atgcaattgt	cttgggcgaa	1194
catactgggt	tgtacaactt	tagcatcggt	gctgctggaa	gggtaaaagg	taaattggaat	1254
ttctcctgct	gtctccattt	cctatgaact	aatgacaact	tgagaaggct	gggaggattg	1314
tgtattttgc	aagtcagatg	gctgcatttt	tgagcattaa	tttgcagcgt	atttcacttt	1374
ttctgttatt	ttcaatttat	tacaacttga	cagctccaag	ctcttattac	taaagtattt	1434
agtatcttgc	agtgatttaa	tatttcatct	tttgccttatt	tctacaagtc	agtgaataaa	1494
attgtattta	ggaagtgtca	ggatgttcaa	aggaaaaggt	aaaaagtggt	catggggaaa	1554
aagctctgtt	tagcacatga	ttttattgta	ttgcgttatt	agctgatttt	actcatttta	1614
tatttgcaaa	atpaaattct	aatatattat	gaaattgtct	aatttgcaca	ccctgtacac	1674
acagaaaatg	gtataaaaata	tgagaacgaa	gtttaaaatt	gtgactctga	ttcattatag	1734
cagaacttta	aatttcccag	ctttttgaag	atttaagcta	cgctattagt	acttcccttt	1794
gtctgtgcca	taagtgtctg	aaaacgttaa	ggttttctgt	tttgttttgt	ttttttaata	1854
tcaaaagagt	cggtgtgaac	cttggttgga	ccccaaagtc	acaagatttt	taaggtgatg	1914
agagcctgca	gacattctgc	ctagattttac	tagcgtgtgc	cttttgccctg	cttctctttg	1974
atttcacaga	atattcattc	agaagtcgcg	tttctgtagt	gtggtggatt	cccactgggc	2034
tctggctcct	cccttggatc	ccgtcagtg	tgctgctcag	cggtctgcac	gtagacttgc	2094
taggaagaaa	tgcagagcca	gctctgtgct	cccactttca	gagttgaact	ctttaagccc	2154
ttgtgagtgg	gcttcaccag	ctactgcaga	ggcattttgc	atttgtctgt	gtcaagaagt	2214
tcacctttct	aagccagtga	aatacagact	taattcgtca	tgactgaacg	aatttgttta	2274
tttcccattha	ggttttagtg	agctacacat	taatatgtat	cgccttagag	caagagctgt	2334
gttccaggaa	ccagatcacg	attttttagc	atggaacaat	atatcccatg	ggagaagacc	2394
tttcagtggt	aactgtttcta	tttttgtgtt	ataattttaa	cttcgatttc	ctcatagctc	2454
ttttaagttga	catttctgct	tactgctact	ggatttttgc	tgcagaataa	tatcagtggc	2514

```

ccacattaaa cataccagtt ggatcatgat aagcaaaatg aaagaaataa tgattaaggg 2574
aaaattaagt gactgtgtta cactgcttct cccatgccag agaataaact ctttcaagca 2634
tcatctttga agagtcgtgt ggtgtgaatt ggtttgtgta cattagaatg tatgcacaca 2694
tccatggaca ctcaggatat agttggccta ataatacggg catgggtaaa acttatgaaa 2754
atttcctcat gctgaattgt aattttctct tacctgtaaa gtaaaattta gatcaattcc 2814
atgtctttgt taagtacagg gatttaatat attttgaata taatgggtat gttctaaatt 2874
tgaactttga gaggcaatac tgttggaatt atgtggattc taactcattt taacaaggta 2934
gcctgacctg cataagatca cttgaatgtt aggtttcata gaactatact aatcttctca 2994
caaaaggtct ataaaatata gtcgttgaaa aaaattttgt atcaaaatgt ttggaaaatt 3054
agaagcttct ccttaacctg tattgatact gacttgaatt attttctaaa attagagcc 3114
gtatacctac ctgtaagtct tttcacatat catttaaact tttgtttgta ttattactga 3174
tttacagctt agttattaat ttttctttat aagaatgccg tcgatgtgca tgcttttatg 3234
tttttcagaa aaggggtgtg ttggatgaaa gtaaaaaaaa aaataaaatc tttcactgtc 3294
tctaattggt gtgctgttta acattttttg accctaaaat tcaccaacag tctcccagta 3354
cataaaatag gcttaattgac tggccctgca ttcttcacaa tatttttccc taagctttga 3414
gcaaagtttt aaaaaaatac actaaaataa tcaaaactgt taagcagtat attagtttgg 3474
ttatataaat tcatctgcaa ttataagat gcattggcga tgttaatttg cttggcaatt 3534
ctgtaatcat taagtgatct cagtgaacaa tgtcaaatgc cttaaattaa ctaagttggg 3594
gaataaaagt gccgatctgg ctaactctta caccatacat actgatagtt tttcatatgt 3654
ttcattttcca tgtgattttt aaaatttaga gtggcaacaa ttttgcttaa tatgggttac 3714
ataagcttta ttttttctct tgttcataat tatattcttt gaataggctc gtgtcaatca 3774
agtgatctaa ctgactgat catagataga aggaaataag gccaagttca agaccagcct 3834
gggcaacata tcgagaacct gtctacaaaa aaattaaaaa aaattagcca ggcatgggtg 3894
cgtacactga gtagtttgtc ccagctactc gggagggtga ggtgggagga tcgcttcagc 3954
ccaggagggt gagattgcag tgagccatgg acataccact gcactacagc ctaggtaaca 4014
gcacgagacc ccaactctta gaaaatgaaa aggaaatata gaaatataaa atttgcttat 4074
tatagacaca cagtaactcc cagatatgta ccacaaaaaa tgtgaaaaga gagagaaatg 4134
tctaccaaag cagtattttg tgtgtataat tgcaagcgca tagtaaaata attttaacct 4194
taatttgttt ttagtagtgt ttagattgaa gattgagtga aatattttct tggcagatat 4254
tccgtatctg gtggaaagct acaatgcaat gtcgttgtag ttttgcatgg cttgctttat 4314
aaacaagatt ttttctccct ccttttgggc cagttttcat tacgagtaac tcacactttt 4374
tgattaaaga acttgaatt acgttatcac ttagtataat tgacattata tagagactat 4434
gtaacatgca atcattagaa tcaaaattag tactttggtc aaaatattta caacattcac 4494
atacttgtca aatattcatg taattaactg aatttaaaac cttcaactat tatgaagtgc 4554
tcgtctgtac aatcgctaatt ttactcagtt tagagtagct acaactcttc gatactatca 4614
tcaatatttg acatcttttc caatttggtg atgaaaagta aatctattcc tgtagcaact 4674
ggggagtgat atatgaggtc aaagacatat accttggtat tataatatgt atactataat 4734
aatagctggg tatcctgagc aggggaaaag gttattttta ggaaaaccac ttcaaataga 4794
aagctgaagt acttctaata tactgaggga agtataatat gtggaacaaa ctctcaacaa 4854
aatgtttatt gatgttgatg aaacagatca gtttttccat cgggattatt attggttcat 4914
gattttatat gtgaatatgt aagatatgtt ctgcaatttt ataaatgttc atgtcttttt 4974
ttaaaaaagg tgttattgaa attctgtgtc tccagcaggc aagaatactt gactaactct 5034
ttttgtctct ttatggtatt ttcagaataa agtctgactt gtgtttttga gattattggg 5094
gcctcattaa ttcagcaata aaggaaaata tgcactctaa aaat 5138

```

&lt;210&gt; 113

&lt;211&gt; 5224

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; 31..33

&lt;223&gt; ATG

&lt;221&gt; misc\_feature

&lt;222&gt; 262..264

&lt;223&gt; TAG

&lt;221&gt; polyA\_signal

&lt;222&gt; 5197..5202

&lt;223&gt; AATAAA

&lt;400&gt; 113

```

ctgctgtccc tgggtgtcca cacgtactcc atg cgc tac ctg ctg ccc agc gtc
Met Arg Tyr Leu Leu Pro Ser Val

```

54



ggatttctaac	tcatttttaac	aaggtagcct	gacctgcata	agatcacttg	aatgttaggt	3054
ttcatagaac	taactaatac	ttctcacaaa	aggtctataa	aatacagtcg	ttgaaaaaaa	3114
ttttgtatca	aaatgttttg	aaaattagaa	gcttctcctt	aacctgtatt	gatactgact	3174
tgaattattt	tctaaaatta	agagccgtat	acctacctgt	aagtcttttc	acatatcatt	3234
taaaactttt	tttctgattat	tactgattta	cagcttagtt	attaattttt	ctttataaga	3294
atgccgtcga	tgtgcatgct	tttatgtttt	tcagaaaagg	gtgtgttttg	atgaaagtaa	3354
aaaaaaaaat	aaaatctttc	actgtctcta	atggctgtgc	tgtttaacat	tttttgacc	3414
taaaattcac	caacagcttc	ccagtacata	aaataggctt	aatgactggc	cctgcattct	3474
tcacaatatt	tttccctaag	ctttgagcaa	agttttaaaa	aaatacacta	aaataatcaa	3534
aactgttaag	cagtatatata	gtttggttat	ataaattcat	ctgcaattta	taagatgcat	3594
ggccgatgtt	aatttgcttg	gcaattctgt	aatcattaag	tgatctcagt	gaaacatgtc	3654
aaatgcctta	aattaactaa	gtttggtgaat	aaaagtgcg	atctgggctaa	ctcttacacc	3714
atacatactg	atagtttttc	atatgtttca	tttccatgtg	atttttataa	tttagagtgg	3774
caacaatttt	gcttaatatg	ggttacataa	gctttatttt	ttcctttgtt	cataattata	3834
ttctttgaat	aggtctgtgt	caatcaagt	atctaactag	actgatcata	gatagaagga	3894
aataaggcca	agttcaagac	cagcctgggc	aacatatacga	gaacctgtct	acaaaaaat	3954
taaaaaaaat	tagccaggca	tggtggcgta	cactgagtag	tttgtcccag	ctactcgga	4014
gggtgaggtg	ggaggatcgc	ttcagcccag	gaggttgaga	ttgcagtggg	ccatggacat	4074
accactgcac	tacagcctag	gtaacagcac	gagaccccaa	ctcttagaaa	atgaaaagga	4134
aatatagaaa	tataaaaattt	gcttattata	gacacacagt	aactcccaga	tatgtaccac	4194
aaaaaatgtg	aaaagagaga	gaaatgtcta	ccaaagcagt	attttgtgtg	tataattgca	4254
agcgcatagt	aaaataattt	taaccttaat	ttgttttttag	tagtgttttag	attgaagatt	4314
gagtgaataa	ttttcttggc	agatattccg	tatctgggtg	aaagctacaa	tgcaatgtcg	4374
ttgtagtttt	gcatggcttg	ctttataaac	aagatttttt	ctccctcctt	ttgggacagt	4434
tttcattacg	agtaactcac	actttttgat	taaagaactt	gaaattacgt	tatcacttag	4494
tataattgac	atpatataga	gactatgtaa	catgcaatca	ttagaatcaa	aattagtact	4554
ttggtcaaaa	tatttacaac	attcacatac	ttgtcaaata	ttcatgtaat	taactgaatt	4614
taaaaccttc	aactattatg	aagtgtcctg	ctgtacaatc	gctaattttac	tcagtttaga	4674
gtagctacaa	cttttcgata	ctatcatcaa	tatttgacat	cttttccaat	ttgtgtatga	4734
aaagtaaata	tattcctgta	gcaactgggg	agtcataatat	gaggtcaaag	acataacac	4794
tggtattata	atatgtatac	tataataata	gctggttata	ctgagcaggg	gaaaagggtta	4854
tttttaggaa	aaccacttca	aatagaaaag	tgaagtactt	ctaataact	gaggggaagta	4914
taatatgtgg	aacaaactct	caacaaaatg	tttattgatg	ttgatgaaac	agatcagttt	4974
ttccatccgg	attattattg	gttcatgatt	ttatatgtga	atatgtaaga	tatgttctgc	5034
aattttataa	atgttcatgt	ctttttttta	aaaagggtg	attgaaattc	tgtgtctcca	5094
gcaggcaaga	atacttgact	aactcttttt	gtctctttat	ggatattttca	gaataaagtc	5154
tgacttgtgt	ttttgagatt	attggtgcct	cattaattca	gcaataaagg	aaaatatgca	5214
tctcaaaaat						5224
<210>	114					
<211>	4863					
<212>	DNA					
<213>	Homo sapiens					
<220>						
<221>	misc_feature					
<222>	31..33					
<223>	ATG					
<221>	misc_feature					
<222>	745..747					
<223>	TAG					
<221>	polyA_signal					
<222>	4836..4841					
<223>	AATAAA					
<400>	114					
ctgctgtccc	tggtgtccca	cacgtactcc	atg cgc tac ctg ctg ccc agc gtc			54
			Met Arg Tyr Leu Leu Pro Ser Val			
			1 5			
gtg ctc ctg ggc acg gcg ccc acc tac gtg ttg gcc tgg ggg gtc tgg						102
Val Leu Leu Gly Thr Ala Pro Thr Tyr Val Leu Ala Trp Gly Val Trp						
10 15 20						
cgg ctg ctc tcc gcc ttc ctg ccc gcc cgc ttc tac caa gcg ctg gac						150
Arg Leu Leu Ser Ala Phe Leu Pro Ala Arg Phe Tyr Gln Ala Leu Asp						

25	30	35	40	
gac cgg ctg tac tgc gtc tac cag agc atg gtg ctc ttc ttc ttc gag				198
Asp Arg Leu Tyr Cys Val Tyr Gln Ser Met Val Leu Phe Phe Phe Glu				
45	50	55		
aat tac acc ggg gtc cag cat gga gga atc tat gta aag cgc agt gcc				246
Asn Tyr Thr Gly Val Gln His Gly Gly Ile Tyr Val Lys Arg Ser Ala				
60	65	70		
aaa ttt aac gag aaa gag atg cga aac aag ttg cag agc tac gtg gac				294
Lys Phe Asn Glu Lys Glu Met Arg Asn Lys Leu Gln Ser Tyr Val Asp				
75	80	85		
gca gga act cca atg tat ctt gtg att ttt cca gaa ggt aca agg tat				342
Ala Gly Thr Pro Met Tyr Leu Val Ile Phe Pro Glu Gly Thr Arg Tyr				
90	95	100		
aat cca gag caa aca aaa gtc ctt tca gct agt cag gca ttt gct gcc				390
Asn Pro Glu Gln Thr Lys Val Leu Ser Ala Ser Gln Ala Phe Ala Ala				
105	110	115		
caa cgt gaa ttt ctc tgc aaa gaa tgt cca aaa att cat att cac att				438
Gln Arg Glu Phe Leu Cys Lys Glu Cys Pro Lys Ile His Ile His Ile				
125	130	135		
gat cgt atc gac aaa aaa gat gtc cca gaa gaa caa gaa cat atg aga				486
Asp Arg Ile Asp Lys Lys Asp Val Pro Glu Glu Gln Glu His Met Arg				
140	145	150		
aga tgg ctg cat gaa cgt ttc gaa atc aaa gat aag atg ctt ata gaa				534
Arg Trp Leu His Glu Arg Phe Glu Ile Lys Asp Lys Met Leu Ile Glu				
155	160	165		
ttt tat gag tca cca gat cca gaa aga aga aaa aga ttt cct ggg aaa				582
Phe Tyr Glu Ser Pro Asp Pro Glu Arg Arg Lys Arg Phe Pro Gly Lys				
170	175	180		
agt gtt aat tcc aaa tta agt atc aag aag act tta cca tca atg ttg				630
Ser Val Asn Ser Lys Leu Ser Ile Lys Lys Thr Leu Pro Ser Met Leu				
185	190	195		
atc tta agt ggt ttg act gca ggc atg ctt atg acc gat gct gga agg				678
Ile Leu Ser Gly Leu Thr Ala Gly Met Leu Met Thr Asp Ala Gly Arg				
205	210	215		
aag ctg tat gtg aac acc tgg ata tat gga acc cta ctt ggc tgc ctg				726
Lys Leu Tyr Val Asn Thr Trp Ile Tyr Gly Thr Leu Leu Gly Cys Leu				
220	225	230		
tgg gtt act att aaa gca tag acaagtagct gtctccagac agtgggatgt				777
Trp Val Thr Ile Lys Ala *				
235				
gctacattgt ctatttttgg cggctgcaca tgacatcaaa ttgtttcctg aatttattaa				837
ggagtgtaaa taagccttg ttgattgaag attggataat agaatttgtg acgaaagctg				897
atatgcaatg gtcttgggca aacataacctg gttgtacaac tttagcatcg gggctgctgg				957
aagggtaaaa gctaaatgga gtttctcctg ctctgtccat ttcctatgaa ctaatgacaa				1017
cttgagaagg ctgggaggat tgtgtatttt gcaagtcaga tggctgcatt tttgagcatt				1077
aatttgcagc gtatttcact ttttctgtta ttttcaattt attacaactt gacagctcca				1137
agctcttatt actaaagtat ttagtatctt gcagctagtt aatatttcat cttttgctta				1197
tttctacaag tcagtgaat aaattgtatt taggaagtgt caggatgttc aaaggaaagg				1257
gtaaaaagtg ttpatgggga aaaagctctg ttttagcacat gattttattg tattgcgtta				1317
ttagctgatt ttactcattt tatatttgca aaataaattt ctaatattta ttgaaattgc				1377
ttaatttgca caccctgtac acacagaaaa tggataaaaa tatgagaacg aagtttaaaa				1437
ttgtgactct gattcattat agcagaactt taaatttccc agctttttga agatttaagc				1497
tacgctatta gtacttccct ttgtctgtgc cataagtgtc tgaaaacggt aaggttttct				1557
gttttgtttt gtttttttaa tatcaaaaga gtcgggtgtga accttggttg gaccccaagt				1617
tcacaagatt ttttaaggta tgagagcctg cagacattct gcctagattt actagcgtgt				1677
gccttttgcc tgcttctctt tgatttcaca gaatttcat tcagaagtcg cgtttctgta				1737
gtgtggtgga ttccactgg gctctggtcc ttcccttggga tcccgtcagt ggtgctgtc				1797
agcggcttgc acgtagactt gctaggaaga aatgcagagc cagcctgtgc tgccacttt				1857
cagagttgaa ctctttaagc ccttgtgagt gggcttcacc agctactgca gaggcatttt				1917
gcatttgtct gtgtcaagaa gttcaccttc tcaagccagt gaaatacaga cttaattcgt				1977
catgactgaa cgaatttgtt tatttcccat taggtttagt ggagctacac attaatatgt				2037

atcgcccttag	agcaagagct	gtgttccagg	aaccagatca	cgatttttag	ccatggaaca	2097
atatatccca	tgggagaaga	cctttcagtg	tgaactgttc	tatttttgtg	ttataattta	2157
aacttcgatt	tcctcatagt	cctttaagtt	gacatttctg	cttactgcta	ctggattttt	2217
gctgcagaaa	tatatcagtg	gccacatta	aacataccag	ttggatcatg	ataagcaaaa	2277
tgaagaaat	aatgattaag	ggaaaattaa	gtgactgtgt	tacactgctt	ctcccatgcc	2337
agagaataaa	ctctttcaag	catcatcttt	gaagagtcgt	gtggtgtgaa	ttggtttgtg	2397
tacattagaa	tgtatgcaca	catccatgga	cactcaggat	atagttggcc	taataatcgg	2457
ggcatgggta	aaacttatga	aaatttcctc	atgctgaatt	gtaattttct	cttacctgta	2517
aagtataaatt	tagatcaatt	ccatgtcttt	gttaagtaca	gggatttaat	atattttgaa	2577
tataatgggt	atgttctaaa	tttgaacttt	gagaggcaat	actggttgaa	ttatgtggat	2637
tctaactcat	tttaacaagg	tagcctgacc	tgcataagat	cacttgatg	ttaggtttca	2697
tagaactata	ctaactctct	cacaaaagg	ctataaaata	cagtcgttga	aaaaaatttt	2757
gtatcaaaa	gttgggaaa	ttagaagctt	ctccttaacc	tgtattgata	ctgacttgaa	2817
ttattttcta	aaattaagag	ccgtatacct	acctgttaagt	cttttcacat	atcattttaaa	2877
cttttggttg	tattattact	gatttacagc	ttagttatta	atttttcttt	ataagaatgc	2937
cgtcgatgtg	catgctttta	tgtttttcag	aaaagggtgt	gtttggatga	aagtaaaaaa	2997
aaaaataaaa	tctttcactg	tctctaattg	ctgtgctgtt	taacattttt	tgaccctaaa	3057
attcaccaac	agtctcccag	tacataaaat	aggcttaattg	actggccctg	cattcttcac	3117
aatatttttc	cctaagcttt	gagcaaaagt	ttaaaaaaat	acactaaaat	aatcaaaaat	3177
gttaagcagt	atattagttt	ggttatataa	attcatctgc	aattttataag	atgcatggcc	3237
gatgttaatt	tgcttggcaa	ttctgtaatc	attaagtgat	ctcagtgaat	catgtcaaat	3297
gccttaaat	aactaagtgt	gtgaataaaa	gtgccgatct	ggctaactct	tacaccatac	3357
atactgatag	tttttcatat	gtttcatttc	catgtgattt	ttaaaattta	gagtggcaac	3417
aattttgctt	aatatgggtt	acataagctt	tattttttcc	tttgttcata	attatattct	3477
ttgaataggt	ctgtgtcaat	caagtgatct	aactgactgt	atcatagata	gaaggaaata	3537
aggccaaggt	caagaccagc	ctgggcaaca	tatcgagaac	ctgtctacaa	aaaaattaaa	3597
aaaaattagc	caggcatggt	ggcgtacact	gagtagtttg	tcccagctac	tggggagggt	3657
gaggtgggag	gatcgcttca	gccaggagg	ttgagattgc	agtgagccat	ggacatacca	3717
ctgcactaca	gcctaggtaa	cagcacgaga	ccccaaactc	tagaaaatga	aaaggaaata	3777
tagaaatata	aaatttgctt	attatagaca	cacagtaact	cccagatatg	taccacaaaa	3837
aatgtgaaaa	gagagagaaa	tgtctaccaa	agcagtattt	tgtgtgtata	attgcaagcg	3897
catagtaaaa	taattttaac	cttaatttgc	tttttagtagt	gttttagattg	aagattgagt	3957
gaaatatttt	cttggcagat	attccgtatc	tgggtggaaa	ctacaatgca	atgtcgttgt	4017
agttttgcat	ggcttgcttt	ataaacaaga	ttttttctcc	ctccttttgg	gccagttttc	4077
attacgagta	actcacactt	tttgattaaa	gaacttgaaa	ttacgttatc	acttagtata	4137
attgacatta	tatagagact	atgtaacatg	caatcattag	aatcaaaaat	agtactttgg	4197
tcaaaatatt	tacaacattc	acatacttgt	caaataattca	tgtaatatac	tgaatttaaa	4257
accttcaact	attatgaagt	gctcgtctgt	acaatcgcta	atttactcag	tttagagtag	4317
ctacaactct	tctgatactat	catcaatatt	tgacatcttt	tccaatttgt	gtatgaaaag	4377
taaatctatt	ctgtagcaa	ctggggagtc	atataatgagg	tcaaagacat	ataccttggt	4437
attataatat	gtataactata	ataatagctg	gttatcctga	gcaggggaaa	aggttatttt	4497
taggaaaacc	acttcaaata	gaaagctgaa	gtacttctaa	tatactgagg	gaagtataat	4557
atgtggaaca	aactctcaac	aaaatgttta	ttgatgttga	tgaacagat	cagtttttcc	4617
atccggatta	ttattgggtc	atgattttat	atgtgaatat	gtaagatatg	ttctgcaatt	4677
ttataaatgt	tcatgtcttt	ttttaaaaaa	ggtgctattg	aaattctgtg	tctccagcag	4737
gcaagaatac	ttgactaact	ctttttgtct	ctttatggta	ttttcagaat	aaagtctgac	4797
ttgtgttttt	gagattattg	gtgcctcatt	aattcagcaa	taaaggaaaa	tatgcatctc	4857
aaaaat						4863
<210>	115					
<211>	5022					
<212>	DNA					
<213>	Homo sapiens					
<220>						
<221>	misc_feature					
<222>	31..33					
<223>	ATG					
<221>	misc_feature					
<222>	904..906					
<223>	TAG					
<221>	polyA_signal					
<222>	4995..5000					



&lt;223&gt; AATAAA

&lt;400&gt; 115

```

ctgctgtccc tgggtgtcca cacgtactcc atg cgc tac ctg ctg ccc agc gtc      54
                               Met Arg Tyr Leu Leu Pro Ser Val
                               1           5

gtg ctc ctg ggc acg gcg ccc acc tac gtg ttg gcc tgg ggg gtc tgg      102
Val Leu Leu Gly Thr Ala Pro Thr Tyr Val Leu Ala Trp Gly Val Trp
10           15           20

cgg ctg ctc tcc gcc ttc ctg ccc gcc cgc ttc tac caa gcg ctg gac      150
Arg Leu Leu Ser Ala Phe Leu Pro Ala Arg Phe Tyr Gln Ala Leu Asp
25           30           35           40

gac cgg ctg tac tgc gtc tac cag agc atg gtg ctc ttc ttc ttc gag      198
Asp Arg Leu Tyr Cys Val Tyr Gln Ser Met Val Leu Phe Phe Phe Glu
45           50           55

aat tac acc ggg gtc cag cat gga gga atc tat gta aag cgc agt gcc      246
Asn Tyr Thr Gly Val Gln His Gly Gly Ile Tyr Val Lys Arg Ser Ala
60           65           70

aaa ttt aac gag aaa gag atg cga aac aag ttg cag agc tac gtg gac      294
Lys Phe Asn Glu Lys Glu Met Arg Asn Lys Leu Gln Ser Tyr Val Asp
75           80           85

gca gga act cca atg tat ctt gtg att ttt cca gaa ggt aca agg tat      342
Ala Gly Thr Pro Met Tyr Leu Val Ile Phe Pro Glu Gly Thr Arg Tyr
90           95           100

aat cca gag caa aca aaa gtc ctt tca gct agt cag gca ttt gct gcc      390
Asn Pro Glu Gln Thr Lys Val Leu Ser Ala Ser Gln Ala Phe Ala Ala
105          110          115          120

caa cgt ggc ctt gca gta tta aaa cat gtg cta aca cca cga ata aag      438
Gln Arg Gly Leu Ala Val Leu Lys His Val Leu Thr Pro Arg Ile Lys
125          130          135

gca act cac gtt gct ttt gat tgc atg aag aat tat tta gat gca att      486
Ala Thr His Val Ala Phe Asp Cys Met Lys Asn Tyr Leu Asp Ala Ile
140          145          150

tat gat gtt acg gtg gtt tat gaa ggg aaa gac gat gga ggg cag cga      534
Tyr Asp Val Thr Val Val Tyr Glu Gly Lys Asp Asp Gly Gly Gln Arg
155          160          165

aga gag tca cgg acc atg acg gaa ttt ctc tgc aaa gaa tgt cca aaa      582
Arg Glu Ser Pro Thr Met Thr Glu Phe Leu Cys Lys Glu Cys Pro Lys
170          175          180

att cat att cac att gat cgt atc gac aaa aaa gat gtc cca gaa gaa      630
Ile His Ile His Ile Asp Arg Ile Asp Lys Lys Asp Val Pro Glu Glu
185          190          195          200

caa gaa cat atg aga aga tgg ctg cat gaa cgt ttc gaa atc aaa gat      678
Gln Glu His Met Arg Arg Trp Leu His Glu Arg Phe Glu Ile Lys Asp
205          210          215

aag atg ctt ata gaa ttt tat gag tca cca gat cca gaa aga aga aaa      726
Lys Met Leu Ile Glu Phe Tyr Glu Ser Pro Asp Pro Glu Arg Arg Lys
220          225          230

aga ttt cct ggg aaa agt gtt aat tcc aaa tta agt atc aag aag act      774
Arg Phe Pro Gly Lys Ser Val Asn Ser Lys Leu Ser Ile Lys Lys Thr
235          240          245

tta cca tca atg ttg atc tta agt ggt ttg act gca ggc atg ctt atg      822
Leu Pro Ser Met Leu Ile Leu Ser Gly Leu Thr Ala Gly Met Leu Met
250          255          260

acc gat gct gga agg aag ctg tat gtg aac acc tgg ata tat gga acc      870
Thr Asp Ala Gly Arg Lys Leu Tyr Val Asn Thr Trp Ile Tyr Gly Thr
265          270          275          280

cta ctt ggc tgc ctg tgg gtt act att aaa gca tag acaagtagct      916
Leu Leu Gly Cys Leu Trp Val Thr Ile Lys Ala *
285          290

gtctccagac agtgggatgt gctacattgt ctatttttgg cggctgcaca tgacatcaaa      976
ttgtttcctg aatttattaa ggagtgtaaa taaagccttg ttgattgaag attggataat      1036

```

agaatttgtg	acgaaagctg	atatgcaatg	gtcttgggca	aacatacctg	gttgataaac	1096
tttagcatcg	gggctgctgg	aagggtaaaa	gctaaatgga	gtttctcctg	ctctgtccat	1156
ttcctatgaa	ctaatagcaa	cttgagaagg	ctgggaggat	tgtgtatttt	gcaagtcaga	1216
tggtgcatt	tttgagcatt	aatttgcagc	gtatttcaact	ttttctgtta	ttttcaattt	1276
attacaactt	gacagctcca	agctcttatt	actaaagtat	ttagtatctt	gcagctagtt	1336
aatatttcat	cttttgctta	tttctacaag	tcagtgaat	aaattgtatt	taggaagtgt	1396
caggatgttc	aaaggaaagg	gtaaaaagtg	ttcatgggga	aaaagctctg	tttagcacat	1456
gattttattg	tattgcgtta	ttagctgatt	ttactcattt	tatatttgca	aaataaattt	1516
ctaattattt	ttgaaattgc	ttaatgtgca	cacctgtac	acacagaaaa	tggtataaaa	1576
tatgagaacg	aagtttaaaa	ttgtgactct	gattcattat	agcagaactt	taaatttccc	1636
agctttttga	agatttaagc	tacgtctatta	gtacttccct	ttgtctgtgc	cataagtgtc	1696
tgaaaacggt	aagggttttct	gttttgtttt	gtttttttta	tatcaaaaga	gtcgggtgtg	1756
accttggttg	gaccccaagt	tcacaagatt	tttaagggtg	tgagagcctg	cagacattct	1816
gcctagattt	actagcgtgt	gccttttgcc	tgcttctctt	tgatttcaca	gaatattcat	1876
tcagaagtcg	cgtttctgta	gtgtggtgga	ttccactggg	gctctggtcc	ttcccttgga	1936
tcccgctcagt	ggtgctgctc	agcggcttgc	acgtagactt	gctaggaaga	aatgcaagagc	1996
cagcctgtgc	tgccactttt	cagagttgaa	ctctttaagc	ccttgtagt	gggcttcacc	2056
agctactgca	gaggcatttt	gcatttgtct	gtgtcaagaa	gttcaccttc	tcaagccagt	2116
gaaatacaga	cttaattcgt	catgactgaa	cgaatttgtt	tatttcccat	taggtttagt	2176
ggagctacac	attaatatgt	atcgcttag	agcaagagct	gtgttccagg	aaccagatca	2236
cgatttttag	ccatggaaca	atataccca	tgggagaaga	cctttcagtg	tgaactgttc	2296
tatttttgtg	ttataattta	aacttcgatt	tcctcatagt	cctttaagtt	gacatttctg	2356
cttactgcta	ctggattttt	gctgcagaaa	tatatcagtg	gcccacatta	aacataccag	2416
ttggatcatg	atpagcaaaa	tgaaagaaat	aatgattaa	ggaaaattaa	gtgactgtgt	2476
tacactgctt	ctbccatgcc	agagaataaa	ctcttcaag	catcatcttt	gaagagtcgt	2536
gtggtgtgaa	ttggtttgtg	tacattagaa	tgtatgcaca	catccatgga	cactcaggat	2596
atagttggcc	taataatcgg	ggcatgggta	aaacttatga	aaatttcctc	atgctgaatt	2656
gtaattttct	cttacctgta	aagtaaaatt	tagatcaatt	ccatgtcttt	gttaagtaca	2716
gggattttaa	atattttgaa	tataatgggt	atgttctaaa	tttgaacttt	gagaggcaat	2776
actgttgga	ttatgtgat	tctaactcat	tttaacaagg	tagcctgacc	tgcaataagat	2836
cacttgaatg	ttaggtttca	tagaactata	ctaattctct	cacaaaaggt	ctataaaata	2896
cagtcgttga	aaaaaatttt	gtatcaaaat	gtttggaaaa	ttagaagctt	ctccttaacc	2956
tgtattgata	ctgacttgaa	ttattttcta	aaattaaagag	ccgtataacct	acctgtaagt	3016
cttttcacat	atcattttaa	cttttgtttg	tattattact	gatttacagc	ttagttatta	3076
atttttcttt	ataagaatgc	cgctgatgtg	catgctttta	tgtttttcag	aaaagggtgt	3136
gtttggatga	aagtaaaaaa	aaaaataaaa	tctttcactg	tctctaattg	ctgtgctgtt	3196
taacattttt	tgaccctaaa	attcaccaac	agtcctccag	tacataaaat	aggcttaatg	3256
actggccctg	cattcttcac	aatatttttc	cctaagcttt	gagcaaaagt	ttaaaaaaat	3316
acactaaaat	aatcaaaact	gttaagcagt	atattagttt	ggttatataa	attcatctgc	3376
aatttataag	atgcatggcc	gatgttaatt	tgcttggcaa	ttctgtaatc	attaagtgat	3436
ctcagtgaat	catgtcaaat	gccttaaat	aactaagttg	gtgaataaaa	gtgccgatct	3496
ggctaactct	tacaccatac	atactgatag	tttttcatat	gtttcatttc	catgtgattt	3556
ttaaaattta	gagtggcaac	aattttgctt	aatatgggtt	acataagctt	tattttttcc	3616
tttgttcata	attatattct	ttgaataggt	ctgtgtcaat	caagtgtatc	aactagactg	3676
atcatagata	gaaggaaata	aggccaagtt	caagaccagc	ctgggcaaca	tatcgagaac	3736
ctgcttacia	aaaaatttaa	aaaaatttagc	caggcatggt	ggcgtaact	gagtagtttg	3796
tccagctac	tcgggagggt	gagggtgggag	gatcgcttca	gccaggagg	ttgagattgc	3856
agtgagccat	ggacatacca	ctgcactaca	gcctaggtaa	cagcacgaga	ccccaaactct	3916
tagaaaatga	aaaggaaata	tagaaatata	aaatttgctt	attatagaca	cacagtaact	3976
cccagatatg	taccacaaaa	aatgtgaaaa	gagagagaaa	tgtctaccaa	agcagtattt	4036
tggtgtgata	attgcaagcg	catagtaaaa	taattttaac	cttaatttgt	tttagtagt	4096
gttttagattg	aagattgagt	gaaatatttt	cttggcagat	attccgtatc	tggtggaaag	4156
ctacaatgca	atgtcgttgt	agttttgcat	ggcttgcttt	ataaacaaga	ttttttctcc	4216
ctccttttgg	gcpagttttc	attacgagta	actcacactt	tttgattaaa	gaacttgaaa	4276
ttacgttatc	acttagtata	attgacatta	tatagagact	atgtaacatg	caatcattag	4336
aatcaaaatt	agtacttttg	tcaaaatatt	tacaacattc	acatacttgt	caaatattca	4396
tgtaatatac	tgaatttaaa	accttcaact	attatgaagt	gctcgtctgt	acaatcgcta	4456
atttactcag	tttagagtag	ctacaactct	tcgatactat	catcaatatt	tgacatcttt	4516
tccaatttgt	gtatgaaaag	taaattctatt	cctgtagcaa	ctggggagtc	atatatgagg	4576
tcaaagacat	ataccttggt	attataatat	gtatactata	ataatagctg	gttatcctga	4636
gcaggggaaa	aggttatttt	taggaaaacc	acttcaaata	gaaagctgaa	gtacttctaa	4696

tatactgagg gaagtataat atgtggaaca aactctcaac aaaatgttta ttgatgttga 4756  
 tgaacacagat cagtttttcc atccggatta ttattggttc atgattttat atgtgaatat 4816  
 gtaagatatg ttctgcaatt ttataaatgt tcatgtcttt ttttaaaaaa ggtgctattg 4876  
 aaattctgtg tctccagcag gcaagaatac ttgactaact ctttttgtct ctttatggta 4936  
 ttttcagaat aaagtctgac ttgtgttttt gagattattg gtgcctcatt aattcagcaa 4996  
 taaaggaaaa tatgcatctc aaaaat 5022  
 <210> 116  
 <211> 4932  
 <212> DNA  
 <213> Homo sapiens  
 <220>  
 <221> misc\_feature  
 <222> 31..33  
 <223> ATG  
 <221> misc\_feature  
 <222> 814..816  
 <223> TAG  
 <221> polyA\_signal  
 <222> 4905..4910  
 <223> AATAAA  
 <400> 116  
 ctgctgtccc tgggtgtcca cacgtactcc atg cgc tac ctg ctg ccc agc gtc 54  
 Met Arg Tyr Leu Leu Pro Ser Val  
 1 5  
 gtg ctc ctg ggc acg gcg ccc acc tac gtg ttg gcc tgg ggg gtc tgg 102  
 Val Leu Leu Gly Thr Ala Pro Thr Tyr Val Leu Ala Trp Gly Val Trp  
 10 15 20  
 cgg ctg ctc tcc gcc ttc ctg ccc gcc cgc ttc tac caa gcg ctg gac 150  
 Arg Leu Leu Ser Ala Phe Leu Pro Ala Arg Phe Tyr Gln Ala Leu Asp  
 25 30 35 40  
 gac cgg ctg tac tgc gtc tac cag agc atg gtg ctc ttc ttc ttc gag 198  
 Asp Arg Leu Tyr Cys Val Tyr Gln Ser Met Val Leu Phe Phe Phe Glu  
 45 50 55  
 aat tac acc ggg gtc cag atg tat ctt gtg att ttt cca gaa ggt aca 246  
 Asn Tyr Thr Gly Val Gln Met Tyr Leu Val Ile Phe Pro Glu Gly Thr  
 60 65 70  
 agg tat aat cca gag caa aca aaa gtc ctt tca gct agt cag gca ttt 294  
 Arg Tyr Asn Pro Glu Gln Thr Lys Val Leu Ser Ala Ser Gln Ala Phe  
 75 80 85  
 gct gcc caa cgt ggc ctt gca gta tta aaa cat gtg cta aca cca cga 342  
 Ala Ala Gln Arg Gly Leu Ala Val Leu Lys His Val Leu Thr Pro Arg  
 90 95 100  
 ata aag gca act cac gtt gct ttt gat tgc atg aag aat tat tta gat 390  
 Ile Lys Ala Thr His Val Ala Phe Asp Cys Met Lys Asn Tyr Leu Asp  
 105 110 115 120  
 gca att tat gat gtt acg gtg gtt tat gaa ggg aaa gac gat gga ggg 438  
 Ala Ile Tyr Asp Val Thr Val Val Tyr Glu Gly Lys Asp Asp Gly Gly  
 125 130 135  
 cag cga aga gag tca ccg acc atg acg gaa ttt ctc tgc aaa gaa tgt 486  
 Gln Arg Arg Glu Ser Pro Thr Met Thr Glu Phe Leu Cys Lys Glu Cys  
 140 145 150  
 cca aaa att cat att cac att gat cgt atc gac aaa aaa gat gtc cca 534  
 Pro Lys Ile His Ile His Ile Asp Arg Ile Asp Lys Lys Asp Val Pro  
 155 160 165  
 gaa gaa caa gaa cat atg aga aga tgg ctg cat gaa cgt ttc gaa atc 582  
 Glu Glu Gln Glu His Met Arg Arg Trp Leu His Glu Arg Phe Glu Ile  
 170 175 180  
 aaa gat aag atg ctt ata gaa ttt tat gag tca cca gat cca gaa aga 630  
 Lys Asp Lys Met Leu Ile Glu Phe Tyr Glu Ser Pro Asp Pro Glu Arg  
 185 190 195 200  
 aga aaa aga ttt cct ggg aaa agt gtt aat tcc aaa tta agt atc aag 678



```

ccccaaactct tagaaaaatga aaaggaaata tagaaatata aaatttgctt attatagaca 3876
cacagtaact cccagatatg taccacaaaa aatgtgaaaa gagagagaaa tgtctaccaa 3936
agcagtatatt tgtgtgtata attgcaagcg catagtaaaa taattttaac ctttaatttgt 3996
tttttagtagt gtttagattg aagattgagt gaaatatatt cttggcagat attccgtatc 4056
tggtggaaag ctacaatgca atgtcgttgt agttttgcat ggcttgcttt ataaacaaga 4116
ttttttctcc ctctttttgg gccagttttc attacgagta actcacactt tttgattaaa 4176
gaacttgaaa ttacgttatc acttagtata attgacatta tatagagact atgtaacatg 4236
caatcattag aatcaaaaatt agtactttgg tcaaaaattt tacaacattc acatacttgt 4296
caaatattca tgaattaac tgaatttaaa accttcaact attatgaagt gctcgtctgt 4356
acaatcgcta atttactcag tttagagtag ctacaactct tcgatactat catcaatatt 4416
tgacatcttt tccaatttgt gtatgaaaag taaatctatt cctgtagcaa ctggggagtc 4476
atatatgagg tcaaagacat ataccttgtt attataatat gtatactata ataatagctg 4536
gttatcctga gcaggggaaa aggttatttt taggaaaacc acttcaata gaaagctgaa 4596
gtacttctaa tatactgagg gaagtataat atgtggaaca aactctcaac aaaatgttta 4656
ttgatgttga tgaacagat cagtttttcc atccggatta ttattggttc atgattttat 4716
atgtgaatat gtaagatatg ttctgcaatt ttataaatgt tcatgtcttt ttttaaaaaa 4776
ggtgctattg aaattctgtg tctccagcag gcaagaatac ttgactaact ctttttgtct 4836
ctttatggta ttttcagaat aaagtctgac ttgtgttttt gagattattg gtgcctcatt 4896
aattcagcaa taaaggaaaa tatgcatctc aaaaaa 4932
<210> 117
<211> 4682
<212> DNA
<213> Homo sapiens
<220>
<221> misc_feature
<222> 31..33
<223> ATG
<221> misc_feature
<222> 301..303
<223> TGA
<221> polyA_signal
<222> 4655..4660
<223> AATAAA
<400> 117
ctgctgtccc tgggtgtcca cacgtactcc atg cgc tac ctg ctg ccc agc gtc 54
Met Arg Tyr Leu Leu Pro Ser Val
1 5
gtg ctc ctg ggc acg gcg ccc acc tac gtg ttg gcc tgg ggg gtc tgg 102
Val Leu Leu Gly Thr Ala Pro Thr Tyr Val Leu Ala Trp Gly Val Trp
10 15 20
cgg ctg ctc tcc gcc ttc ctg ccc gcc cgc ttc tac caa gcg ctg gac 150
Arg Leu Leu Ser Ala Phe Leu Pro Ala Arg Phe Tyr Gln Ala Leu Asp
25 30 35 40
gac cgg ctg tac tgc gtc tac cag agc atg gtg ctc ttc ttc ttc gag 198
Asp Arg Leu Tyr Cys Val Tyr Gln Ser Met Val Leu Phe Phe Phe Glu
45 50 55
aat tac acc ggg gtc cag aat ttc tct gca aag aat gtc caa aaa ttc 246
Asn Tyr Thr Gly Val Gln Asn Phe Ser Ala Lys Asn Val Gln Lys Phe
60 65 70
ata ttc aca ttg atc gta tcg aca aaa aag atg tcc cag aag aac aag 294
Ile Phe Thr Leu Ile Val Ser Thr Lys Lys Met Ser Gln Lys Asn Lys
75 80 85
aac ata tga gaagatggct gcatgaacgt ttcgaaatca aagataagat 343
Asn Ile *
90
gcttatagaa ttttatgagt caccagatcc agaagaaga aaaagatttc ctgggaaaag 403
tggttaattcc aaattaagta tcaagaagac ttaccatca atgttgatct taagtgggtt 463
gactgcaggc atgttatga ccgatgctgg aaggaagctg tatgtgaaca cctggatata 523
tggaacccta ctggctgcc tgtgggttac tattaaagca tagacaagta gctgtctcca 583
gacagtggga tgtgctacat tgtctatttt tggcggctgc acatgacatc aaattgtttc 643
ctgaatttat taaggagtgt aaataaagcc ttgttgattg aagattggat aatagaattt 703

```

gtgacgaaag	ctgatatgca	atgggtcttg	gcaaacatac	ctgggtgtac	aacttttagca	763
tccgggctgc	tggaagggtg	aaagctaaat	ggagtttctc	ctgctctgtc	catttcctat	823
gaactaatga	caacttgaga	aggctgggag	gattgtgtat	tttgcaagtc	agatggctgc	883
atttttgagc	atgaatttgc	agcgtatttc	actttttctg	ttattttcaa	tttattacaa	943
cttgacagct	ccaagctctt	attactaaag	tatttagtat	cttgacagct	gttaaatatt	1003
catcttttgc	ttattttctac	aagtcagtga	aataaattgt	atttaggaag	tgtaggatg	1063
ttcaaaggaa	agggtaaaaa	gtgttcattg	ggaaaaagct	ctgttttagca	catgatttta	1123
ttgtattgcg	ttatttagctg	attttactca	ttttatattt	gcaaaataaa	tttctaata	1183
ttattgaaat	tgcttaattt	gcacaccctg	tacacacaga	aaatgggtata	aaatatgaga	1243
acgaagttta	aaattgtgac	tctgattcat	tatagcagaa	ctttaaattt	cccagctttt	1303
tgaagattta	agctacgcta	ttagtacttc	cctttgtctg	tgccataagt	gcttgaaaaac	1363
gttaagggtt	tctgttttgt	tttgtttttt	taatatcaaa	agagtcggtg	tgaaccttg	1423
ttggaccca	agttcacaa	atttttaagg	tgatgagagc	ctgcagacat	tctgcctaga	1483
tttactagcg	tgtgcctttt	gcctgcttct	ctttgatttc	acagaatatt	cattcagaag	1543
tgcgctttct	gtagtgtggt	ggattcccat	tgggtctctg	tccttccctt	ggatcccgct	1603
agtggtgctg	ctcagcggct	tgcacgtaga	cttgctagga	agaaatgcag	agccagcctg	1663
tgctgcccac	tttcagagtt	gaactcttta	agcccttggt	agtgggcttc	accagctact	1723
gcagaggcat	tttgcatatt	tctgtgtcaa	gaagttcacc	ttctcaagcc	agtgaatac	1783
agacttaatt	cgtcatgact	gaacgaattt	gtttatttcc	cattagggtt	agtggagcta	1843
cacattaata	gttagcgct	tagagcaaga	gctgtgttcc	aggaaccaga	tcacgatttt	1903
tagccatgga	acaatatatc	ccatgggaga	agacctttca	gtgtgaactg	ttctattttt	1963
gtgttataat	ttaaacttcg	atttctctcat	agtcctttta	gttgacattt	ctgcttactg	2023
ctactggatt	tttgctgcag	aaatatatca	gtggccacac	ttaaacatac	cagttggatc	2083
atgataagca	aaatgaaaga	aataatgatt	aagggaatat	taagtgactg	tgttacactg	2143
cttctcccat	gpcagagaat	aaactctttc	aagcatcatc	tttgaagagt	cgtgtggtgt	2203
gaattgggtt	gtgtacatta	gaatgtatgc	acacatccat	ggacactcag	gatatagttg	2263
gcctaataat	cggggcatgg	gtaaaactta	tgaataattc	ctcatgctga	attgtaat	2323
tctcttacct	gtaaagtaaa	atttagatca	attccatgtc	tttgtaagt	acagggattt	2383
aatatatttt	gaatataatg	ggtatgttct	aaatttgaac	tttgagaggg	aatactgttg	2443
gaattatgtg	gatttctaact	cattttaaca	aggtagcctg	acctgcataa	gatcacttga	2503
atgttagggt	tcatagaact	atactaactc	tctcacaaaa	ggtctataaa	atacagtcgt	2563
tgaaaaaaat	tttgtatcaa	aatgtttgga	aaattagaag	cttctcctta	acctgtattg	2623
atactgactt	gaattatttt	ctaaaattaa	gagccgtata	cctacctgta	agtcctttca	2683
catatcattt	aaacttttgt	ttgtattatt	actgatttac	agcttagtta	ttaatTTTTt	2743
tttataagaa	tgccgtcgat	gtgcatgctt	ttatgttttt	cagaaaaggg	tgtgtttgga	2803
tgaagtaaa	aaaaaaaata	aaatctttca	ctgtctctaa	tggctgtgct	gtttaacatt	2863
ttttgacct	aaaattcacc	aacagtcctc	cagtacataa	aataggctta	atgactggcc	2923
ctgcattctt	cacaatattt	ttccctaagc	tttgagcaaa	gttttaaaaa	aatacactaa	2983
aataatcaaa	actgttaagc	agtatattag	tttggttata	taaattcctc	tgcaatttat	3043
aagatgcatg	gccgatgtta	atttgcttgg	caattctgtg	atcattaagt	gatctcagtg	3103
aaacatgtca	aatgccctaa	attaactaag	ttggtgaata	aaagtgccga	tctggctaac	3163
tcttacacca	tacatactga	tagtttttca	tatgtttcat	ttccatgtga	tttttaaaat	3223
ttagagtggc	aacaattttg	cttaatatgg	gttacataag	ctttattttt	tcctttgttc	3283
ataattatat	tctttgaata	ggtctgtgtc	aatcaagtga	tctaactaga	ctgatcatag	3343
atagaaggaa	ataaggccaa	gttcaagacc	agcctgggca	acataatcgag	aacctgtcta	3403
caaaaaaatt	aaaaaaaatt	agccaggcat	ggtagcgtac	actgagtagt	ttgtcccagc	3463
tactcgggag	ggtgaggtgg	gaggatcgct	tcagcccagg	agggtgagat	tgtagtgagc	3523
catggacata	ccactgcact	acagcctagg	taacagcacg	agaccccaac	tcttagaaaa	3583
tgaaggagaa	atatagaaat	ataaaatttg	cttatttatag	acacacagta	actcccagat	3643
atgtaccaca	aaaaatgtga	aaagagagag	aaatgtctac	caaagcagta	ttttgtgtgt	3703
ataattgcaa	gcgcatagta	aaataatttt	aaccttaatt	tgTTTTtagt	agtgtttaga	3763
ttgaagattg	agtgaatat	tttcttggca	gatattccgt	atctggtgga	aagctacaat	3823
gcaatgtcgt	tgtagttttg	cttggtctgc	tttataaaca	agattttttc	tcctctcttt	3883
tgggcccagtt	ttcattacga	gtaactcaca	ctttttgatt	aaagaacttg	aaattacgtt	3943
atcacttagt	ataattgaca	ttatatagag	actatgtaac	atgcaatcat	tagaatcaaa	4003
attagtactt	tggtcaaaat	atttacaaca	ttcacatact	tgtaaaatat	tcatgtaatt	4063
aactgaattt	aaaaccttca	actattatga	agtgtctgct	tgtacaatcg	ctaatttact	4123
cagtttttag	tagctacaac	tcttcgatac	tatcatcaat	atttgacatc	ttttccaatt	4183
tgtgtatgaa	aagtaaatct	attcctgtag	caactgggga	gtcataatat	aggtcaaaga	4243
catatacctt	gttattataa	tatgtatact	ataataatag	ctgggttatcc	tgagcagggg	4303
aaaagggttat	ttttaggaaa	accacttcaa	atagaaagct	gaagtacttc	taatatactg	4363

```

agggaagtat aatatgtgga acaaactctc aacaaaatgt ttattgatgt tgatgaaaca 4423
gatcagtttt tccatccgga ttattattgg ttcatgattt tatatgtgaa tatgtaagat 4483
atgttctgca attttataaa tgttcatgtc tttttttaa aaaggtgcta ttgaaattct 4543
gtgtctccag caggcaagaa tacttgacta actcttttg tctctttatg gtattttcag 4603
aataaagtct gacttgtgtt tttgagatta ttggtgcctc attaatcag caataaagga 4663
aaatatgcat ctcaaaaat 4682
<210> 118
<211> 4558
<212> DNA
<213> Homo sapiens
<220>
<221> misc_feature
<222> 31..33
<223> ATG
<221> misc_feature
<222> 235..237
<223> TGA
<221> polyA_signal
<222> 4531..4536
<223> AATAAA
<400> 118
ctgctgtccc tgggtgtcca cacgtactcc atg cgc tac ctg ctg ccc agc gtc 54
Met Arg Tyr Leu Leu Pro Ser Val
1 5
gtg ctc ctg ggc acg gcg ccc acc tac gtg ttg gcc tgg ggg gtc tgg 102
Val Leu Leu Gly Thr Ala Pro Thr Tyr Val Leu Ala Trp Gly Val Trp
10 15 20
cgg ctg ctc tcc gcc ttc ctg ccc gcc cgc ttc tac caa gcg ctg gac 150
Arg Leu Leu Ser Ala Phe Leu Pro Ala Arg Phe Tyr Gln Ala Leu Asp
25 30 35 40
gac cgg ctg tac tgc gtc tac cag agc atg gtg ctc ttc ttc ttc gag 198
Asp Arg Leu Tyr Cys Val Tyr Gln Ser Met Val Leu Phe Phe Phe Glu
45 50 55
aat tac acc ggg gtc cag gat gct tat aga att tta tga gtcaccagat 247
Asn Tyr Thr Gly Val Gln Asp Ala Tyr Arg Ile Leu *
60 65
ccagaaagaa gaaaaagatt tcctgggaaa agtggttaatt ccaaattaag tatcaagaag 307
actttaccat caatgttgat cttaagtggg ttgactgcag gcatgcttat gaccgatgct 367
ggaaggaagc tgtatgtgaa cacctggata tatggaaccc tacttggtcg cctgtgggtt 427
actattaaag catagacaag tagctgtctc cagacagtgg gatgtgtctac attgtctatt 487
tttggcggct gcacatgaca tcaaattggt tcctgaattt attaaggagt gtaataaag 547
cctgtttgat tgaagattgg ataatagaat ttgtgacgaa agctgatatg caatgggtctt 607
gggcaaacat acctgggtgt acaactttag catcggggct gctggaaggg taaaagctaa 667
atggagtttc tctgtctctg tccatttcct atgaactaat gacaacttga gaaggctggg 727
aggatttgtt attttgcaag tcagatgggt gcatttttga gcattaattt gcagcgtatt 787
tcacttttct tgttattttc aattttattac aacttgacag ctccaagctc ttattactaa 847
agtatttagt atcttgcagc tagttaatat ttcatctttt gcttatttct acaagtcagt 907
gaaataaatt gtatttagga agtgtcagga tgttcaaagg aaagggtaaa aagtgttcac 967
ggggaaaaag ctctgtttag cacatgattt tattgtattg cgttattagc tgattttact 1027
cattttatat ttgcaaaata aattttcta atttattgaa attgcttaac ttgcacaccc 1087
tgtacacaca gaaaatggta taaaatatga gaacgaagtt taaaattgtg actctgatcc 1147
attatagcag aacttttaaa ttcccagctt ttggaagatt taagctacgc tattagtact 1207
tccctttgtc tgtgccataa gtgcttgaaa acgttaagggt tttctgtttt gttttgtttt 1267
tttaatatca aaagagtcgg tgtgaacctt ggttggaacc caagttcaca agatttttaa 1327
ggtgatgaga gcctgcagac attctgccta gatttactag cgtgtgcctt ttgcctgctt 1387
ctctttgatt tcacagaata ttcattcaga agtcgcgttt ctgtagtggt gtggattccc 1447
actgggctct ggtccttccc ttggatcccg tcagtgggtg tgctcagcgg cttgcacgta 1507
gacttgctag gaagaaatgc agagccagcc tagtctgccc actttcagag ttgaactctt 1567
taagcccttg tgagtgggtc tcaccagcta ctgcagaggc attttgcatt tgtctgtgtc 1627
aagaagtcca ccttctcaag ccagtgaat acagacttaa ttcgtcatga ctgaacgaat 1687
ttgtttattt cccattaggt ttagtggagc tacacattaa tatgtatcgc cttagagcaa 1747

```

```

gagctgtgtt ccaggaacca gatcacgatt tttagccatg gaacaatata tcccatggga 1807
gaagaccttt cagtgtgaac tgttctatct ttgtgttata atttaaactt cgatttcctc 1867
atagtccttt aagttgacat ttctgcttac tgctactgga tttttgctgc agaaatatat 1927
cagtggtccca cattaacat accagttgga tcatgataag caaaatgaaa gaaataatga 1987
ttaagggaaa attaagtgac tgtgttacac tgcttctccc atgccagaga ataaactctt 2047
tcaagcatca tctttgaaga gtcgtgtggt gtgaattggt ttgtgtacat tagaatgtat 2107
gcacacatcc atggacactc aggatatagt tggcctaata atcggggcat gggtaaaact 2167
tatgaaaatt tctctcatgt gaattgtaat ttctcttac ctgtaaagta aaatttagat 2227
caattccatg tctttgttaa gtacagggat ttaatatatt ttgaatataa tgggtatggt 2287
ctaaatttga actttgagag gcaatactgt tgggaattatg tggattctaa ctcatcttaa 2347
caaggtagcc tgacctgcat aagatcactt gaatgttagg ttcatagaa ctatactaact 2407
cttctcacaa aaggtctata aaatacagtc gttgaaaaaa attttgtatc aaaatgtttg 2467
gaaaattaga agcttctcct taacctgtat tgatactgac ttgaattatt ttctaaaatt 2527
aagagccgta acctacctg taagtctttt cacatatcat ttaaaactttt gtttgtatta 2587
ttactgattt acagcttagt tattaatttt tctttataag aatgccgtcg atgtgcatgc 2647
ttttatgttt ttgagaaaag ggtgtgtttg gatgaaagta aaaaaaaaaa taaaatcttt 2707
cactgtctct aatggctgtg ctgtttaaca ttttttgacc ctaaaattca ccaacagtct 2767
cccagtacat aaaataggct taatgactgg cctgcattc ttcacaatat tttccctaa 2827
gctttgagca aagtttttaa aaaatacact aaaataatca aaactgttaa gcagtatat 2887
agtttggtta tataaattca tctgcaattt ataagatgca tggccgatgt taatttgctt 2947
ggcaattctg taatcattaa gtgatctcag tgaacatgt caaatgcctt aaattaacta 3007
agttggtgaa taaaagtgcc gatctggcta actcttacac catacatact gatagttttt 3067
catatgtttc atttccatgt gattttttaa atttagagtg gcaacaattt tgcttaatat 3127
gggttacata agctttattt ttctcttgtt tcataattat attctttgaa taggtctgtg 3187
tcaatcaagt gatctaacta gactgatcat agatagaagg aaataaggcc aagttcaaga 3247
ccagcctggg caacatatcg agaacctgtc taaaaaaaaa taaaaaaaaa ttagccaggc 3307
atggtggcgt aactgagta gtttgccca gctactcggg aggggtgagg gggaggatcg 3367
cttcagccca ggaggttgag attgcagtga gccatggaca taccactgca ctacagccta 3427
ggtaacagca cgagacccca actcttagaa aatgaaaagg aaatatagaa atataaaatt 3487
tgcttattat agacacacag taactccag atatgtacca caaaaaatgt gaaaagagag 3547
agaaatgtct accaaagcag tattttgtgt gtataattgc aagcgcatag taaaataatt 3607
ttaaccttaa tttgttttta gtagtgttta gattgaagat tgagtgaat attttcttgg 3667
cagatattcc gtatctggtg gaaagctaca atgcaatgtc gttgtagttt tgcattggct 3727
gctttataaa caagattttt tctccctcct tttgggccag ttttcattac gagtaactca 3787
cactttttga ttaaagaact tgaaattacg ttatcactta gtataattga cattatatag 3847
agactatgta acatgcaatc attagaatca aaattagtag tttggtcaaa atatttacaa 3907
cattcacata ctgtcaaat attcatgtaa ttaactgaat ttaaaacctt caactattat 3967
gaagtgtctg tctgtacaat cgctaattta ctgagtttag agtagctaca actcttcgat 4027
actatcatca atatttgaca tcttttccaa tttgtgtatg aaaagtaaat ctattcctgt 4087
agcaactggg gagtcatata tgaggtcaaa gacatatacc ttgttattat aatatgtata 4147
ctataataat agctggttat cctgagcagg ggaaaagggt atttttagga aaaccacttc 4207
aaatagaaaag ctgaagtact tctaataatc tgagggaagt ataatatgtg gaacaaactc 4267
tcaacaaaat gtttattgat gttgatgaaa cagatcagtt ttccatccg gattattatt 4327
ggttcatgat ttatattgtg aatatgtaag atatgttctg caattttata aatgttcagt 4387
tcttttttta aaaaagggtc tattgaaatt ctgtgtctcc agcaggcaag aatacttgac 4447
taactctttt tgtctcttta tgggtatttc agaataaagt ctgacttgtg tttttgagat 4507
tattggtgcc tcattaattc agcaataaag gaaaatatgc atctcaaaaa t 4558

```

<210> 119  
<211> 5270  
<212> DNA  
<213> Homo sapiens  
<220>  
<221> misc\_feature  
<222> 31..33  
<223> ATG  
<221> misc\_feature  
<222> 229..231  
<223> TAG  
<221> polyA\_signal  
<222> 5243..5248  
<223> AATAAA



&lt;400&gt; 119

ctgctgtccc	tggtgctcca	cacgtactcc	atg	cgc	tac	ctg	ctg	ccc	agc	gtc	54
			Met	Arg	Tyr	Leu	Leu	Pro	Ser	Val	
			1			5					
gtg	ctc	ctg	ggc	acg	gcg	ccc	acc	tac	gtg	ttg	102
Val	Leu	Leu	Gly	Thr	Ala	Pro	Thr	Tyr	Val	Leu	
	10				15				20		
cgg	ctg	ctc	ttc	gcc	ttc	ctg	ccc	gcc	cgc	ttc	150
Arg	Leu	Leu	Ser	Ala	Phe	Leu	Pro	Ala	Arg	Phe	
	25				30				35		
gac	cgg	ctg	tac	tgc	gtc	tac	cag	agc	atg	gtg	198
Asp	Arg	Leu	Tyr	Cys	Val	Tyr	Gln	Ser	Met	Val	
				45					50		
aat	tac	acc	ggg	gtc	cag	aga	ttg	gat	tcg	tag	251
Asn	Tyr	Thr	Gly	Val	Gln	Arg	Leu	Asp	Ser	*	
			60						65		
cataaaagtg	gaaggccctc	tttaacaata	ttgctatatg	gagatttgcc	aaaaataaaa						311
gaaaatataa	tatatattagc	aaatcatcaa	agcacagttg	actggattgt	tgctgacatc						371
ttggccatca	ggcagaatgc	gctaggacat	gtgcgctacg	tgctgaaaga	agggttaaaa						431
tggtctgcat	tgatatgggtg	ttactttgct	cagcatggag	gaatctatgt	aaagcgcagt						491
gccaaattta	acgagaaaga	gatgcgaac	aagttgcaga	gctacgtgga	cgcaggaact						551
ccaatgtatc	ttgtgatttt	tccagaaggt	acaaggtata	atccagagca	aacaaaagtc						611
ctttcagcta	gtcaggcatt	tgctgcccac	cgtggccttg	cagtattaaa	acatgtgcta						671
acaccacgaa	taaaggcaac	tcacgttgct	tttgattgca	tgaagaatta	tttagatgca						731
atttatgatg	ttacggtggt	ttatgaagg	aaagacgatg	gagggcagcg	aagagagtca						791
ccgaccatga	cggattttct	ctgcaaagaa	tgtccaaaaa	ttcatattca	cattgatcgt						851
atcgacaaaa	aagatgtccc	agaagaacaa	gaacatatga	gaagatggct	gcatgaacgt						911
ttcgaaatca	aagataagat	gcttatagaa	ttttatgagt	caccagatcc	agaaagaaga						971
aaaagatttc	ctgggaaaag	tgtaattccc	aaattaagta	tcaagaagac	tttaccatca						1031
atgttgatct	taagtggttt	gactgcaggc	atgcttatga	ccgatgctgg	aaggaagctg						1091
tatgtgaaca	ccggatata	tggaacccta	cttggctgcc	tgtgggttac	tattaaagca						1151
tagacaagta	gctgtctcca	gacagtggga	tgtgctacat	tgtctatttt	tgccggctgc						1211
acatgacatc	aaattgtttc	ctgaatttat	taaggagtgt	aaataaagcc	ttgttgattg						1271
aagattggat	aatagaattt	gtgacgaaag	ctgatatgca	atggctcttg	gcaaacatac						1331
ctgggtgtac	aaatttagca	tcggggctgc	tggaagggtg	aaagctaaat	ggagtctctc						1391
ctgctctgtc	catttcctat	gaactaatga	caacttgaga	aggctgggag	gatttgtgtat						1451
tttgcaagtc	agatggctgc	atttttgagc	attaatttgc	agcgtatttc	actttttctg						1511
ttattttcaa	tttattacaa	cttgacagct	ccaagctctt	attactaaag	tatttagtat						1571
cttgacagta	gttaatatatt	catcttttgc	ttatttctac	aagtcagtga	aataaattgt						1631
atttaggaag	tgtcaggatg	ttcaaaggaa	agggtaaaaa	gtgttcattg	ggaaaaagct						1691
ctgtttagca	catgatttta	ttgtattgct	ttatttagctg	attttactca	ttttatattt						1751
gcaaaataaa	tttctaatat	ttattgaaat	tgcttaattt	gcacaccctg	tacacacaga						1811
aaatggtata	aaatatgaga	acgaagttta	aaattgtgac	tctgattcat	tatagcagaa						1871
ctttaaattt	cccagctttt	tgaagattta	agctacgcta	ttagtacttc	cctttgtctg						1931
tgccataagt	gcttgaaaac	gttaagggtt	tctgttttgt	tttgtttttt	taatatcaaa						1991
agagtcggtg	tgaaccttgg	ttggacccca	agttcacaaag	atttttaagg	tgatgagagc						2051
ctgcagacat	tctgcctaga	tttactagcg	tgtgcctttt	gcctgcttct	ccttgatttc						2111
acagaatatt	cattcagaag	tcgcgtttct	gtagtgtggt	ggattccac	tggtgctctg						2171
tccttccctt	ggatcccgtc	agtgtgtgct	ctcagcggtc	tgacagtaga	cctgctagga						2231
agaaatgcag	agccagcctg	tgctgcccac	tttcagagtt	gaactcttta	agcccttggt						2291
agtgggcttc	accagctact	gcagaggcat	tttgcatttg	tctgtgtcaa	gaagttcacc						2351
ttctcaagcc	agtgaataac	agacttaatt	cgtcatgact	gaacgaattt	gtttattttc						2411
cattagggtt	agtggagcta	cacattaata	tgtatcgctt	tagagcaaga	gctgtgttcc						2471
aggaaccaga	tcacgatttt	tagccatgga	acaatatatc	ccatgggaga	agacctttca						2531
gtgtgaactg	ttctattttt	gtgttataat	ttaaacttcg	atttcctcat	agtcctttaa						2591
gttgacattt	ctgcttactg	ctactggatt	tttctgcag	aaatatatca	gtggccaca						2651
ttaaacatac	cagttggatc	atgataagca	aaatgaaaga	aataatgatt	aagggaaaat						2711
taagtgactg	tggtacactg	cttctcccat	gccagagaat	aaactctttc	aagcatcatc						2771
tttgaagagt	cgtgtggtgt	gaattgggtt	gtgtacatta	gaatgtatgc	acacatccat						2831
ggacactcag	gatatagttg	gcctaataat	cggggcatgg	gtaaaactta	tgaaaatttc						2891
ctcatgctga	attgtaattt	tctcttacct	gtaaagtaaa	atttagatca	attccatgtc						2951

```

tttgtaagt acaggggattt aatataatgt gaatataatg ggtatgttct aaatttgaac 3011
tttgagaggc aatctactgttg gaattatgtg gattctaaact cattttaaca aggtagcctg 3071
acctgcataa gatcacttga atgttagggt tcatagaact atactaatct tctcacaaaa 3131
ggtctataaa atacagtcgt tgaaaaaaat tttgtatcaa aatgttttga aaattagaag 3191
cttctcctta acctgtattg atactgactt gaattatgtt ctaaaattaa gagccgtata 3251
cctacctgta agtcttttca catatcatgt aaacttttgt ttgtattatt actgatttac 3311
agcttagtta ttaatttttc tttataagaa tgccgtcgat gtgcatgctt ttatgttttt 3371
cagaaaaggg tgtgtttgga tgaaagtaaa aaaaaaaata aaatctttca ctgtctctaa 3431
tggtctgtgct gttaaacatt ttttgacctt aaaattcacc aacagtctcc cagtacataa 3491
aataggctta atgactggcc ctgcattctt cacaatattt tccctaagc tttgagcaaa 3551
gttttaaaaa aataactata aataatcaaa actgttaagc agtatattag tttggttata 3611
taaatccatc tgcaatttat aagatgcgat gccgatgtta atttgcttgg caattctgta 3671
atcattaagt gatctcagtg aaacatgtca aatgccttaa attaaactaag ttggtgaata 3731
aaagtgcgga tctggctaac tcttacacca tacatactga tagtttttca tatgtttcat 3791
ttccatgtga tttttaaaat ttagagtggc aacaattttg cttaatatgg gttacataag 3851
ctttattttt tctttgttcc ataattatat tctttgaata ggtctgtgtc aatcaagtga 3911
tctaactaga ctgatcatag atagaaggaa ataaggccaa gttcaagacc agcctgggca 3971
acatatcgag aacctgtcta caaaaaaatt aaaaaaaatt agccaggcat ggtggcgta 4031
actgagtagt ttgtccagc tactcgggag ggtgaggtgg gaggatcgct tcagccagg 4091
aggttgagat tgcagtggc catggacata ccactgcact acagcctagg taacagcacg 4151
agaccccaac tcttagaaaa tgaaaaggaa atatagaaat ataaaatttg cttattatag 4211
acacacagta actcccagat atgtaccaca aaaaatgtga aaagagagag aaatgtctac 4271
caaagcagta ttttgtgtgt ataattgcaa gcgcatagta aaataatttt aaccttaatt 4331
tgtttttagt agtggttaga ttgaagattg agtgaaatat tttcttggca gatattccgt 4391
atctggtgga aagctacaat gcaatgtcgt tgtagttttg catggcttgc tttataaaca 4451
agattttttc tccctccttt tgggccagtt ttcattacga gtaactcaca ctttttgatt 4511
aaagaacttg aaattacgtt atcacttagt ataattgaca ttatatagag actatgtaac 4571
atgcaatcat tagaatcaaa attagtactt tgggtcaaaat atttacaaca ttcacatact 4631
tgtcaaatat tcatgtaatt aactgaattt aaaaccttca actattatga agtgctcgct 4691
tgtacaatcg ctaatttact cagtttagag tagctacaac tcttcgatac tatcatcaat 4751
atgtgacatc ttttccaatt tgtgtatgaa aagtaaatct attcctgtag caactgggga 4811
gtcatatatg aggtcaaaga catatacctt gttattataa tatgtatact ataataatag 4871
ctggttatcc tgagcagggg aaaagggttat ttttaggaaa accacttcaa atagaaagct 4931
gaagtacttc taatatactg aggggaagtat aatatgtgga acaaactctc aacaaaatgt 4991
ttattgatgt tgatgaaaca gatcagtttt tccatccgga ttattattgg ttcattgatt 5051
tatatgtgaa tatgttaagt atgttctgca attttataaa tgttcatgtc tttttttaa 5111
aaaggtgcta ttgaaattct gtgtctccag caggcaagaa tacttgacta actctttttg 5171
tctctttatg gtattttcag aataaagtct gacttgtgtt tttgagatta ttggtgcctc 5231
attaattcag caataaagga aaatatgcat ctcaaaaaa 5270
<210> 120
<211> 5002
<212> DNA
<213> Homo sapiens
<220>
<221> misc_feature
<222> 31..33
<223> ATG
<221> misc_feature
<222> 322..324
<223> TAA
<221> polyA_signal
<222> 4975..4980
<223> AATAAA
<400> 120
ctgctgtccc tgggtgctcca cacgtactcc atg cgc tac ctg ctg ccc agc gtc 54
Met Arg Tyr Leu Leu Pro Ser Val
1 5
gtg etc ctg gcc acg gcg ccc acc tac gtg ttg gcc tgg ggg gtc tgg 102
Val Leu Leu Gly Thr Ala Pro Thr Tyr Val Leu Ala Trp Gly Val Trp
10 15 20
cgg ctg etc tcc gcc ttc ctg ccc gcc cgc ttc tac caa gcg ctg gac 150

```



```

ggctgtgctg tttaacattt tttgacccta aaattcacca acagtctccc agtacataaa 3224
ataggcttaa tgactggccc tgcattcttc acaatatattt tccctaagct ttgagcaaaag 3284
ttttaaaaaa atacactaaa ataatacaaaa ctgttaagca gtatattagt ttggttatat 3344
aaattcatct gcaatttata agatgcatgg ccgatgttaa tttgcttggc aattctgtaa 3404
tcattaagtg atctcagtg aacatgtcaa atgccttaaa ttaactaagt tgggtgaataa 3464
aagtgccgat ctggctaact cttacaccat acatactgat agtttttcat atgtttcatt 3524
tccatgtgat ttttaaaatt tagagtggca acaattttgc ttaatatggg ttacataaagc 3584
tttatttttt cttttgttca taattatatt ctttgaatag gtctgtgtca atcaagtgat 3644
ctaactagac tgatcataga tagaaggaaa taaggccaag ttcaagacca gcctgggcaa 3704
catatcgaga acctgtctac aaaaaaatta aaaaaaatta gccaggcatg gtggcgtaaa 3764
ctgagtagtt tgtcccagct actcgggagg gtgagggtggg aggatcgctt cagcccagga 3824
ggttgagatt gcagttagcc atggacatac cactgcacta cagcctaggt aacagcacga 3884
gaccccaact cttagaaaat gaaaaggaaa tatagaaaata taaaatttgc ttattataga 3944
cacacagtaa ctcccagata tgtaccacaa aaaatgtgaa aagagagaga aatgtctacc 4004
aaagcagtat tttgtgtgta taattgcaag cgcatagtaa aataatttta accttaattt 4064
gttttttagta gtgttttagat tgaagattga gtgaaatatt ttcttggcag atattccgta 4124
tctggtggaa agttacaatg caatgtcgtt gtagttttgc atggcttgct ttataaacia 4184
gattttttct ccttcttttt gggccagttt tcattacgag taactcacac tttttgatta 4244
aagaacttga aattacgtta tcacttagta taattgacat tatatagaga ctatgtaaca 4304
tgcaatcatt agaatacaaaa ttagtacttt ggtcaaaaata tttacaacat tcacatactt 4364
gtcaaatatt catgtaatta actgaattta aaaccttcaa ctattatgaa gtgctcgtct 4424
gtacaatcgc taattttactc agtttagagt agtacaact cttcgatact atcatcaata 4484
tttgacatct tttccaattt gtgtatgaaa agtaaatcta ttctgttagc aactggggag 4544
tcataatga ggtcaaagac atataccttg ttattataat atgtatacta taataatagc 4604
tggttatcct gagcagggga aaagggttatt tttaggaaaa ccacttcaaa tagaaagctg 4664
aagtacttct aatatactga gggaagtata atatgtggaa caaactctca acaaaatggt 4724
tattgatggt gatgaaacag atcagttttt ccatccggat tattattggt tcatgatttt 4784
atatgtgaat atgtaagata tgttctgcaa ttttataaat gtatcatgtct ttttttaaaa 4844
aagggtctat tgaaattctg tgtctccagc aggcaagaat acttgactaa ctctttttgt 4904
ctctttatgg tattttcaga ataaagtctg acttgtgttt ttgagattat tgggtgcctca 4964
ttaattcagc aataaaggaa aatatgcacg tcaaaaat 5002

<210> 121
<211> 4958
<212> DNA
<213> Homo sapiens
<220>
<221> misc_feature
<222> 31..33
<223> ATG
<221> misc_feature
<222> 577..579
<223> TGA
<221> polyA_signal
<222> 4931..4936
<223> AATAAA
<400> 121
ctgctgtccc tgggtgtcca cacgtactcc atg cgc tac ctg ctg ccc agc gtc 54
Met Arg Tyr Leu Leu Pro Ser Val
1 5
gtg ctc ctg ggc acg gcg ccc acc tac gtg ttg gcc tgg ggg gtc tgg 102
Val Leu Leu Gly Thr Ala Pro Thr Tyr Val Leu Ala Trp Gly Val Trp
10 15 20
cgg ctg ctc tcc gcc ttc ctg ccc gcc cgc ttc tac caa gcg ctg gac 150
Arg Leu Leu Ser Ala Phe Leu Pro Ala Arg Phe Tyr Gln Ala Leu Asp
25 30 35 40
gac cgg ctg tac tgc gtc tac cag agc atg gtg ctc ttc ttc ttc gag 198
Asp Arg Leu Tyr Cys Val Tyr Gln Ser Met Val Leu Phe Phe Phe Glu
45 50 55
aat tac acc ggg gtc cag ata ttg cta tat gga gat ttg cca aaa aat 246
Asn Tyr Thr Gly Val Gln Ile Leu Leu Tyr Gly Asp Leu Pro Lys Asn
60 65 70

```

aaa gaa aat ata ata tat tta gca aat cat caa agc aca gtt gac tgg	294
Lys Glu Asn Ile Ile Tyr Leu Ala Asn His Gln Ser Thr Val Asp Trp	
75 80 85	
att gtt gct gac atc ttg gcc atc agg cag aat gcg cta gga cat gtg	342
Ile Val Ala Asp Ile Leu Ala Ile Arg Gln Asn Ala Leu Gly His Val	
90 95 100	
cgc tac gtg ctg aaa gaa ggg tta aaa tgg ctg cca ttg tat ggg tgt	390
Arg Tyr Val Leu Lys Glu Gly Leu Lys Trp Leu Pro Leu Tyr Gly Cys	
105 110 115 120	
tac ttt gct cag cat gga gga atc tat gta aag cgc agt gcc aaa ttt	438
Tyr Phe Ala Gln His Gly Gly Ile Tyr Val Lys Arg Ser Ala Lys Phe	
125 130 135	
aac gag aaa gag atg cga aac aag ttg cag agc tac gtg gac gca gga	486
Asn Glu Lys Glu Met Arg Asn Lys Leu Gln Ser Tyr Val Asp Ala Gly	
140 145 150	
act cca aat ttc tct gca aag aat gtc caa aaa ttc ata ttc acg ttg	534
Thr Pro Asn Phe Ser Ala Lys Asn Val Gln Lys Phe Ile Phe Thr Leu	
155 160 165	
atc gta tgc aca aaa aag atg tcc cag aag aac aag aac ata tga	579
Ile Val Ser Thr Lys Lys Met Ser Gln Lys Asn Lys Asn Ile *	
170 175 180	
gaagatggct gcatgaacgt ttcgaaatca aagataagat gcttatagaa ttttatgagt	639
caccagatcc agaaagaaga aaaagatttc ctgggaaaag tgtaattcc aaattaagta	699
tcaagaagac ttaccatca atgttgatct taagtgggtt gactgcaggc atgcttatga	759
ccgatgctgg aaggaagctg tatgtgaaca cctggatata tggaacccta ctggctgcc	819
tgtgggttac tattaagca tagacaagta gctgtctcca gacagtggga tgtgtacat	879
tgtctatattt tggcggctgc acatgacatc aaattgtttc ctgaatttat taaggagtgt	939
aaataaagcc ttgttgattg aagattggat aatagaattt gtgacgaaag ctgatatgca	999
atggctcttg gcaaacatac ctgggtgtac aacttttagca tcggggctgc tggaagggtg	1059
aaagctaaat ggagtttctc ctgctctgtc catttcttat gaactaatga caacttgaga	1119
aggctgggag gattgtgtat ttgtcaagtc agatggctgc atttttgagc attaatgtg	1179
agcgtatttc actttttctg ttattttcaa ttattacaa cttgacagct ccaagctctt	1239
attactaaag tattagtat cttgcagcta gttaatatct catcttttgc ttatttctac	1299
aagtcagtga aataaattgt atttaggaag tgcaggatg ttcaaaggaa agggtaaaaa	1359
gtgttcattg ggaaaaagct ctgttttagca catgatttta ttgtattgag ttattagctg	1419
attttactca ttatatattt gcaaaataaa ttctaatat ttattgaaat tgcttaattt	1479
gcacaccctg tacacacaga aaatgggtata aaatatgaga acgaagtta aaattgtgac	1539
tctgattcat tatagcagaa ctttaaattt cccagctttt tgaagattta agctacgcta	1599
ttagtacttc cctttgtctg tgccataagt gcttgaaaac gtttaagggtt tctgttttgt	1659
tttgtttttt taatatcaaa agagtccgtg tgaaccttgg ttggacccca agttcacaag	1719
atttttaagg tgatgagagc ctgcagacat tctgcctaga ttactagcg tgtgcctttt	1779
gctgcttct ctttgatttc acagaatatt ccttcagaag tcgcgtttct gtagtgtggt	1839
ggattccac tgggctctgg tcttccctt ggatcccgtc agtgggtgctg ctgagcggt	1899
tgcacgtaga cttgctagga agaaatgcag agccagcctg tgctgcccac tttcagagtt	1959
gaactcttta agcccttggt agtgggcttc accagctact gcagaggcat ttgtcatttg	2019
tctgtgtcaa gaagttcacc ttctcaagcc agtgaaatac agacttaatt cgtcatgact	2079
gaacgaattt gtttatttcc cattaggttt agtggagcta cacattaata tgtatcgct	2139
tagagcaaga gctgtgttcc aggaaccaga tcacgatttt tagccatgga acaatatatc	2199
ccatgggaga agacctttca gtgtgaactg ttctattttt gtgttataat ttaaacttcg	2259
atttcctcat agtcctttaa gttgacattt ctgcttactg ctactggatt ttgtctgcag	2319
aaatatatca gtggcccaaca ttaaacatac cagttggatc atgataagca aaatgaaaga	2379
aataatgatt aagggaataa taagtgactg tgttacctg cttctcccat gccagagaat	2439
aaactcttcc aagcatcatc tttgaagagt cgtgtggtgt gaattggttt gtgtacatta	2499
gaatgtatgc acacatccat ggacactcag gatatagttg gcctaataat cggggcatgg	2559
gtaaaactta tgaaaatttc ctcatgctga attgtaattt tctcttacct gtaaaagtaaa	2619
atttagatca attccatgtc tttgttaagt acagggattt aatatatttt gaatataatg	2679
ggtatgttct aaatttgaac tttgagaggc aatactgttg gaattatgtg gattctaact	2739
cattttaaca aggtagcctg acctgcataa gatcacttga atgttaggtt tcatagaact	2799
ataactaatct tctcacaataa ggtctataaa atacagtcgt tgaaaaaaat tttgtatcaa	2859
aatgttttga aaattagaag cttctcctta acctgtattg atactgactt gaattatttt	2919
ctaaaattaa gagccgtata cctacctgta agtcttttca catatcattt aaacttttgt	2979

```

ttgtattatt actgatttac agcttagtta ttaatttttc tttataagaa tgccgtcgat 3039
gtgcatgctt ttatgttttt cagaaaaggg tgtgtttgga tgaaagtaaa aaaaaaata 3099
aaatctttca ctgtctctaa tggctgtggt gtttaacatt ttttgacctt aaaattcacc 3159
aacagctctcc cagtacataa aataggctta atgactggcc ctgcattctt cacaatattt 3219
ttccctaagc ttgagcaaaa gttttaaaaa aatacactaa aataatcaaa actgttaagc 3279
agtatattag ttgggttata taaattcatc tgcaatttat aagatgcatg gccgatgtta 3339
atttgcttgg caattctgta atcattaagt gatctcagtg aaacatgtca aatgccttaa 3399
attaactaag ttgggtgaata aaagtgccga tctgggtaac tcttacacca tacatactga 3459
tagtttttca tatgtttcat ttccatgtga tttttaaaat ttagagtggc aacaattttg 3519
cttaatatgg gttacataag ctttattttt tctttgttc ataattatat tctttgaata 3579
ggctctgtgc aatcaagtga tctaactaga ctgatcatag atagaaggaa ataaggccaa 3639
gttcaagacc agcctgggca acatatcgag aacctgtcta caaaaaatt aaaaaaatt 3699
agccaggcat ggtggcgtag actgagtagt ttgtccagc tactcgggag ggtgaggtgg 3759
gaggatcgct tcagcccagg aggttgagat tgcagtggc catggacata ccactgcact 3819
acagcctagg taacagcacg agaccccaac tcttagaaaa tgaaaaggaa atatagaaat 3879
ataaaatttg cttattatag acacacagta actcccagat atgtaccaca aaaaatgtga 3939
aaagagagag aaatgtctac caaagcagta ttttgtgtgt ataattgcaa gcgcatagta 3999
aaataatttt aaccttaatt tgtttttagt agtgtttaga ttgaagattg agtgaaatat 4059
tttcttgggc gatattccgt atctgggtga aagctacaat gcaatgtcgt tgtagttttg 4119
catggcttgc tttataaaca agattttttc tccctccttt tgggccagtt ttcattacga 4179
gtaactcaca ctttttgatt aaagaacttg aaattacgtt atcacttagt ataattgaca 4239
ttatatagag actatgtaac atgcaatcat tagaatcaaa attagtactt tgggtcaaaat 4299
atttacaaca ttcacatact tgtcaaatat tcatgtaatt aactgaattt aaaaccttca 4359
actattatga agtgcctgct tgtacaatcg ctaatttact cagtttagag tagctacaac 4419
tcttcgatac tatcatcaat atttgacatc ttttccaatt tgtgtatgaa aagtaaatct 4479
attcctgtag caactgggga gtcatatatg aggtcaaaga catatacctt gttattataa 4539
tatgtatact ataataatag ctgggttatcc tgagcagggg aaaaggttat ttttaggaaa 4599
accacttcaa atagaaagct gaagtacttc taatatactg aggggaagtat aatatgtgga 4659
acaaactctc aacaaaatgt ttattgatgt tgatgaaaca gatcagtttt tccatccgga 4719
ttattattgg ttpatgattt tatatgtgaa tatgtaagat atgttctgca attttataaa 4779
tgttcatgtc tttttttaaa aaagggtgcta ttgaaattct gtgtctccag caggcaagaa 4839
tacttgacta actctttttg tctcttttat gtattttcag aataaagtct gacttgtgtt 4899
tttgagatta ttggtgcctc attaatccag caataaagga aaatatgcat ctcaaaaat 4958

<210> 122
<211> 5094
<212> DNA
<213> Homo sapiens
<220>
<221> misc_feature
<222> 31..33
<223> ATG
<221> misc_feature
<222> 976..978
<223> TAG
<221> polyA_signal
<222> 5067..5072
<223> AATAAA
<400> 122
ctgctgtccc tgggtgctcca cacgtactcc atg cgc tac ctg ctg ccc agc gtc 54
Met Arg Tyr Leu Leu Pro Ser Val
1 5
gtg ctc ctg gcc acg gcg ccc acc tac gtg ttg gcc tgg ggg gtc tgg 102
Val Leu Leu Gly Thr Ala Pro Thr Tyr Val Leu Ala Trp Gly Val Trp
10 15 20
cgg ctg ctc tcc gcc ttc ctg ccc gcc cgc ttc tac caa gcg ctg gac 150
Arg Leu Leu Ser Ala Phe Leu Pro Ala Arg Phe Tyr Gln Ala Leu Asp
25 30 35 40
gac cgg ctg tac tgc gtc tac cag agc atg gtg ctc ttc ttc ttc gag 198
Asp Arg Leu Tyr Cys Val Tyr Gln Ser Met Val Leu Phe Phe Phe Glu
45 50 55
aat tac acc ggg gtc cag ata ttg cta tat gga gat ttg cca aaa aat 246

```

Asn	Tyr	Thr	Gly	Val	Gln	Ile	Leu	Leu	Tyr	Gly	Asp	Leu	Pro	Lys	Asn	
			60					65					70			
aaa	gaa	aat	ata	ata	tat	tta	gca	aat	cat	caa	agc	aca	gtt	gac	tgg	294
Lys	Glu	Asn	Ile	Ile	Tyr	Leu	Ala	Asn	His	Gln	Ser	Thr	Val	Asp	Trp	
			75				80					85				
att	gtt	gct	gac	atc	ttg	gcc	atc	agg	cag	aat	gcg	cta	gga	cat	gtg	342
Ile	Val	Ala	Asp	Ile	Leu	Ala	Ile	Arg	Gln	Asn	Ala	Leu	Gly	His	Val	
			90				95				100					
cgc	tac	gtg	ctg	aaa	gaa	ggg	tta	aaa	tgg	ctg	cca	ttg	tat	ggg	tgt	390
Arg	Tyr	Val	Leu	Lys	Glu	Gly	Leu	Lys	Trp	Leu	Pro	Leu	Tyr	Gly	Cys	
					110					115					120	
tac	ttt	gct	cag	cat	gga	gga	atc	tat	gta	aag	cgc	agt	gcc	aaa	ttt	438
Tyr	Phe	Ala	Gln	His	Gly	Gly	Ile	Tyr	Val	Lys	Arg	Ser	Ala	Lys	Phe	
				125					130					135		
aac	gag	aaa	gag	atg	cga	aac	aag	ttg	cag	agc	tac	gtg	gac	gca	gga	486
Asn	Glu	Lys	Glu	Met	Arg	Asn	Lys	Leu	Gln	Ser	Tyr	Val	Asp	Ala	Gly	
			140					145					150			
act	cca	atg	tat	ctt	gtg	att	ttt	cca	gaa	ggg	aca	agg	tat	aat	cca	534
Thr	Pro	Met	Tyr	Leu	Val	Ile	Phe	Pro	Glu	Gly	Thr	Arg	Tyr	Asn	Pro	
			155				160					165				
gag	caa	aca	aaa	gtc	ctt	tca	gct	agt	cag	gca	ttt	gct	gcc	caa	cgt	582
Glu	Gln	Thr	Lys	Val	Leu	Ser	Ala	Ser	Gln	Ala	Phe	Ala	Ala	Gln	Arg	
			170				175				180					
ggg	aaa	gac	gat	gga	ggg	cag	cga	aga	gag	tca	ccg	acc	atg	acg	gaa	630
Gly	Lys	Asp	Asp	Gly	Gln	Arg	Arg	Glu	Ser	Pro	Thr	Met	Thr	Thr	Glu	
				190					195						200	
ttt	ctc	tgc	aaa	gaa	tgt	cca	aaa	att	cat	att	cac	att	gat	cgt	atc	678
Phe	Leu	Cys	Lys	Glu	Cys	Pro	Lys	Ile	His	Ile	His	Ile	Asp	Arg	Ile	
				205					210					215		
gac	aaa	aaa	gat	gtc	cca	gaa	gaa	caa	gaa	cat	atg	aga	aga	tgg	ctg	726
Asp	Lys	Lys	Asp	Val	Pro	Glu	Glu	Gln	Glu	His	Met	Arg	Arg	Trp	Leu	
			220					225					230			
cat	gaa	cgt	ttc	gaa	atc	aaa	gat	aag	atg	ctt	ata	gaa	ttt	tat	gag	774
His	Glu	Arg	Phe	Glu	Ile	Lys	Asp	Lys	Met	Leu	Ile	Glu	Phe	Tyr	Glu	
			235				240					245				
tca	cca	gat	cca	gaa	aga	aga	aga	aga	ttt	cct	ggg	aaa	agt	gtt	aat	822
Ser	Pro	Asp	Pro	Glu	Arg	Arg	Lys	Arg	Phe	Pro	Gly	Lys	Ser	Val	Asn	
			250				255				260					
tcc	aaa	tta	agt	atc	aag	aag	act	tta	cca	tca	atg	ttg	atc	tta	agt	870
Ser	Lys	Leu	Ser	Ile	Lys	Lys	Thr	Leu	Pro	Ser	Met	Leu	Ile	Leu	Ser	
			265			270				275					280	
ggg	ttg	act	gca	ggc	atg	ctt	atg	acc	gat	gct	gga	agg	aag	ctg	tat	918
Gly	Leu	Thr	Ala	Gly	Met	Leu	Met	Thr	Asp	Ala	Gly	Arg	Lys	Leu	Tyr	
				285					290					295		
gtg	aac	acc	tgg	ata	tat	gga	acc	cta	ctt	ggc	tgc	ctg	tgg	gtt	act	966
Val	Asn	Thr	Trp	Ile	Tyr	Gly	Thr	Leu	Leu	Gly	Cys	Leu	Trp	Val	Thr	
			300					305				310				
att	aaa	gca	tag	acaagtagct	gtctccagac	agtgaggatgt	gctacattgt									1018
Ile	Lys	Ala	*													
			315													
ctat	tttt	ttg	cggt	gcaca	tgacat	caaaa	ttgt	ttcctg	aatt	tatt	taa	ggag	tgtaaa			1078
taa	agc	ctt	tg	att	gaag	att	ggataat	aga	att	ttgtg	acgaa	agctg	atat	gcaatg		1138
gtc	ttggg	gca	aacata	cctg	gtt	gtacaac	tttag	catcg	ggg	ctgctg	g	aagg	gtaaaa			1198
gcta	aat	gga	gttt	ctcctg	ctct	gtccat	ttc	ctat	gaa	cta	atgacaa	cttg	agaagg			1258
ctg	ggg	aggat	tgt	gtat	ttt	gcaagtcaga	tggt	ctgcatt	ttt	gagcatt	aatt	ttgcagc				1318
gtat	ttt	cact	tttt	ctgtta	tttt	caattt	att	acaact	gac	agctcca	agct	cttatt				1378
act	aa	agtat	ttag	tatctt	gcag	tagtt	aat	atttcat	cttt	tgctta	ttt	ctacaag				1438
tcag	tgaaat	aaatt	gtatt	tag	gaagtg	g	caggat	gttc	aaag	gaaagg	gtaaaa	aagt				1498
ttcat	gggga	aaa	agctctg	tttag	cacat	gatt	tttattg	tatt	gcgtta	ttag	ctgatt					1558
ttact	cattt	tat	atttgca	aaataa	attt	cta	atattta	ttgaa	attgc	tta	atttgca					1618
cacct	gtac	acacaga	aaaa	tggt	ataaaa	tat	gagaacg	aag	tttaaaa	ttgt	gactct					1678

gattcattat	agcagaactt	taaatttccc	agctttttga	agattttaagc	tacgctatta	1738
gtacttccct	ttgtctgtgc	cataagtgtc	tgaaaacggt	aagggtttct	gttttgtttt	1798
gtttttttta	tatcaaaaga	gtcgggtgtg	accttgggtg	gaccccaagt	tcacaagatt	1858
tttaagggtga	tgagagcctg	cagacattct	gcctagattt	actagcgtgt	gccttttgcc	1918
tgcttctctt	tgatttcaca	gaatattcat	tcagaagtcg	cgtttctgta	gtgtgggtga	1978
ttcccactgg	gctctggtcc	ttcccttgga	tcccgtcagt	ggtgctgctc	agcggcttgc	2038
acgtagactt	gctaggaaga	aatgcagagc	cagcctgtgc	tgcccacttt	cagagttgaa	2098
ctctttaagc	ccttgtgagt	gggcttcacc	agctactgca	gaggcatttt	gcatttgtct	2158
gtgtcaagaa	gttcaccttc	tcaagccagt	gaaatacaga	cttaattcgt	catgactgaa	2218
cgaattttgt	tatttcccat	taggtttagt	ggagctacac	attaatatgt	atcgcttag	2278
agcaagagct	gtgttccagg	aaccagatca	cgatttttag	ccatggaaca	atatatccca	2338
tgggagaaga	cctttcagtg	tgaactgttc	tatttttgtg	ttataattta	aacttcgatt	2398
tcctcatagt	cctttaagtt	gacatttctg	cttactgcta	ctggattttt	gctgcagaaa	2458
tatatcagtg	gcccacatta	aacataccag	ttggatcatg	ataagcaaaa	tgaagaaaat	2518
aatgattaag	ggaaaattaa	gtgactgtgt	tacactgctt	ctcccatgcc	agagaataaa	2578
ctctttcaag	catcatcttt	gaagagtcgt	gtgggtgtgaa	ttgggttgtg	tacattagaa	2638
tgtatgcaca	catccatgga	cactcaggat	atagttggcc	taataatcgg	ggcatgggta	2698
aaacttatga	aaatttcctc	atgctgaatt	gtaattttct	cttacctgta	aagtaaaatt	2758
tagatcaatt	ccatgtcttt	gttaagtaca	gggattttaat	atattttgaa	tataatgggt	2818
atgttctaaa	tttgaacttt	gagaggcaat	actgttggaa	ttatgtggat	tctaactcat	2878
tttaacaagg	tagcctgacc	tgcataagat	cacttgaatg	ttaggtttca	tagaactata	2938
ctaactctct	cacaaaaggt	ctataaaata	cagtcggtga	aaaaaatttt	gtatcaaaaat	2998
gtttggaaaa	ttagaagcct	ctccttaacc	tgtattgata	ctgacttgaa	ttattttcta	3058
aaattaagag	ccgtatacct	acctgtaagt	cttttcacat	atcattttaa	cttttgtttg	3118
tattattact	gatttacagc	ttagttatta	atttttcttt	ataagaatgc	cgctcagtgtg	3178
catgctttta	tggttttcag	aaaagggtgt	gtttggatga	aagtaaaaaa	aaaaataaaa	3238
tctttcactg	tctctaattg	ctgtgctgtt	taacattttt	tgaccctaaa	attcaccaac	3298
agtctcccag	tacataaaaat	aggcttaatg	actggccctg	cattcttcac	aatatttttc	3358
cctaagcttt	gagcaaagtt	ttaaaaaaat	acactaaaaat	aatcaaaaact	gttaagcagt	3418
atatttagttt	ggttatataa	attcatctgc	attttataag	atgcattggcc	gatgttaatt	3478
tgcttggcaa	ttctgtaatc	attaagtgat	ctcagtgaat	catgtcaaat	gccttaaat	3538
aactaagttg	gtgaataaaa	gtgccgatct	ggctaactct	tacaccatac	atactgatag	3598
tttttcatat	gtttcatttc	catgtgattt	ttaaaaattt	gagtggaac	aattttgctt	3658
aatatgggtt	acataagcct	tattttttcc	tttgttcata	attatattct	ttgaataggt	3718
ctgtgtcaat	caagtgatct	aactagactg	atcatagata	gaaggaaata	aggccaagtt	3778
caagaccagc	ctgggcaaca	tatcgagaac	ctgtctacaa	aaaaatttaa	aaaaattagc	3838
caggcatggt	ggcgtacact	gagtagtttg	tcccagctac	tcggggagggt	gaggtgggag	3898
gatcgcttca	gpcaggagg	ttgagattgc	agttagccat	ggacatacca	ctgcaactaca	3958
gcctaggtaa	cagcacgaga	ccccaaactc	tagaaaatga	aaaggaaata	tagaaatata	4018
aaatttgctt	attatagaca	cacagtaact	cccagatatg	taccacaaaa	aatgtgaaaa	4078
gagagagaaa	tgctaccaa	agcagtattt	tgtgtgtata	attgcaagcg	catagtaaaa	4138
taatttttaac	cttaatttgt	tttttagtagt	gttttagattg	aagattgagt	gaaatatttt	4198
cttggcagat	atcccgatc	tggtggaaag	ctacaatgca	atgtcgttgt	agttttgcat	4258
ggcttgcttt	ataaacaaga	ttttttctcc	ctccttttgg	gccagttttc	attacgagta	4318
actcacactt	tttgattaaa	gaacttgaaa	ttacgttatc	acttagtata	attgacatta	4378
tatagagact	atgtaacatg	caatcattag	aatcaaaatt	agtacttttg	tcaaaatatt	4438
tacaacattc	acatacttgt	caaataattca	tgtaattaac	tgaattttaa	accttcaact	4498
attatgaagt	gctcgtctgt	acaatcgcta	atttactcag	tttagagtag	ctacaactct	4558
tcgatactat	catcaatatt	tgacatcttt	tccaatttgt	gtatgaaaag	taaatctatt	4618
cctgtagcaa	ctggggagtc	atatatgagg	tcaaagacat	ataccttgtt	attataatat	4678
gtatactata	ataatagctg	gttatcctga	gcaggggaaa	aggttatttt	taggaaaacc	4738
acttcaaata	gaaagctgaa	gtacttctaa	tatactgagg	gaagtataat	atgtggaaca	4798
aactctcaac	aaaatgttta	ttgatgttga	tgaacacagat	cagtttttcc	atccggatta	4858
ttattgggtc	atgattttat	atgtgaatat	gtaagatatg	ttctgcaatt	ttataaatgt	4918
tcatgtcttt	ttttaaaaaa	ggtgctattg	aaattctgtg	tctccagcag	gcaagaatac	4978
ttgactaact	ctttttgtct	ctttatggta	ttttcagaat	aaagtcctgac	ttgtgttttt	5038
gagattattg	gtgcctcatt	aattcagcaa	taaaggaaaa	tatgcatctc	aaaaat	5094

<210> 123  
 <211> 5049  
 <212> DNA  
 <213> Homo sapiens



```

<220>
<221> misc_feature
<222> 31..33
<223> ATG
<221> misc_feature
<222> 931..933
<223> TAG
<221> polyA_signal
<222> 5022..5027
<223> AATAAA
<400> 123
ctgctgtccc tgggtgtcca cacgtactcc atg cgc tac ctg ctg ccc agc gtc      54
                               1      5
                               Met Arg Tyr Leu Leu Pro Ser Val
gtg ctc ctg ggc acg gcg ccc acc tac gtg ttg gcc tgg ggg gtc tgg      102
Val Leu Leu Gly Thr Ala Pro Thr Tyr Val Leu Ala Trp Gly Val Trp
    10                15                20
cgg ctg ctc tcc gcc ttc ctg ccc gcc cgc ttc tac caa gcg ctg gac      150
Arg Leu Leu Ser Ala Phe Leu Pro Ala Arg Phe Tyr Gln Ala Leu Asp
    25                30                35                40
gac cgg ctg tac tgc gtc tac cag agc atg gtg ctc ttc ttc ttc gag      198
Asp Arg Leu Tyr Cys Val Tyr Gln Ser Met Val Leu Phe Phe Phe Glu
    45                50                55
aat tac acc ggg gtc cag ata ttg cta tat gga gat ttg cca aaa aat      246
Asn Tyr Thr Gly Val Gln Ile Leu Leu Tyr Gly Asp Leu Pro Lys Asn
    60                65                70
aaa gaa aat ata ata tat tta gca aat cat caa agc aca gtt gac tgg      294
Lys Glu Asn Ile Ile Tyr Leu Ala Asn His Gln Ser Thr Val Asp Trp
    75                80                85
att gtt gct gac atc ttg gcc atc agg cag aat gcg cta gga cat gtg      342
Ile Val Ala Asp Ile Leu Ala Ile Arg Gln Asn Ala Leu Gly His Val
    90                95                100
cgc tac gtg ctg aaa gaa ggg tta aaa tgg ctg cca ttg tat ggg tgt      390
Arg Tyr Val Leu Lys Glu Gly Leu Lys Trp Leu Pro Leu Tyr Gly Cys
    105                110                115                120
tac ttt gct cag cat gga gga atc tat gta aag cgc agt gcc aaa ttt      438
Tyr Phe Ala Gln His Gly Gly Ile Tyr Val Lys Arg Ser Ala Lys Phe
    125                130                135
aac gag aaa gag atg cga aac aag ttg cag agc tac gtg gac gca gga      486
Asn Glu Lys Glu Met Arg Asn Lys Leu Gln Ser Tyr Val Asp Ala Gly
    140                145                150
act cca atg tat ctt gtg att ttt cca gaa ggt aca agg tat aat cca      534
Thr Pro Met Tyr Leu Val Ile Phe Pro Glu Gly Thr Arg Tyr Asn Pro
    155                160                165
gag caa aca aaa gtc ctt tca gct agt cag gca ttt gct gcc caa cgt      582
Glu Gln Thr Lys Val Leu Ser Ala Ser Gln Ala Phe Ala Ala Gln Arg
    170                175                180
gaa ttt ctc tgc aaa gaa tgt cca aaa att cat att cac att gat cgt      630
Glu Phe Leu Cys Lys Glu Cys Pro Lys Ile His Ile His Ile Asp Arg
    185                190                195                200
atc gac aaa aaa gat gtc cca gaa gaa caa gaa cat atg aga aga tgg      678
Ile Asp Lys Lys Asp Val Pro Glu Glu Gln Glu His Met Arg Arg Trp
    205                210                215
ctg cat gaa cgt ttc gaa atc aaa gat aag atg ctt ata gaa ttt tat      726
Leu His Glu Arg Phe Glu Ile Lys Asp Lys Met Leu Ile Glu Phe Tyr
    220                225                230
gag tca cca gat cca gaa aga aga aaa aga ttt cct ggg aaa agt gtt      774
Glu Ser Pro Asp Pro Glu Arg Arg Lys Arg Phe Pro Gly Lys Ser Val
    235                240                245
aat tcc aaa tta agt atc aag aag act tta cca tca atg ttg atc tta      822
Asn Ser Lys Leu Ser Ile Lys Lys Thr Leu Pro Ser Met Leu Ile Leu

```

250	255	260	
agt ggt ttg act gca ggc atg ctt atg acc gat gct gga agg aag ctg			870
Ser Gly Leu Thr Ala Gly Met Leu Met Thr Asp Ala Gly Arg Lys Leu			
265	270	275	280
tat gtg aac acc tgg ata tat gga acc cta ctt ggc tgc ctg tgg gtt			918
Tyr Val Asn Thr Trp Ile Tyr Gly Thr Leu Leu Gly Cys Leu Trp Val			
285	290	295	
act att aaa gca tag acaagtagct gtctccagac agtgggatgt gctacattgt			973
Thr Ile Lys Ala *			
300			
ctatTTTTtg cggctgcaca tgacatcaaa ttgtttcctg aatttattaa ggagtgtaaa			1033
taaaagccttg ttgattgaag attggataat agaatttttg acgaaagctg atatgcaatg			1093
gtcttgggca aacatacctg gttgtacaac tttagcatcg gggctgctgg aagggtaaaa			1153
gctaaatgga gtttctcctg ctctgtccat ttccatgaa ctaatgacaa cttgagaagg			1213
ctgggaggat tgtgtatttt gcaagtcaga tggctgcatt tttgagcatt aatttgcagc			1273
gtatttctact ttttctgtta ttttcaattt attacaactt gacagctcca agctcttatt			1333
actaaagtat ttagtatctt gcagctagt t aatatttcat cttttgctta tttctacaag			1393
tcagtgaat aaattgtatt taggaagtgt caggatgttc aaaggaaaagg gtaaaaagt			1453
ttcatgggga aaaagctctg ttttagcacat gattttattg tattgcgtta ttagctgatt			1513
ttactcattt tatatttgca aaataaattt ctaatattta ttgaaattgc ttaatttgca			1573
caccctgtac acacagaaaa tgggtataaaa tatgagaacg aagtttataaa ttgtgactct			1633
gattcattat agcagaactt taaattttccc agcttttttga agatttgaag tacgctatta			1693
gtacttccct ttgtctgtgc cataagtgtc tgaaaacgtt aagggttttct gttttgtttt			1753
gtttttttaa tatcaaaaga gtccgtgtga acctgtgttg gaccccaagt tcacaagatt			1813
tttaaggtag tgagagcctg cagacattct gcctagattt actagcgtgt gccttttgcc			1873
tgcttctctt tgatttcaca gaatattcat tcagaagtcg cgtttctgta gtgtgggtga			1933
ttccactgg gctctgtgctc ttcccttgga tcccgctcag ggtgctgctc agcggcttgc			1993
acgtagactt gctaggaaga aatgcagagc cagcctgtgc tgcccacttt cagagttgaa			2053
ctctttaagc ccttgtgagt gggcttcacc agctactgca gaggcatttt gcatttgtct			2113
gtgtcaagaa gttcaccttc tcaagccagt gaaatacaga ctttaattcgt catgactgaa			2173
cgaatttggt tatttcccat taggtttagt ggagctacac attaatatgt atcgcttag			2233
agcaagagct gtgttccagg aaccagatca cgatttttag ccatggaaca atatatccca			2293
tgggagaaga cctttcagtg tgaactgttc tatttttgtg ttataattta aacttcgatt			2353
tcctcatagt cctttaagtt gacattttctg cttactgcta ctggattttt gctgcagaaa			2413
tatatcagt gcccacatta aacataccag ttggatcatg ataagcaaaa tgaaagaaat			2473
aatgaagtga ggaaaattaa gtgactgtgt tacactgctt ctcccatgcc agagaataaa			2533
ctctttcaag catcatcttt gaagagtcgt gtgggtgtgaa ttgggttgtg tacattagaa			2593
tgatgcaca catccatgga cactcaggat atagttggcc taataatcgg ggcatgggt			2653
aaacttatga aaatttcctc atgctgaatt gtaattttct cttacctgta aagtaaaatt			2713
tagatcaatt ccatgtcttt gttaagtaca gggatttaat atattttgaa tataatgggt			2773
atgttctaaa tttgaacttt gagaggcaat actgttgga ttatgtggat tctaactcat			2833
tttaacaagg tagcctgacc tgcataagat cacttgaatg ttaggtttca tagaactata			2893
ctaattctct cacaaaaggt ctataaaata cagtcgttga aaaaaatttt gtatcaaaat			2953
gtttggaaaa ttagaagctt ctcccttaacc tgtattgata ctgacttgaa ttattttcta			3013
aaattaagag ccgtatacct acctgtgaat cttttcacat atcattttaa cttttgtttg			3073
tattattact gatttacagc ttagttatta atttttcttt ataagaatgc cgtcgatgtg			3133
catgctttta tgtttttcag aaaagggtgt gtttggatga aagtaaaaaa aaaaataaaa			3193
tccttctactg tctctaattg ctgtgctgtt taacattttt tgaccctaaa attcaccaac			3253
agtctcccag tacataaaat aggtctaatt actggccctg cattcttcac aatatttttc			3313
cctaagcttt gagcaaagtt ttaaaaaaat acactaaaa atcaaaaact gttaagcagt			3373
atattagttt ggttatataa attcatctgc aatttataag atgcatggcc gatgttaatt			3433
tgcttggcaa ttctgtaatc attaatgtgat ctgagtgaat catgtcaaat gccttaatt			3493
aactaagttg gtgaataaaa gtgccgatct ggctaactct tacaccatac atactgatag			3553
tttttcatat gtttcatttc catgtgattt ttaaaattta gagtggcaac aattttgctt			3613
aatatgggtt acataagctt tattttttcc tttgttcata attatattct ttgaataggt			3673
ctgtgtcaat caagtgatct aactagactg atcatagata gaaggaaata aggccaagt			3733
caagaccagc ctgggcaaca tatcgagaac ctgtctacaa aaaaattaaa aaaaattagc			3793
caggcatggg ggggtacact gtagtagttt tcccagctac tggggagggt gaggtgggag			3853
gatcgcttca gccaggagg ttgagattgc agtgagccat ggacatacca ctgcactaca			3913
gcctaggtaa cagcacgaga ccccaactct tagaaaatga aaaggaaata tagaaatata			3973
aaatttgctt attatagaca cacagtaact cccagatatg taccacaaaa aatgtgaaaa			4033

gagagagaaa tgtctaccaa agcagtatTT tgtgtgtata attgcaagcg catagtaaaaa 4093  
 taattttaac cttaatttTgt ttttagtagt gtttagattg aagattgagt gaaatatttt 4153  
 cttggcagat attccgtatc tgggtgaaag ctacaatgca atgtcgtTgt agttttgcat 4213  
 ggcttgcttt ataaacaaga ttttttctcc ctcttttTgg gccagttttc attacgagta 4273  
 actcacactt tttgattaaa gaacttgaaa ttacgttatc acttagtata attgacatta 4333  
 tatagagact atgtaacatg caatcattag aatcaaaatt agtactttTg tcaaaatatt 4393  
 tacaacattc acatacttTgt caaatattca tgtaattaac tgaatttaaa accttcaact 4453  
 attatgaagt gctcgtctTgt acaatcgcta atttactcag ttttagagtag ctacaactct 4513  
 tcgatactat catcaatatt tgacatcttt tccaattTgt gtatgaaaag taaatctatt 4573  
 cctgtagcaa ctggggagtc atatatgagg tcaaagacat atacctTgt attataatat 4633  
 gtatactata ataatagctg gttatcctga gcaggggaaa aggttatTTt taggaaaacc 4693  
 acttcaaata gaaagctgaa gtacttctaa tatactgagg gaagtataat atgtggaaca 4753  
 aactctcaac aaaatgttta ttgatgtTga tgaaacagat cagtttttcc atccggatta 4813  
 ttattggTtc atgattttat atgtgaatat gtaagatatg ttctgcaatt ttataaatgt 4873  
 tcatgtcttt ttttaaaaaa ggtgctattg aaattctTg tctccagcag gcaagaatac 4933  
 ttgactaact ctttttTgt ctttatggta ttttcagaat aaagtctgac ttgtgttttt 4993  
 gagattattg gtgcctcatt aattcagcaa taaaggaaaa tatgcatctc aaaaat 5049  
 <210> 124  
 <211> 5324  
 <212> DNA  
 <213> Homo sapiens  
 <220>  
 <221> misc\_feature  
 <222> 31..33  
 <223> ATG  
 <221> misc\_feature  
 <222> 586..588  
 <223> TAA  
 <221> polyA\_signal  
 <222> 5297..5302  
 <223> AATAAA  
 <400> 124  
 ctgctgtccc tgggtgtcca cacgtactcc atg cgc tac ctg ctg ccc agc gtc 54  
 Met Arg Tyr Leu Leu Pro Ser Val  
 1 5  
 gtg ctc ctg ggc acg gcg ccc acc tac gtg ttg gcc tgg ggg gtc tgg 102  
 Val Leu Leu Gly Thr Ala Pro Thr Tyr Val Leu Ala Trp Gly Val Trp  
 10 15 20  
 cgg ctg ctc tcc gcc ttc ctg ccc gcc cgc ttc tac caa gcg ctg gac 150  
 Arg Leu Leu Ser Ala Phe Leu Pro Ala Arg Phe Tyr Gln Ala Leu Asp  
 25 30 35 40  
 gac cgg ctg tac tgc gtc tac cag agc atg gtg ctc ttc ttc ttc gag 198  
 Asp Arg Leu Tyr Cys Val Tyr Gln Ser Met Val Leu Phe Phe Phe Glu  
 45 50 55  
 aat tac acc ggg gtc cag ata ttg cta tat gga gat ttg cca aaa aat 246  
 Asn Tyr Thr Gly Val Gln Ile Leu Leu Tyr Gly Asp Leu Pro Lys Asn  
 60 65 70  
 aaa gaa aat ata ata tat tta gca aat cat caa agc aca gtt gac tgg 294  
 Lys Glu Asn Ile Ile Tyr Leu Ala Asn His Gln Ser Thr Val Asp Trp  
 75 80 85  
 att gtt gct gac atc ttg gcc atc agg cag aat gcg cta gga cat gtg 342  
 Ile Val Ala Asp Ile Leu Ala Ile Arg Gln Asn Ala Leu Gly His Val  
 90 95 100  
 cgc tac gtg ctg aaa gaa ggg tta aaa tgg ctg cca ttg tat ggg tgt 390  
 Arg Tyr Val Leu Lys Glu Gly Leu Lys Trp Leu Pro Leu Tyr Gly Cys  
 105 110 115 120  
 tac ttt gct cag cat gga gga atc tat gta aag cgc agt gcc aaa ttt 438  
 Tyr Phe Ala Gln His Gly Gly Ile Tyr Val Lys Arg Ser Ala Lys Phe  
 125 130 135  
 aac gag aaa gag atg cga aac aag ttg cag agc tac gtg gac gca gga 486  
 Asn Glu Lys Glu Met Arg Asn Lys Leu Gln Ser Tyr Val Asp Ala Gly

	140	145	150	
act cca atg tat ctt gtg att ttt cca gaa ggt aca agg tat aat cca				534
Thr Pro Met Tyr Leu Val Ile Phe Pro Glu Gly Thr Arg Tyr Asn Pro				
	155	160	165	
gag caa aca aaa gtc ctt tca gct agt cag gca ttt gct gcc caa cgt				582
Glu Gln Thr Lys Val Leu Ser Ala Ser Gln Ala Phe Ala Ala Gln Arg				
	170	175	180	
ggc taa agcagtcctc ctgagtagtt aggactacag acatacacgt gccaccgcgc				638
Gly *				
185				
ccagctccgt gttctctttg tttccctgcc tcctgctctt ccacttatct ttgcatggca				698
ggccttgtag tattaaaaca tgtgctaaca ccacgaataa aggcaactca cgttgctttt				758
gattgcatga agaattatct agatgcaatt tatgatgtta cgggtggtta tgaaggga				818
gacgatggag ggcagcgaag agagtcaccg accatgacgg aatttctctg caaagaatgt				878
ccaaaaattc atattcacat tgatcgtatc gacaaaaaag atgtcccaga agaacaagaa				938
catatgagaa gatggctgca tgaacgtttc gaaatcaaag ataagatgct tatagaatct				998
tatgagtcac cagatccaga aagaagaaaa agatttcctg ggaaaagtgt taattccaaa				1058
ttaagtatca agaagacttt accatcaatg ttgatcttaa gtggtttgac tgcaggcatg				1118
cttatgaccg atgctggaag gaagctgtat gtgaacacct ggatatatgg aacctactt				1178
ggctgcctgt gggttactat taaagcatag acaagtagct gtctccagac agtgggatgt				1238
gctacattgt ctatttttgg cggtgcaca tgacatcaaa ttgtttcctg aattttattaa				1298
ggagtgtaaa taaagccttg ttgattgaag attggataat agaatttgtg acgaaagctg				1358
atatgcaatg gtcttgggca aacataacctg gttgtacaac tttagcatcg gggctgctg				1418
aagggtaaaa gctaaatgga gtttctcctg ctctgtccat ttcctatgaa ctaatgacaa				1478
cttgagaagg ctgggaggat tgtgtatttt gcaagtcaga tggctgcatt ttgagcatt				1538
aatttgcagc gtatttcact ttttctgtta ttttcaattt attacaactt gacagctcca				1598
agctcttatt actaaagtat ttagtatctt gcagctagtt aatatttcat cttttgctta				1658
tttctacaag tcagtgaat aaattgtatt taggaagtgt caggatgttc aaaggaaagg				1718
gtaaaaagtg ttcatgggga aaaagctctg tttagcacat gattttattg tattgcgtta				1778
ttagctgatt ttactcattt tatatttgca aaataaattt ctaatatatta ttgaaattgc				1838
ttaatttgca caccctgtac acacagaaaa ttgtataaaa tatgagaacg aagtttaaaa				1898
ttgtgactct gattcattat agcagaactt taaatttccc agctttttga agatttaagc				1958
tacgctatta gtacttccct ttgtctgtgc cataagtgct tgaaaacggt aaggttttct				2018
gttttgtttt gtttttttaa tatcaaaaga gtcggtgtga accttggttg gaccccaagt				2078
tcacaagatt ttttaaggta tgagagcctg cagacattct gcctagattt actagcgtgt				2138
gccttttgcc tgcttctctt tgatttcaca gaatttccat tcagaagtcg cgtttctgta				2198
gtgtggtgga ttcccaactgg gctctggtcc ttcccttgga tcccgtcagt ggtgctgctc				2258
agcggcttgc acgtagactt gctaggaaga aatgcagagc cagcctgtgc tgcccaactt				2318
cagagttgaa ctctttaagc cctgtgaggt gggcttcacc agctactgca gaggcatttt				2378
gcatttgtct gtgtcaagaa gttcaccttc tcaaggcagt gaaatacaga cttaattcgt				2438
catgactgaa cgaatttgtt tatttcccat taggtttagt ggagctacac attaatatgt				2498
atcgcttag agpaagagct gtgttcagg aaccagatca cgatttttag ccatggaaca				2558
atatatccca tgggagaaga cctttcagtg tgaactgttc tatttttggt ttataattta				2618
aacttcgatt tcctcatagt cctttaagtt gacatttctg cttactgcta ctggattttt				2678
gctgcagaaa tatatcagtg gccacatta aacataccag ttggatcatg ataagcaaaa				2738
tgaagaaaaa aatgattaag ggaaaattaa gtgactgtgt tacactgctt ctcccagcc				2798
agagaataaa ctctttcaag catcatcttt gaagagtcgt gtggtgtgaa ttggtttgtg				2858
tacattagaa tgtatgcaca catccatgga cactcaggat atagtggcc taataatcgg				2918
ggcatgggta aaacttatga aaatttcctc atgctgaatt gtaattttct cttacctgta				2978
aagtaaaatt tagatcaatt ccatgtcttt gttaagtaca gggatttaaat atattttgaa				3038
tataatgggt atgttctaaa tttgaacttt gagaggcaat actgttgga ttatgtggat				3098
tctaactcat tttacaagg tagcctgacc tgcataagat cacttgaatg ttaggtttca				3158
tagaactata ctaatcttct cacaaaagg ctataaaata cagtcgttga aaaaaatttt				3218
gtatcaaaat gtttggaataa ttagaagctt ctcttaacc tgtattgata ctgacttgaa				3278
ttattttcta aaattaagag cgtataacct acctgtaagt cttttcacat atcatttaaa				3338
cttttgtttg tattattact gatttacagc tagttatta atttttctt ataagaatgc				3398
cgctgatgtg catgctttta tgtttttcag aaaagggtgt gtttgatga aagtaaaaaa				3458
aaaaataaaa tctttcactg tctctaattg ctgtgctgt taacattttt tgacctaaa				3518
attcaccaac agtctcccag tacataaaat aggtctaatt actggccctg cattcttcac				3578
aatatttttc cctaagcttt gagcaaagtt ttaaaaaaat aactaaaaat aatcaaaact				3638
gtaagcagt atattagttt ggttatataa attcatctgc aatttataag atgcatggcc				3698

```

gatgttaatt tgcttggcaa ttctgtaatc attaatgat ctcagtgaac catgtcaaat 3758
gccttaaatt aactaagttg gtgaataaaa gtgccgatct ggctaactct tacaccatac 3818
atactgatag tttttcatat gtttcatttc catgtgattt ttaaaattta gagtggcaac 3878
aattttgctt aatatgggtt acataagctt tattttttcc tttgttcata attatatctt 3938
ttgaatagggt ctgtgtcaat caagtgatct aactagactg atcatagata gaaggaaata 3998
aggccaagtt capgaccagc ctgggcaaca tatcgagaac ctgtctacaa aaaaattaaa 4058
aaaaattagc caggcatggt ggcgtacact gagtagtttg tcccagctac tcggggaggt 4118
gaggtgggag gatcgcttca gccaggaggg ttgagattgc agtgagccat ggacatacca 4178
ctgcactaca gcctaggtaa cagcacgaga cccaactct tagaaaatga aaaggaaata 4238
tagaaatata aaatttgctt attatagaca cacagtaact cccagatatg taccacaaaa 4298
aatgtgaaaa gagagagaaa tgtctaccaa agcagtattt tgtgtgtata attgcaagcg 4358
catagtaaaa taattttaac cttaatttgt ttttagtagt gtttagattg aagattgagt 4418
gaaatatttt ctgggcagat attccgtatc tgggtgaaaag ctacaatgca atgtcgttgt 4478
agttttgcat ggcttgcttt ataaacaaga ttttttctcc ctcttttgg gccagttttc 4538
attacgagta actcacactt tttgattaaa gaacttgaaa ttacgttatc acttagtata 4598
attgacatta tatagagact atgtaacatg caatcattag aatcaaaatt agtactttgg 4658
tcaaaatatt tacaacattc acatacttgt caaatattca tgtaattaac tgaatttaaa 4718
accttcaact attatgaagt gctcgtctgt acaatcgcta atttactcag tttagagtag 4778
ctacaactct tcgatactat catcaatatt tgacatcttt tccaatttgt gtatgaaaag 4838
taaacttatt cctgtagcaa ctggggagtc atatatgagg tcaaagacat ataccttgtt 4898
attataatat gtatactata ataatagctg gttatcctga gcaggggaaa aggttatatt 4958
taggaaaacc acttcaataa gaaagctgaa gtacttctaa tatactgagg gaagtataat 5018
atgtggaaca aactctcaac aaaatgttta ttgatgttga tgaaacagat cagtttttcc 5078
atccggatta ttattgggtc atgattttat atgtgaatat gtaagatatg ttctgcaatt 5138
ttataaatgt tcatgtcttt ttttaaaaaa ggtgctattg aaattctgtg tctccagcag 5198
gcaagaatac ttgactaact ctttttgtct ctttatggta ttttcagaat aaagtctgac 5258
ttgtgttttt gaqattattg gtgcctcatt aattcagcaa taaaggaaaa tatgcatctc 5318
aaaaat 5324

```

&lt;210&gt; 125

&lt;211&gt; 77

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 125

```

Met Arg Tyr Leu Leu Pro Ser Val Val Leu Leu Gly Thr Ala Pro Thr
1          5          10          15
Tyr Val Leu Ala Trp Gly Val Trp Arg Leu Leu Ser Ala Phe Leu Pro
20          25          30
Ala Arg Phe Tyr Gln Ala Leu Asp Asp Arg Leu Tyr Cys Val Tyr Gln
35          40          45
Ser Met Val Leu Phe Phe Phe Glu Asn Tyr Thr Gly Val Gln Leu Thr
50          55          60
Gly Leu Leu Leu Thr Ser Trp Pro Ser Gly Arg Met Arg
65          70          75

```

&lt;210&gt; 126

&lt;211&gt; 238

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; 98..103

&lt;223&gt; Box II

&lt;400&gt; 126

```

Met Arg Tyr Leu Leu Pro Ser Val Val Leu Leu Gly Thr Ala Pro Thr
1          5          10          15
Tyr Val Leu Ala Trp Gly Val Trp Arg Leu Leu Ser Ala Phe Leu Pro
20          25          30
Ala Arg Phe Tyr Gln Ala Leu Asp Asp Arg Leu Tyr Cys Val Tyr Gln
35          40          45
Ser Met Val Leu Phe Phe Phe Glu Asn Tyr Thr Gly Val Gln His Gly
50          55          60
Gly Ile Tyr Val Lys Arg Ser Ala Lys Phe Asn Glu Lys Glu Met Arg

```



Ser Pro Asp Pro Glu Arg Arg Lys Arg Phe Pro Gly Lys Ser Val Asn  
 225 230 235 240  
 Ser Lys Leu Ser Ile Lys Lys Thr Leu Pro Ser Met Leu Ile Leu Ser  
 245 250 255  
 Gly Leu Thr Ala Gly Met Leu Met Thr Asp Ala Gly Arg Lys Leu Tyr  
 260 265 270  
 Val Asn Thr Trp Ile Tyr Gly Thr Leu Leu Gly Cys Leu Trp Val Thr  
 275 280 285  
 Ile Lys Ala  
 290  
 <210> 128  
 <211> 261  
 <212> PRT  
 <213> Homo sapiens  
 <220>  
 <221> SITE  
 <222> 68..73  
 <223> Box II  
 <221> SITE  
 <222> 119..127  
 <223> Box III  
 <400> 128  
 Met Arg Tyr Leu Leu Pro Ser Val Val Leu Leu Gly Thr Ala Pro Thr  
 1 5 10 15  
 Tyr Val Leu Ala Trp Gly Val Trp Arg Leu Leu Ser Ala Phe Leu Pro  
 20 25 30  
 Ala Arg Phe Tyr Gln Ala Leu Asp Asp Arg Leu Tyr Cys Val Tyr Gln  
 35 40 45  
 Ser Met Val Leu Phe Phe Phe Glu Asn Tyr Thr Gly Val Gln Met Tyr  
 50 55 60  
 Leu Val Ile Phe Pro Glu Gly Thr Arg Tyr Asn Pro Glu Gln Thr Lys  
 65 70 75 80  
 Val Leu Ser Ala Ser Gln Ala Phe Ala Ala Gln Arg Gly Leu Ala Val  
 85 90 95  
 Leu Lys His Val Leu Thr Pro Arg Ile Lys Ala Thr His Val Ala Phe  
 100 105 110  
 Asp Cys Met Lys Asn Tyr Leu Asp Ala Ile Tyr Asp Val Thr Val Val  
 115 120 125  
 Tyr Glu Gly Lys Asp Asp Gly Gly Gln Arg Arg Glu Ser Pro Thr Met  
 130 135 140  
 Thr Glu Phe Leu Cys Lys Glu Cys Pro Lys Ile His Ile His Asp  
 145 150 155 160  
 Arg Ile Asp Lys Lys Asp Val Pro Glu Glu Gln Glu His Met Arg Arg  
 165 170 175  
 Trp Leu His Glu Arg Phe Glu Ile Lys Asp Lys Met Leu Ile Glu Phe  
 180 185 190  
 Tyr Glu Ser Pro Asp Pro Glu Arg Arg Lys Arg Phe Pro Gly Lys Ser  
 195 200 205  
 Val Asn Ser Lys Leu Ser Ile Lys Lys Thr Leu Pro Ser Met Leu Ile  
 210 215 220  
 Leu Ser Gly Leu Thr Ala Gly Met Leu Met Thr Asp Ala Gly Arg Lys  
 225 230 235 240  
 Leu Tyr Val Asn Thr Trp Ile Tyr Gly Thr Leu Leu Gly Cys Leu Trp  
 245 250 255  
 Val Thr Ile Lys Ala  
 260  
 <210> 129  
 <211> 90  
 <212> PRT  
 <213> Homo sapiens  
 <400> 129

```

Met Arg Tyr Leu Leu Pro Ser Val Val Leu Leu Gly Thr Ala Pro Thr
1          5          10          15
Tyr Val Leu Ala Trp Gly Val Trp Arg Leu Leu Ser Ala Phe Leu Pro
          20          25          30
Ala Arg Phe Tyr Gln Ala Leu Asp Asp Arg Leu Tyr Cys Val Tyr Gln
          35          40          45
Ser Met Val Leu Phe Phe Phe Glu Asn Tyr Thr Gly Val Gln Asn Phe
          50          55          60
Ser Ala Lys Asn Val Gln Lys Phe Ile Phe Thr Leu Ile Val Ser Thr
65          70          75          80
Lys Lys Met Ser Gln Lys Asn Lys Asn Ile
          85          90

```

&lt;210&gt; 130

&lt;211&gt; 68

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 130

```

Met Arg Tyr Leu Leu Pro Ser Val Val Leu Leu Gly Thr Ala Pro Thr
1          5          10          15
Tyr Val Leu Ala Trp Gly Val Trp Arg Leu Leu Ser Ala Phe Leu Pro
          20          25          30
Ala Arg Phe Tyr Gln Ala Leu Asp Asp Arg Leu Tyr Cys Val Tyr Gln
          35          40          45
Ser Met Val Leu Phe Phe Phe Glu Asn Tyr Thr Gly Val Gln Asp Ala
          50          55          60

```

Tyr Arg Ile Leu

65

&lt;210&gt; 131

&lt;211&gt; 66

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 131

```

Met Arg Tyr Leu Leu Pro Ser Val Val Leu Leu Gly Thr Ala Pro Thr
1          5          10          15
Tyr Val Leu Ala Trp Gly Val Trp Arg Leu Leu Ser Ala Phe Leu Pro
          20          25          30
Ala Arg Phe Tyr Gln Ala Leu Asp Asp Arg Leu Tyr Cys Val Tyr Gln
          35          40          45
Ser Met Val Leu Phe Phe Phe Glu Asn Tyr Thr Gly Val Gln Arg Leu
          50          55          60

```

Asp Ser

65

&lt;210&gt; 132

&lt;211&gt; 97

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; SITE

&lt;222&gt; 81..83

&lt;223&gt; Box I

&lt;400&gt; 132

```

Met Arg Tyr Leu Leu Pro Ser Val Val Leu Leu Gly Thr Ala Pro Thr
1          5          10          15
Tyr Val Leu Ala Trp Gly Val Trp Arg Leu Leu Ser Ala Phe Leu Pro
          20          25          30
Ala Arg Phe Tyr Gln Ala Leu Asp Asp Arg Leu Tyr Cys Val Tyr Gln
          35          40          45
Ser Met Val Leu Phe Phe Phe Glu Asn Tyr Thr Gly Val Gln Ile Leu
          50          55          60
Leu Tyr Gly Asp Leu Pro Lys Asn Lys Glu Asn Ile Ile Tyr Leu Ala
65          70          75          80

```



<400> 133

<400> 134

Met	Arg	Tyr	Leu	Leu	Pro	Ser	Val	Val	Leu	Leu	Gly	Thr	Ala	Pro	Thr
1				5					10					15	
Tyr	Val	Leu	Ala	Trp	Gly	Val	Trp	Arg	Leu	Leu	Ser	Ala	Phe	Leu	Pro
			20					25					30		
Ala	Arg	Phe	Tyr	Gln	Ala	Leu	Asp	Asp	Arg	Leu	Tyr	Cys	Val	Tyr	Gln
		35					40					45			
Ser	Met	Val	Leu	Phe	Phe	Phe	Glu	Asn	Tyr	Thr	Gly	Val	Gln	Ile	Leu
	50					55					60				
Leu	Tyr	Gly	Asp	Leu	Pro	Lys	Asn	Lys	Glu	Asn	Ile	Ile	Tyr	Leu	Ala
65					70					75				80	
Asn	His	Gln	Ser	Thr	Val	Asp	Trp	Ile	Val	Ala	Asp	Ile	Leu	Ala	Ile
				85					90					95	
Arg	Gln	Asn	Ala	Leu	Gly	His	Val	Arg	Tyr	Val	Leu	Lys	Glu	Gly	Leu

100 105 110  
 Lys Trp Leu Pro Leu Tyr Gly Cys Tyr Phe Ala Gln His Gly Gly Ile  
 115 120 125  
 Tyr Val Lys Arg Ser Ala Lys Phe Asn Glu Lys Glu Met Arg Asn Lys  
 130 135 140  
 Leu Gln Ser Tyr Val Asp Ala Gly Thr Pro Met Tyr Leu Val Ile Phe  
 145 150 155 160  
 Pro Glu Gly Thr Arg Tyr Asn Pro Glu Gln Thr Lys Val Leu Ser Ala  
 165 170 175  
 Ser Gln Ala Phe Ala Ala Gln Arg Gly Lys Asp Asp Gly Gly Gln Arg  
 180 185 190  
 Arg Glu Ser Pro Thr Met Thr Glu Phe Leu Cys Lys Glu Cys Pro Lys  
 195 200 205  
 Ile His Ile His Ile Asp Arg Ile Asp Lys Lys Asp Val Pro Glu Glu  
 210 215 220  
 Gln Glu His Met Arg Arg Trp Leu His Glu Arg Phe Glu Ile Lys Asp  
 225 230 235 240  
 Lys Met Leu Ile Glu Phe Tyr Glu Ser Pro Asp Pro Glu Arg Arg Lys  
 245 250 255  
 Arg Phe Pro Gly Lys Ser Val Asn Ser Lys Leu Ser Ile Lys Lys Thr  
 260 265 270  
 Leu Pro Ser Met Leu Ile Leu Ser Gly Leu Thr Ala Gly Met Leu Met  
 275 280 285  
 Thr Asp Ala Gly Arg Lys Leu Tyr Val Asn Thr Trp Ile Tyr Gly Thr  
 290 295 300  
 Leu Leu Gly Cys Leu Trp Val Thr Ile Lys Ala  
 305 310 315  
 <210> 135  
 <211> 300  
 <212> PRT  
 <213> Homo sapiens  
 <220>  
 <221> SITE  
 <222> 81..83  
 <223> Box I  
 <221> SITE  
 <222> 160..165  
 <223> Box II  
 <400> 135  
 Met Arg Tyr Leu Leu Pro Ser Val Val Leu Leu Gly Thr Ala Pro Thr  
 1 5 10 15  
 Tyr Val Leu Ala Trp Gly Val Trp Arg Leu Leu Ser Ala Phe Leu Pro  
 20 25 30  
 Ala Arg Phe Tyr Gln Ala Leu Asp Arg Arg Leu Tyr Cys Val Tyr Gln  
 35 40 45  
 Ser Met Val Leu Phe Phe Phe Glu Asn Tyr Thr Gly Val Gln Ile Leu  
 50 55 60  
 Leu Tyr Gly Asp Leu Pro Lys Asn Lys Glu Asn Ile Ile Tyr Leu Ala  
 65 70 75 80  
 Asn His Gln Ser Thr Val Asp Trp Ile Val Ala Asp Ile Leu Ala Ile  
 85 90 95  
 Arg Gln Asn Ala Leu Gly His Val Arg Tyr Val Leu Lys Glu Gly Leu  
 100 105 110  
 Lys Trp Leu Pro Leu Tyr Gly Cys Tyr Phe Ala Gln His Gly Gly Ile  
 115 120 125  
 Tyr Val Lys Arg Ser Ala Lys Phe Asn Glu Lys Glu Met Arg Asn Lys  
 130 135 140  
 Leu Gln Ser Tyr Val Asp Ala Gly Thr Pro Met Tyr Leu Val Ile Phe  
 145 150 155 160  
 Pro Glu Gly Thr Arg Tyr Asn Pro Glu Gln Thr Lys Val Leu Ser Ala  
 165 170 175

Ser Gln Ala Phe Ala Ala Gln Arg Glu Phe Leu Cys Lys Glu Cys Pro  
 180 185 190  
 Lys Ile His Ile His Ile Asp Arg Ile Asp Lys Lys Asp Val Pro Glu  
 195 200 205  
 Glu Gln Glu His Met Arg Arg Trp Leu His Glu Arg Phe Glu Ile Lys  
 210 215 220  
 Asp Lys Met Leu Ile Glu Phe Tyr Glu Ser Pro Asp Pro Glu Arg Arg  
 225 230 235 240  
 Lys Arg Phe Pro Gly Lys Ser Val Asn Ser Lys Leu Ser Ile Lys Lys  
 245 250 255  
 Thr Leu Pro Ser Met Leu Ile Leu Ser Gly Leu Thr Ala Gly Met Leu  
 260 265 270  
 Met Thr Asp Ala Gly Arg Lys Leu Tyr Val Asn Thr Trp Ile Tyr Gly  
 275 280 285  
 Thr Leu Leu Gly Cys Leu Trp Val Thr Ile Lys Ala  
 290 295 300  
 <210> 136  
 <211> 185  
 <212> PRT  
 <213> Homo sapiens  
 <220>  
 <221> SITE  
 <222> 81..83  
 <223> Box I  
 <221> SITE  
 <222> 160..165  
 <223> Box II  
 <400> 136  
 Met Arg Tyr Leu Leu Pro Ser Val Val Leu Leu Gly Thr Ala Pro Thr  
 1 5 10 15  
 Tyr Val Leu Ala Trp Gly Val Trp Arg Leu Leu Ser Ala Phe Leu Pro  
 20 25 30  
 Ala Arg Phe Tyr Gln Ala Leu Asp Asp Arg Leu Tyr Cys Val Tyr Gln  
 35 40 45  
 Ser Met Val Leu Phe Phe Phe Glu Asn Tyr Thr Gly Val Gln Ile Leu  
 50 55 60  
 Leu Tyr Gly Asp Leu Pro Lys Asn Lys Glu Asn Ile Ile Tyr Leu Ala  
 65 70 75 80  
 Asn His Gln Ser Thr Val Asp Trp Ile Val Ala Asp Ile Leu Ala Ile  
 85 90 95  
 Arg Gln Asn Ala Leu Gly His Val Arg Tyr Val Leu Lys Glu Gly Leu  
 100 105 110  
 Lys Trp Leu Pro Leu Tyr Gly Cys Tyr Phe Ala Gln His Gly Gly Ile  
 115 120 125  
 Tyr Val Lys Arg Ser Ala Lys Phe Asn Glu Lys Glu Met Arg Asn Lys  
 130 135 140  
 Leu Gln Ser Tyr Val Asp Ala Gly Thr Pro Met Tyr Leu Val Ile Phe  
 145 150 155 160  
 Pro Glu Gly Thr Arg Tyr Asn Pro Glu Gln Thr Lys Val Leu Ser Ala  
 165 170 175  
 Ser Gln Ala Phe Ala Ala Gln Arg Gly  
 180 185  
 <210> 137  
 <211> 19  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> misc\_binding  
 <222> 1..19  
 <223> amplification oligonucleotide PG1ASel3  
 <400> 137

accgggggtcc agttgactg 19  
<210> 138  
<211> 17  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..17  
<223> amplification oligonucleotide PG1ASe14  
<400> 138  
cggggtccag catggag 17  
<210> 139  
<211> 16  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..16  
<223> amplification oligonucleotide PG1ASe15  
<400> 139  
ccgggggtcca ggcctt 16  
<210> 140  
<211> 16  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..16  
<223> amplification oligonucleotide PG1ASe16  
<400> 140  
cggggtccag gccttg 16  
<210> 141  
<211> 21  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..21  
<223> amplification oligonucleotide PG1ASe17  
<400> 141  
accgggggtcc agaatttctc t 21  
<210> 142  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..19  
<223> amplification oligonucleotide PG1ASe18  
<400> 142  
cggggtccag gatgcttat 19  
<210> 143  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..23  
<223> amplification oligonucleotide PG1ASe24  
<400> 143  
aatcatcaaa gcacagcatg gag 23

```

<210> 144
<211> 28
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..28
<223> amplification oligonucleotide PG1ASe25
<400> 144
caaatcatca aagcacagat gtatcttg                28
<210> 145
<211> 20
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..20
<223> amplification oligonucleotide PG1ASe26
<400> 145
atcaaagcac aggccttgca                20
<210> 146
<211> 26
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..26
<223> amplification oligonucleotide PG1ASe27
<400> 146
agcaaatcat caaagcacag aatttc                26
<210> 147
<211> 28
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..28
<223> amplification oligonucleotide PG1ASe28
<400> 147
atcatcaaag cacaggatgc ttatagaa                28
<210> 148
<211> 31
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..31
<223> amplification oligonucleotide PG1ASe35
<400> 148
gtgttacttt gctcagatgt atcttgat t                31
<210> 149
<211> 23
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..23
<223> amplification oligonucleotide PG1ASe36
<400> 149
tactttgctc aggccttgca gta                23
<210> 150

```

<211> 27  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..27  
<223> amplification oligonucleotide PG1ASe37  
<400> 150  
gggtgttact ttgctcagaa tttctct 27  
<210> 151  
<211> 29  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..29  
<223> amplification oligonucleotide PG1ASe38  
<400> 151  
ggtgttactt tgctcaggat gcttataga 29  
<210> 152  
<211> 20  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..20  
<223> amplification oligonucleotide PG1ASe46  
<400> 152  
caggaactcc agccttgag 20  
<210> 153  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..23  
<223> amplification oligonucleotide PG1ASe47  
<400> 153  
caggaactcc aaatttctct gca 23  
<210> 154  
<211> 25  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..25  
<223> amplification oligonucleotide PG1ASe48  
<400> 154  
cgcaggaact ccagatgctt ataga 25  
<210> 155  
<211> 22  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..22  
<223> amplification oligonucleotide PG1ASe57  
<400> 155  
ctgccaacg tgaatttctc tg 22  
<210> 156  
<211> 22

<212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> misc\_binding  
 <222> 1..22  
 <223> amplification oligonucleotide PG1ASe58  
 <400> 156  
 gcccaacgtg gatgcttata ga 22  
 <210> 157  
 <211> 23  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> misc\_binding  
 <222> 1..23  
 <223> amplification oligonucleotide PG1ASe68  
 <400> 157  
 cgaccatgac gggatgctta tag 23  
 <210> 158  
 <211> 19  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> misc\_binding  
 <222> 1..19  
 <223> amplification oligonucleotide PG1ASe1X  
 <400> 158  
 ccggggtcca gagattgga 19  
 <210> 159  
 <211> 26  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> misc\_binding  
 <222> 1..26  
 <223> amplification oligonucleotide PG1ASeX2  
 <400> 159  
 aaagtggaag gccctcttta acaata 26  
 <210> 160  
 <211> 25  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> misc\_binding  
 <222> 1..25  
 <223> amplification oligonucleotide PG1Ae1b3  
 <400> 160  
 gccctcttta acattgactg gattg 25  
 <210> 161  
 <211> 24  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> misc\_binding  
 <222> 1..24  
 <223> amplification oligonucleotide PG1Ae1b4  
 <400> 161  
 gccctcttta acacatggag gaat 24  
 <210> 162  
 <211> 28  
 <212> DNA

<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..28  
<223> amplification oligonucleotide PG1Aelb5  
<400> 162  
ggccctcttt aacaatgtat cttgtgat 28  
<210> 163  
<211> 25  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..25  
<223> amplification oligonucleotide PG1Aelb6  
<400> 163  
gccctcttta acagccttgc agtat 25  
<210> 164  
<211> 25  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..25  
<223> amplification oligonucleotide PG1Aelb7  
<400> 164  
ggccctcttt aacaaatttc tctgc 25  
<210> 165  
<211> 28  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..28  
<223> amplification oligonucleotide PG1Aelb8  
<400> 165  
gaaggccctc tttaacagat gcttatag 28  
<210> 166  
<211> 26  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..26  
<223> amplification oligonucleotide PG1Ae3b4  
<400> 166  
atgctggatt atagcatgga ggaatc 26  
<210> 167  
<211> 31  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> misc\_binding  
<222> 1..31  
<223> amplification oligonucleotide PG1Ae3b5  
<400> 167  
caaatgctg gattatagat gtatcttgtg a 31  
<210> 168  
<211> 23  
<212> DNA  
<213> Homo Sapiens



```

<220>
<221> misc_binding
<222> 1..23
<223> amplification oligonucleotide PG1Ae3b6
<400> 168
tgctggatta taggccttgc agt 23
<210> 169
<211> 28
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..28
<223> amplification oligonucleotide PG1Ae3b7
<400> 169
tgctggatta tagaatttct ctgcaaag 28
<210> 170
<211> 30
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..30
<223> amplification oligonucleotide PG1Ae3b8
<400> 170
ccaaaatgct ggattatagg atgcttatag 30
<210> 171
<211> 21
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..21
<223> amplification oligonucleotide PG1Ae5b6
<400> 171
tatctttgca tggcagcctt g 21
<210> 172
<211> 23
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..23
<223> amplification oligonucleotide PG1Ae5b7
<400> 172
ctttgcatgg caaatttctc tgc 23
<210> 173
<211> 27
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..27
<223> amplification oligonucleotide PG1Ae5b8
<400> 173
ttatctttgc atggcagatg cttatag 27
<210> 174
<211> 20
<212> DNA
<213> Homo Sapiens
<220>

```

```

<221> misc_binding
<222> 1..20
<223> amplification oligonucleotide PG1Ae56b
<400> 174
ctgccaacg tgggaaagac
<210> 175
<211> 21
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..21
<223> amplification oligonucleotide PG1Ae46b
<400> 175
gcaggaactc caggaaagac g
<210> 176
<211> 25
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..25
<223> amplification oligonucleotide PG1Ae36b
<400> 176
tggtactttg ctcagggaac gacga
<210> 177
<211> 22
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..22
<223> amplification oligonucleotide PG1Ae26b
<400> 177
atcaaagcac agggaaagac ga
<210> 178
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> misc_binding
<222> 1..19
<223> amplification oligonucleotide PG1Ae16b
<400> 178
ccgggggtcca gggaaagac
<210> 179
<211> 56520
<212> DNA
<213> Homo sapiens
<220>
<221> exon
<222> 2001..2216
<223> exon1
<221> exon
<222> 18196..18265
<223> exon2
<221> exon
<222> 23716..23831
<223> exon3
<221> exon
<222> 25570..25659

```

<223> exon4  
<221> exon  
<222> 34668..34758  
<223> exon5  
<221> exon  
<222> 40685..40843  
<223> exon6  
<221> exon  
<222> 48067..48190  
<223> exon7  
<221> exon  
<222> 50179..54519  
<223> exon8  
<221> polyA\_signal  
<222> 54493..54498  
<223> AATAAA  
<221> primer\_bind  
<222> 1991..2008  
<223> upstream amplification primer 5-63  
<221> primer\_bind  
<222> 2505..2525  
<223> downstream amplification primer 5-63 , complement  
<221> primer\_bind  
<222> 4091..4111  
<223> downstream amplification primer 99-622  
<221> primer\_bind  
<222> 4528..4546  
<223> upstream amplification primer 99-622 , complement  
<221> primer\_bind  
<222> 5475..5495  
<223> downstream amplification primer 99-621  
<221> primer\_bind  
<222> 5927..5947  
<223> upstream amplification primer 99-621 , complement  
<221> primer\_bind  
<222> 8127..8144  
<223> downstream amplification primer 99-619  
<221> primer\_bind  
<222> 8560..8578  
<223> upstream amplification primer 99-619 , complement  
<221> primer\_bind  
<222> 11622..11639  
<223> upstream amplification primer 4-76  
<221> primer\_bind  
<222> 12018..12037  
<223> downstream amplification primer 4-76 , complement  
<221> primer\_bind  
<222> 11930..11947  
<223> upstream amplification primer 4-77  
<221> primer\_bind  
<222> 12339..12358  
<223> downstream amplification primer 4-77 , complement  
<221> primer\_bind  
<222> 12915..12932  
<223> upstream amplification primer 4-71  
<221> primer\_bind  
<222> 13317..13334  
<223> downstream amplification primer 4-71 , complement  
<221> primer\_bind  
<222> 13216..13233  
<223> upstream amplification primer 4-72

<221> primer\_bind  
<222> 13617..13636  
<223> downstream amplification primer 4-72 , complement  
<221> primer\_bind  
<222> 13547..13564  
<223> upstream amplification primer 4-73  
<221> primer\_bind  
<222> 13962..13981  
<223> downstream amplification primer 4-73 , complement  
<221> primer\_bind  
<222> 15994..16011  
<223> downstream amplification primer 99-610  
<221> primer\_bind  
<222> 16463..16480  
<223> upstream amplification primer 99-610 , complement  
<221> primer\_bind  
<222> 17304..17324  
<223> downstream amplification primer 99-609  
<221> primer\_bind  
<222> 17814..17832  
<223> upstream amplification primer 99-609 , complement  
<221> primer\_bind  
<222> 18008..18027  
<223> upstream amplification primer 4-90  
<221> primer\_bind  
<222> 18423..18442  
<223> downstream amplification primer 4-90 , complement  
<221> primer\_bind  
<222> 18699..18716  
<223> downstream amplification primer 99-607  
<221> primer\_bind  
<222> 19164..19182  
<223> upstream amplification primer 99-607 , complement  
<221> primer\_bind  
<222> 22589..22609  
<223> downstream amplification primer 99-602  
<221> primer\_bind  
<222> 23111..23129  
<223> upstream amplification primer 99-602 , complement  
<221> primer\_bind  
<222> 25098..25118  
<223> downstream amplification primer 99-600  
<221> primer\_bind  
<222> 25657..25674  
<223> upstream amplification primer 99-600 , complement  
<221> primer\_bind  
<222> 26537..26557  
<223> downstream amplification primer 99-598  
<221> primer\_bind  
<222> 27022..27040  
<223> upstream amplification primer 99-598 , complement  
<221> primer\_bind  
<222> 32262..32281  
<223> downstream amplification primer 99-592  
<221> primer\_bind  
<222> 32823..32841  
<223> upstream amplification primer 99-592 , complement  
<221> primer\_bind  
<222> 34215..34233  
<223> upstream amplification primer 99-217  
<221> primer\_bind

<222> 34624..34644  
<223> downstream amplification primer 99-217 , complement  
<221> primer\_bind  
<222> 34473..34491  
<223> upstream amplification primer 5-47  
<221> primer\_bind  
<222> 34916..34936  
<223> downstream amplification primer 5-47 , complement  
<221> primer\_bind  
<222> 34702..34722  
<223> downstream amplification primer 99-589  
<221> primer\_bind  
<222> 35182..35200  
<223> upstream amplification primer 99-589 , complement  
<221> primer\_bind  
<222> 39591..39611  
<223> upstream amplification primer 99-12899  
<221> primer\_bind  
<222> 39971..39991  
<223> downstream amplification primer 99-12899 , complement  
<221> primer\_bind  
<222> 40531..40549  
<223> upstream amplification primer 4-12  
<221> primer\_bind  
<222> 40932..40950  
<223> downstream amplification primer 4-12 , complement  
<221> primer\_bind  
<222> 40629..40649  
<223> downstream amplification primer 99-582  
<221> primer\_bind  
<222> 41058..41078  
<223> upstream amplification primer 99-582 , complement  
<221> primer\_bind  
<222> 45729..45746  
<223> downstream amplification primer 99-576  
<221> primer\_bind  
<222> 46186..46203  
<223> upstream amplification primer 99-576 , complement  
<221> primer\_bind  
<222> 47879..47896  
<223> upstream amplification primer 4-13  
<221> primer\_bind  
<222> 48217..48236  
<223> downstream amplification primer 4-13 , complement  
<221> primer\_bind  
<222> 48902..48922  
<223> upstream amplification primer 99-12903  
<221> primer\_bind  
<222> 49331..49351  
<223> downstream amplification primer 99-12903 , complement  
<221> primer\_bind  
<222> 49830..49848  
<223> upstream amplification primer 5-56  
<221> primer\_bind  
<222> 50271..50290  
<223> downstream amplification primer 5-56 , complement  
<221> primer\_bind  
<222> 50172..50189  
<223> upstream amplification primer 4-61  
<221> primer\_bind  
<222> 50573..50591

<223> downstream amplification primer 4-61 , complement  
<221> primer\_bind  
<222> 50541..50560  
<223> upstream amplification primer 4-62  
<221> primer\_bind  
<222> 50940..50959  
<223> downstream amplification primer 4-62 , complement  
<221> primer\_bind  
<222> 50555..50572  
<223> upstream amplification primer 4-63  
<221> primer\_bind  
<222> 50964..50983  
<223> downstream amplification primer 4-63 , complement  
<221> primer\_bind  
<222> 50774..50792  
<223> upstream amplification primer 4-64  
<221> primer\_bind  
<222> 51183..51202  
<223> downstream amplification primer 4-64 , complement  
<221> primer\_bind  
<222> 51146..51165  
<223> upstream amplification primer 4-65  
<221> primer\_bind  
<222> 51479..51496  
<223> downstream amplification primer 4-65 , complement  
<221> primer\_bind  
<222> 51593..51610  
<223> upstream amplification primer 4-67  
<221> primer\_bind  
<222> 29734..29744  
<223> upstream amplification primer 4-67 , complement  
<221> primer\_bind  
<222> 51167..51185  
<223> upstream amplification primer 5-50  
<221> primer\_bind  
<222> 51667..51687  
<223> downstream amplification primer 5-50 , complement  
<221> primer\_bind  
<222> 51387..51403  
<223> upstream amplification primer 5-71  
<221> primer\_bind  
<222> 51826..51843  
<223> downstream amplification primer 5-71 , complement  
<221> primer\_bind  
<222> 51772..51789  
<223> upstream amplification primer 5-30  
<221> primer\_bind  
<222> 52199..52217  
<223> downstream amplification primer 5-30 , complement  
<221> primer\_bind  
<222> 51850..51867  
<223> upstream amplification primer 5-58  
<221> primer\_bind  
<222> 52382..52400  
<223> downstream amplification primer 5-58 , complement  
<221> primer\_bind  
<222> 52507..52527  
<223> upstream amplification primer 5-53  
<221> primer\_bind  
<222> 52997..53017  
<223> downstream amplification primer 5-53 , complement

```

<221> primer_bind
<222> 52703..52721
<223> upstream amplification primer 5-60
<221> primer_bind
<222> 53142..53162
<223> downstream amplification primer 5-60 , complement
<221> primer_bind
<222> 53001..53018
<223> upstream amplification primer 5-68
<221> primer_bind
<222> 53521..53538
<223> downstream amplification primer 5-68 , complement
<221> primer_bind
<222> 53459..53476
<223> upstream amplification primer 5-66
<221> primer_bind
<222> 53920..53940
<223> downstream amplification primer 5-66 , complement
<221> primer_bind
<222> 54202..54220
<223> upstream amplification primer 5-62
<221> primer_bind
<222> 54681..54701
<223> downstream amplification primer 5-62 , complement
<400> 179
gtggatctgt gactgttcgc aggaagagag gagcgggagc aggacagaca ataactgata 60
gtcaggagct ggggtttggag ataaagaggg aacaagagaa agttaagttc tgtgttttca 120
tggcaaacat tgcacaaaag tttacaactt cgtgactaac agtaatctgg ggtgattcac 180
aacaaattta cacataaaca catatcttact gactttatac acagcaatcc taacgtgaac 240
acagaacctg ctttatcttt tgcacactg ttctagtgtg gagatgtctg gtctcagtta 300
aagaaagcat aaggagcatt agttgtgcac actgtccaca cccgtgactt tttccacca 360
gtactaaacc tagtgcttct tacagtacag ggcaatgaca gccacagaaa gagagaagct 420
ccttttactg tgtaatgctt cctgctggcc ttcaaatact tgttacttga gagatctcca 480
ttcacctggc tttgtcccca aaggatcatca tctaccaatg atgttggtat ttgatgttaa 540
tcatgtataa agaaagtagc taccatcctg gccctgatta gaacttccca ctgaaatacc 600
gtcctgccta aaggtagcac aggtttccat tatggtgggt gtggggaggg ggcgggaata 660
tatatatata tatatatata tatatatatg gtaaagcatt cggcattctt ttaaagtaca 720
actatccttg aaaagggtta catattaaac catttttacc acagccaaag gggaggagaa 780
agatccaaaa gtccctgtgga tctgctttaa catcaataaa acagttatcc acccttcgta 840
gcttttagtg aaggctacaa agtatgctt ttatggatt acacatgtgc acgcaactac 900
tttaattact acagaaaaaa acgaggctcc ttattaaaaa aaaatcagaa acaagtccaa 960
cagactctga ggaaatgaag caagagtga tttctgaaaag gtctaataaa cagtatggaa 1020
atatacctgt gggattgttc ttcagctatg cataaacatg taattatcat cattactgtg 1080
atggggaaaa acacggaccc taattctgaa acaccctggt agcgagagac gggcaggagg 1140
ggctgctgag cactcagagc ggaggctgag gagcgcgct ccccttgcaa aggactggca 1200
gtgagcagat ggggacactc gagctgcccc gcgacctggg ccgagctgcc tacaacctgg 1260
gcccaggctg ctgcaagaat tagacctccg ataactgtaa caccacttt ctactgctc 1320
taattgtgtg catcccggcg cccaggggct tgtgagcagc aggtgcgctg tccaggcagc 1380
tccagcgacc cttaaacctg accgcgcgca cgtccggccc gagggagcag aacaagaggc 1440
accgggaccc tccctcggcc agcaccaccc ttcaccagct tccgtcagtc gccaccacct 1500
cccttcccgc gtcgcgagcc ggcccagctg gggagcatgc gcagtggccg gagccgggtt 1560
gcccgcgcca cagcaggtag ctgtactgca actgtcggcc caaaccaacc aatcaagaga 1620
cgtgttattg ccgcccagggt ggaactatgg caacgggcga ccaatcagaa ggcgcgttgt 1680
tgccgcggag cccctgccc cggcaggggg atgtggcgat ggggtgaggt catggggtgt 1740
gagcatccct gagccatcga tccgggaggg ccgcgggttc ccttgctttg ccgcggggag 1800
cggcgcacgc agccccgcac tgcctaccc ggccccgggc ggccccgggc cgggtgattgg 1860
tgggggcgga ggcctgggagc ggggtggcgg cgcgcgggcc cggggccggg cgggtgattgg 1920
ccgcctgctg ccgcgcactg agggccggga ggcggggcgg gagcgaggc ggagctcgct 1980
gccgcgagc tgagaagatg ctgctgtccc tgggtgctcca cactgtactcc atgcgctacc 2040
tgctgcccag cgtcgtgctc ctgggcacgg cgcccaccta cgtgttggtc tggggggtct 2100
ggcggtgct ctccgccttc ctgcccggcc gcttctacca agcgtggac gaccggctgt 2160

```

actgcgtcta	ccagagcatg	gtgctcttct	tcttcgagaa	ttacaccggg	gtccaggtga	2220
gccgcctccc	gctccccggg	ctcggcgtcc	acccgagctc	ccggggggcg	ggacctctcc	2280
gctccccac	agctggcgag	ggtcaccggg	ccggcccgcc	ggaccagca	cggagagcac	2340
gtgcgcctc	ccgccttcc	tctccgcatg	cttccctgcc	ttotgccgag	atcgctctct	2400
aggaagctgt	ggctgcgtcg	tcctgaggct	acgagtggga	ccgcgcgcc	ctttcccgcc	2460
ccctgcctcg	ggctgatgc	tgcttagcaa	agtgggtgca	gatgcacgtt	ttaaataata	2520
gggcacgcgt	ttagcagttt	ctggcctttg	gtccaaagag	gtgggtcatgt	tggaacagat	2580
cggagacgtc	tacactccga	agtgcgcttt	tacagtgacc	tcttgaaaca	gaagtacaat	2640
tccgtcttgt	gttctttccc	ctggacaagt	gaaagctggg	cgaagaaatg	aatacatttg	2700
ttaacgtag	aagcctaact	agatacaatt	cttgccaact	ttactgggc	ttgaatgtgt	2760
gggtgatctg	ttgtctgatt	actttctttc	tggtactgtt	tctctgtaga	gattggattc	2820
gtagattaaa	cttgagaaac	aaaccataaa	agtgggaagg	cctctttaac	agtaggtatt	2880
tgaagtgtta	taaaaaaaaa	aaaggtgaat	ttttctttta	tttctcagtt	tgaagaaca	2940
gctttattct	tggttattcc	taatgtccac	ctagtctctc	tttacttttc	ttggtagggg	3000
taggggtgca	tgggggaaatg	ggacgggtatc	attttgtctt	tttaactttt	tttttttcca	3060
cctacagcag	ctgtttttac	cctgtgggtca	gtcagggtact	atatttagtt	tgcagttgca	3120
ctgctgatcg	acccttgatg	gccccagttg	gaagttgttt	ggggggaagg	aactaggaga	3180
ggccagggcc	tccattttaa	ccagtgtctg	taagtgtctc	cttggaagga	aaaaaagata	3240
ctgttccagg	tcattggtttc	ctggtagttg	acgtttaaaa	tgggcctcat	ttaaaaattt	3300
caataattca	ggcataattt	ttccctttat	atggtaactc	caccaagttt	gtctaattgt	3360
atgattttta	tcattgattaa	gtttttactt	ccacatcatg	tgacaactgg	cctgggatgg	3420
gatataagct	cagaacacaa	agtcatcac	ctgttaaaaa	aataattcta	tctgtggcgg	3480
gttatgttat	ttttgttcaa	agaggacaca	atatgatgca	gaatacacca	ttgaaggatt	3540
ttttggtttg	gcaagttctt	atttttttaa	atggctgtaa	aacctagcag	tgtttctgaa	3600
attgcatacc	ttacctgatg	ttcagagatc	cgattttact	cttgatttcc	cagcaagtga	3660
ttttgaaaac	atttaactca	atcattcccc	ccaccgtctg	ttcaaatcaa	aggaagtggc	3720
atccagcact	aattttcatg	cattttatgaa	aggatgcctg	aggaccctta	agtataattc	3780
aaaattttgt	ttaatgtgtg	ttccttgatg	aagttcttta	ggagtcgtag	aacgaactga	3840
ttgcccactg	atcatcaaat	gcaagttatg	aacattttaa	aaaaatttaa	aaccaagagt	3900
ttcttgttcc	tgcattttta	tttttattgt	atggagggga	caaataatta	ttttctgttt	3960
agtaacagag	caggggtatt	tgaatttatt	agggctcttt	tctgcagtct	gggtttcctg	4020
tgtacacaaa	gctacctttc	aataattttt	attgtttctg	ttaaagattaa	atcaatagag	4080
gaataaatag	ctatcttcaa	acataagacc	caaaggaaaa	agatttatag	tgatgttctg	4140
tcaccttatt	ttttacctgt	gactttgtac	cattaacttt	gtcactgaga	tgttttgatt	4200
aaaattttta	gcttgccttt	cttgttttgt	taggacactc	ttttttctt	gaattgtttt	4260
tatcagcttt	cgtttgcaag	gctagtgatg	atctcttgt	tctgtataaa	gtattgttga	4320
ctcatttctg	aaggaggttt	tagtaattta	agaggttata	agttttttaa	taaaagggtt	4380
attaatttat	atatatttaa	gaggcatttt	aaaataaaat	ttttttttaa	tgacattttt	4440
acacctttca	actctagggt	taaaaaataa	gtggttcaca	gtagtctctg	cagaagaata	4500
ttttctttta	catagaattt	ttaagctgaa	gagaagtagt	agtaggtcca	tgagatttat	4560
gatctgtgct	tggcaggtaa	acctgcttcc	aacaaaattta	gttggtattt	tcttggtatc	4620
tgggtaaaata	cccttttctt	ccccagtttc	actactttat	tttcatatgt	atctctgaga	4680
tagagaaaata	tttcagtcag	tgctgctaaa	attgttccct	ataactcggt	tatcctttta	4740
ggctcctcca	gaatctctca	ttggtactga	aactcaaatg	ggtactttct	tcaccattta	4800
tttcttttaga	ataagtaata	agaattttat	aagctttttt	atatttcacg	taatttgaga	4860
ctattgaaa	tccagttaag	tctctctact	gtgttgagag	gcattgatcc	aagtacctgt	4920
gttactttcc	tgtgctgcca	aaacagatca	cctcaaaacta	agcggcttaa	aataatagaa	4980
cttaagttct	cgtgattctg	gaggccagca	ctttgaaatc	aaggtgtagg	ctcaattttta	5040
ctccctctgg	aggccctagg	gggaatctgt	tcttgtgggt	ttcaacttct	ggtgactggg	5100
ggcattctct	ggcttggggc	cccatcactt	caacctctgc	cttacagtcc	ttgctgccac	5160
ctcttctgtc	tcacatctca	ctctcccttt	ctcttagaag	gatgcttgct	attgggttta	5220
gagccacact	ggatattccg	ggatgatctc	ttcatctcaa	gatccttaat	tataactgca	5280
aagagccttt	ttccaaaata	gaaaacattc	acaggttcca	gggcttagga	tgtggacaca	5340
ttttttgagg	ggctgccctt	cattccccca	caacaatgaa	ctccatagtt	ctgcctattc	5400
agtattttgt	agttatttctg	tagtttaact	tgcttatttt	cttttaggtat	ttacgtatta	5460
aagcattttg	gtctctgctt	tctttaacag	agaacctggg	tttctgtaat	aagtttactt	5520
actttcccat	aatcttttag	tttcttattt	acagatttac	cttcacatat	cccttaagta	5580
gaacattttga	ttaactgttt	tattttcgga	acaaatctgc	attctgtata	ataaccaact	5640
tattcatatt	tccgtattct	tttaattctt	atctgattct	gaaattacca	tcttgtgatt	5700
atatatatat	atatatggaa	ataactgaaa	tcttgataaa	ttaaaggtga	tataacttct	5760
aagacaatta	attatgtatg	atgtgggtgaa	tatactgggtg	tttgggttgt	ttgccactta	5820



aaagccctat	ctataggata	ggaagtaact	tgaatgtgga	atgcttagag	actcagagta	5880
agaggccgta	tatatatcct	tgagctggag	tttaaggaaa	acttatggga	aattaaaagg	5940
aaagtggag	tactgacaga	ggattgcgta	ggactcatga	aaaaggaatg	aagttacctt	6000
aaattctatc	atcgtgagtt	aacgtgaaac	tagatttatg	ttagtttata	gcctagaatt	6060
ctatcctagg	aatctagata	tatcctaaat	gttgagatag	ctgcataaac	aataactgta	6120
atcgttatga	taaataatga	caaattctttt	tagcatgttt	tgtgaagctg	ataaatgtta	6180
ataggatgtc	ttcaaagtgc	agaattctttt	tttctttgct	tcttttttaa	aaaatttctt	6240
ttccccatt	cctatgcaat	acactgaaaa	ctgatcattg	aaatttgtag	gccaaaaaat	6300
taatcaacac	gtaatatagatt	ggggtttggg	tttttttgag	tcagggtctt	cttctgtcac	6360
ccaggctctg	gtgagggtggc	accatcatgg	ctcattgcag	ccttgaatgc	ctgggttcaa	6420
gtgatcctcc	ggagtagctg	ccgtgccatt	atttctagct	aattttttaa	agtttttgta	6480
gaaatggggt	ctttctgtgt	tgcccaggct	ggtcttgaat	tcctggcctc	aggtgatcct	6540
tctgccttgg	cctcccaaag	tgctgggatt	acagggtgta	gccaccatgc	ctagccccta	6600
ataaatattc	taattaccga	tttatcttgc	ttaaatcagt	tggtaacact	tggaatttac	6660
ttcagaatat	attttacatt	agtggctctg	actgctaatt	cccccttctc	caaatgctaa	6720
tgtaataata	caataaaatg	cacagttctt	aagtttatat	aaaaataaca	ggttttcagt	6780
tgacctgctt	taagtgtaaa	atagtgtgaa	aaacacaaga	aagaagataa	agaatttaag	6840
attttgacat	ttctctataa	tgcccttaac	ttctccaagg	attcatactt	ttttttgtaa	6900
gacagaatct	cacactgttg	cccaaaccag	aggtgcagtg	gtgcagtcct	cactcactgc	6960
aaactctgcc	cccgggtcca	agcggctctc	ccacctcagc	ctctgagta	gctgggacta	7020
caggtagaca	gcaccatgcc	cagctaattt	tttttttggg	tatttttttag	tgggggtaga	7080
gacgagattt	tgccatattg	cccagtcctg	ttttgagctc	ctgggctcaa	gtgatccgtc	7140
cttgatccac	catgcttagc	tgattcatac	tcttaactga	aacattgttc	caagtttctc	7200
agaacacgtc	aaggcttttt	atctagagaa	cattataaac	tggatctttc	tttggttagc	7260
actgattcat	caaactaate	ctaaactcct	aatgagttaa	atttatatct	tgaatcttgc	7320
tgtaaaagca	gccattcatt	agaatgaaac	atgtttactt	agaattggag	aagggagctt	7380
ataagtcac	tagtctactc	ccttttatga	cacttctaca	ttctttctgc	acttctgcca	7440
aaatgttgcc	cagcgtcgtc	tctgatacct	atagtcctaa	caagaatatg	aatcatacct	7500
tgatccttta	attttactct	tctctgctta	tttgccattc	atgtgaagac	cttaaataga	7560
tcttaaattg	cttccttcac	tttagctgag	agtgacagga	ctgtgtagg	gtgggtgtgt	7620
ttctgcattt	gcttatttaa	gcaggataat	aaaaactttt	actataggaa	attaaacatt	7680
tcccaatcaa	atacaattcc	agtctaacac	aattaaattc	tggttaggga	actgcttaac	7740
ttactagact	tataggaaaa	tactaaaaaa	atgtaactag	aactctattt	ttacacttta	7800
taaatataaa	cctctgtgaa	caaaccagtt	atttcaggtt	gcatttgtgt	atagtttttt	7860
aatgcctgat	ttttctattt	taaaatcaca	gatgcaatta	tacattcaaa	cactgccaca	7920
atactttgag	aaagttaaag	tttcccttac	tcctacactg	cgtaacacct	tcctaggtac	7980
atcccagttt	gggtgtgtaac	tttagatttc	ttccaagagc	ttttgagtaa	gtgtttgaa	8040
tgtgggaagg	tttttttagtt	aaatgaactt	cttacagatc	agtttttttag	tacagtagca	8100
cgaaatatac	ctgcataacct	atgggggatac	ctctgtgcca	ttacgatgga	aggcacggga	8160
aaacagcact	ccgtatatac	ctagtttact	ttccctcttt	tgtatatttg	tctgattttg	8220
tggaagctgat	gcttctcaag	tggaaacaga	agtttaacttt	tcctttacta	ttttctcatt	8280
ttattatggt	ttcttaacta	gaggttgatg	tttagtggtg	gaccattcaa	tagtaagtaa	8340
tgacttttca	gtaagggatc	tctagaaccc	agatccctta	attcctgcaa	tattcccgtg	8400
tgtacattgt	tccaggtgct	gtcctgggta	ccaagggata	caatgtttga	tagacaatgt	8460
acctgccatt	atggagggtca	cattctagtg	tgggaagaca	aacaataaca	agaaaatgaa	8520
aattttactgt	gccatgccag	gttgtttagc	ctgggtgggtg	agaggtaggg	gtttggaaaa	8580
tcttactgag	caagtgcacat	ttgtgtggag	ctctgtaaaa	gggccagctt	ggaaggtaat	8640
gtagtcatcc	aggtagagaaa	tgatggttag	gggagtgga	agagtggatg	ttaagattga	8700
aaagaattcc	aaatctattt	tagtggtagc	tgatagggtc	ttgtgattga	atgtggagga	8760
aaaagaagag	ggtgggttag	taacacactc	agtcgcagtt	agtgaagtgc	gctgtgtgca	8820
agtattgttc	tattatgtaa	ataattccat	ctttacaaag	taggcaccat	tcttccctct	8880
ttacagacaa	ggaaaaggga	acacccatgg	ttcacatctg	tagtagccta	gccaggagtt	8940
tcaggcactt	attttctgaa	gatgctctgc	ctggcaatgt	ggttatattg	gttgaaatga	9000
gacccctac	tttcaaggta	ttcatctagg	aaagacatga	actgccaatt	acaatatagg	9060
ataaactgta	aattagagac	gtgtttatta	actttgccat	acagaggtaa	agtaactctt	9120
taaagtaact	ctttgcttgg	gttagtggag	aaggctataa	aaattacttg	gagtttttac	9180
tttgaacatg	cgtaattaac	atggaatggt	tagggaaaag	aggttttcaa	ttgataacat	9240
aataaacatg	aggagtgtga	agcatggcat	tcaagggttt	ctaaattctg	ccccgggttaa	9300
cttttccatt	cggttggtttc	attctagtct	agcttttctc	tctgggcccgc	ccctcccac	9360
attagaccgc	tctctctctg	aattccaact	caagcccttg	cttttctcca	tctgtcatga	9420
tgttaccoca	tctcattgtc	agggttaactt	ttatgtaata	ttaacatata	taatactgat	9480

ataacattag	catatttttaa	tgtatggatc	atctcctctg	caacattgta	acctcttggg	9540
gatggcaata	atgggaagaa	tgacttgatt	ttactttttc	ttttaacaaa	aatgggtggag	9600
tagtctgggc	acggtgtggc	tcatgcctgt	aatcccagca	ttttgggagg	ccaaggaggg	9660
tggatcactt	gaggtcaggc	attcgagacc	agtctggcca	acattgtgaa	accccatctc	9720
taccaaaaaa	atacaaacac	ttactgggca	tggtgggtgtg	tgctgtagt	ccttagctact	9780
caggaggctg	aggtgggaga	atcacttgaa	catgggaggt	agaggctcca	gcttgggcca	9840
cagagtgaga	ccctgtctca	aaagaaaaaa	aaggtaaaag	ggccaggtgc	ggaggctcac	9900
gctggtaatc	caagcacttt	gggaggctga	ggcaattggat	cacctgaggt	cgggagttcg	9960
agatcagcct	gaccaacatg	gagaaacccc	ttctctacta	aaaatacaaa	attagccggg	10020
cgtgggtggg	cctgcctgta	atctaagcta	catgggaggt	tgaggcagga	gaatcacttg	10080
aaccaggag	acagagggtg	tggtgagcca	agatggcacc	attgcactcc	cgactgggca	10140
acaagagcga	aattccgtct	caaaacaaac	aaacaaacaa	aacaaaacag	agagaaaagg	10200
cagagtactc	tagggaattc	tagtctgtgt	ttctgtggaa	atgtatatga	atctcacttt	10260
taagggtatg	agatttttga	atggcataac	tagttgataa	gttttgcctc	aacagggtac	10320
ccaagtctag	tgagtcgat	tcattctttc	cttaaataga	tgaaggagga	agaaacatga	10380
ctccaccctc	aagagtaagg	cagaatgagc	aaagtcagag	aagttaaaaa	agaattctca	10440
cgcagccagc	agtgcagaga	aaccttggtt	tagttgtgaa	tcaaaaccag	tactttttgt	10500
aatttttgag	cctatgcaat	tctccaaggt	tttatgttgt	ttcttctgtt	tctctgtagg	10560
caccagaaat	caaaacccca	aataagaaag	tgttacttga	agatttttaga	gtacttattt	10620
gtgtataagt	gtaagtgata	tttgggaagc	gactttactg	cgctcctcca	gcttggcatg	10680
agaattccag	gggcggaag	aaaggagggt	gatgggtac	ggaaaggaga	gtcatgttaa	10740
gtcccagcca	catattaagt	gctaaccacc	tactgttaaa	aggtgtaatg	ttctagactg	10800
acaaaataca	tagtctctac	cgtaaagtaa	cacataat	agcagtgag	aaagatgtca	10860
cttaaaagaa	aacttgaata	tatgctgaga	tagttcacaa	attaaagaaa	tgaacaaaga	10920
actgaggaaa	taaaggagga	atacaactgt	gtccaaatga	atacttaact	gggtgggagc	10980
tggttcata	gtaagcaggt	ggttcaccta	aaagttggat	gtaacgtagt	taacgccagc	11040
tcttggtgca	cttaccatatt	gcattgcttc	cgggctta	ttgtgttc	ataggaataa	11100
attttttgtt	ggttttta	tttactcctt	gtaattccgt	ggttgatatt	caaagtga	11160
aaaattacat	aagcttctaa	tatatgagaa	gtcttctcac	ttgacatttt	ttatttggaa	11220
tttttgcaga	gagtagtttt	gtcacagtca	aaagattttg	ggatcttgca	gtgagaaacc	11280
taggtgta	tcctatttct	ctgccattcc	gtatgtcatc	tggattaagt	gtcaacttct	11340
cagttctcaag	attctcgtcc	ttaaatggaa	tactttttgt	catgctattt	tgaagacaaa	11400
atgagataat	acgtgaaact	gcctagctca	gtgaatggta	catcatagat	actcagaaaa	11460
aacacaccct	ctaaaataag	aacagtacca	aaagacagga	tgtaaaataa	gggcagtagc	11520
aaaagacaca	tgcatgctga	gtgtatgaga	agaactttg	tggccttctt	gggtggcaca	11580
ggccatggca	gttccacagc	atgacgtggt	tgctgtgggt	ggtagagcag	acatgccgct	11640
ccccgtca	ccttggtgtt	gatgcttgct	ttcttcagct	gagaggagc	agctgtgata	11700
tgaagggtctt	gtgtgtacag	tcgtgacctc	acatttccaa	tttctctgtg	gcagaaccca	11760
cagtctacaa	cgtacgagca	ccagagttga	cgtgagacag	acagcataca	gaggcttgta	11820
acatccttct	ggaaaacact	gtgtgaagctt	tcagtgcgaa	taaacaatgat	cagtggcaag	11880
ttctgttaga	tgtagtctgc	aagcatcctg	attttactgg	gcaagactat	gttgatttac	11940
aggcggtga	tgattccatg	gatagccac	tactagtatt	ttcacaat	tcacaagaca	12000
ttcttactgg	aagattgccc	tggtcttatg	atactgctgc	ccttttagct	tcatttgc	12060
ttcagactaa	acttgagagc	tacagtcagt	cagagaactt	gctaggccac	ctctcaggtt	12120
attctttcat	tcctgatcat	cctcaaaat	ttgaaaaaga	aattgtaaaa	attacatcag	12180
caacatatag	gcttatgtcc	ttgagaagca	gcagttaatt	acctaaccac	agcaagtacc	12240
ttagaactct	gtggagttga	attgcactat	gcaaggatc	aagtaacaat	aaaattatga	12300
ttggaatgat	gtcaagagga	attctgattt	ataacaggct	atgaatgagt	acctttccat	12360
ggtcgaagat	tgtaaaaatt	tgttttta	gcaaacagtt	ttttattcag	ctttgaaaat	12420
gacttgcata	aatctggaga	aagattatca	ggattttaata	tggtgaatta	tatggcatgt	12480
aaacatttgt	ggaaagcaag	tttagaacat	cacatattct	tctgttttga	cagaccactt	12540
ccaactagaa	agaatttttt	tgacacattat	tttacattag	gttcaaaat	cctaattgcat	12600
ggtgggagaa	ctgaagtcca	gttagttcag	tatggcaag	aaaaggcaaa	taaagacaga	12660
ctacttgcag	gatcctcaag	taagccattg	acgtggaaat	taatagtttg	ggaagttagta	12720
ggcaggaatt	caatatctga	tgaaaagatt	agaaacataa	agccttccat	cacaattccc	12780
accgggaaca	ggaattccta	ctcatcaaaa	ttctgcattc	atacaagagg	gaacctgatt	12840
atgaccatct	tctgttggtc	atttggtaga	ttatgtggtt	cacacttctt	ccaaatattt	12900
gcaaatcaga	catcaccatt	atcagcacaa	gctaatagca	tcattctgga	atcatcacta	12960
ttacaggaga	cccctggaga	tggttagcct	ccagctttac	cacccaaaca	agctaagaaa	13020
aactgttgga	acpaaattca	ttattttacat	tttcaacaag	atctggaaga	tcatattaat	13080
gaaacgttga	tgttctatct	tctcttaaaa	aatctgctcc	taatgggtgt	attctacatg	13140

ataatcgtgt	tctaataccga	gtgaacctga	cgaaaatgga	aggtttggag	tcaatgcaaa	13200
gggggatatg	atpagaagat	gtctgtgatc	gtgtccctgag	aagcaccagg	aacacctttg	13260
acctcagtga	ctctcgattg	aagagaagac	caagttgtat	tgatcagtg	ttgggacttt	13320
acagaacaca	cccatgattg	ggttgtccctg	ctttttaaag	ccaactgtga	gagacattct	13380
ggggaactca	tgcttctagt	tctacctatg	ctgcataatga	tgtagtggaa	gaagtgtctag	13440
aaaatgagac	agacttccag	tacattcttg	agaaagcccc	actagatagt	gtccaccagg	13500
atgacctgt	gctgtgggag	tcagtgatcc	agctaaccga	gggcttatcg	ctggaacatt	13560
ctggacacaa	tttgatcaac	ttatcaaaaa	aaaacttggga	atgacaattt	ctgggtgccag	13620
attaccttag	aacctttgca	aaaatagata	gagatagttt	tccttatgat	gttacatggc	13680
ttatttttaa	aggtaatgaa	aactacatca	gtgtaattcc	agcatcataa	gtcagaacag	13740
tgcttgtcaa	ggggcgttac	cacacacttg	aacagatttt	tggcagatga	cttgggaaca	13800
aggctcctcc	atgtttgtaa	tgttgaccac	acaagttgaa	tgtggcagag	ttaaatgacc	13860
ccaatattgg	ccagaaccca	caggaagttc	atcctatgga	tgctaccaag	ccttctgcca	13920
ctgagaagaa	ggaagcactg	tctttatcct	caggaagatc	acactgctgt	ttaaccaaga	13980
gaaaaattag	agagtcacat	atcacgcaga	tccagtacag	agggtggcct	gaccatggag	14040
acctgatga	ttcagtgaact	ttctggattt	tgcttttcat	atgcaaaata	agagggtctag	14100
caaggaaaaa	cccttgtgtg	tttcttgag	tgctggagtt	ggaagaacca	gcgttcttaa	14160
tactatggaa	acagccatgt	gtctcattga	tctcattgaa	tgcatcagc	cagtttattc	14220
actagacatg	gtaagaacaa	tgagagagca	gtgagccgtg	atggtccaaa	cacctagttca	14280
ttacagtttt	gcgtgtgaag	tactattttg	aaagccttatg	aagaaggcct	tgctgaagaa	14340
agcaaaagga	aaaaaagaac	tttgtcatct	gttaggttcc	atttattgca	tgataattgt	14400
gtttgtattg	attattgggc	aagtagctgt	ttgtcatttt	gatcttattt	cagaagggca	14460
taataatttt	actattcaat	gaaacgtttt	aaacggggta	gaaaaagact	agtttttgta	14520
tgctttacag	cagaaatcct	ataatgatta	actggaata	tatttcgttg	gcataaaaat	14580
acatttaaaa	gttcaagtaa	ttataaacat	tgtaaatgt	atatgtaatc	atattgaaat	14640
tgaaattctt	tatagctgta	cttctgtgta	atcaaagact	ggggagagat	agactagcta	14700
gctctttctc	ttatccatta	atcacttaac	agagttttga	ataaaaagtt	ccatttcatg	14760
ggataagaat	aatgacaggt	taacctattt	tagttgggta	ctatgttcta	ggtgtgtgat	14820
gaagtagttt	acatagtttc	actgatttca	ctacaatccc	aggaggagta	gttactatta	14880
ttacactcat	tttacaggca	aagaaatagg	tttggagggg	ttgggtgttt	tgccccagtt	14940
ctcatcgtaa	aatgacagat	gaggattcaa	attcaagtct	taattgaagt	ccattacttt	15000
agaacctacc	tcttagtggc	tcttatgtta	cagtataagg	gagagcagac	tgttccttta	15060
ccctttagag	gtagctaggg	cttgtgaatt	aagagactga	ttaacaggag	aagaggcata	15120
cacattttat	tgacgttagt	atttttacat	gcacagggaa	ggagggtttt	atttttattt	15180
ttatttttat	ctttatttta	aagagacagg	ggtcttgctg	tgtagccagg	gctggactca	15240
aactcctgaa	gccaagcgat	tcttctgctt	gagattcctg	agtagcaggg	actatagggtg	15300
tgctcctctg	tgcttggcta	aagaaggggt	ttgtatgtga	tttttaacaa	aggctgataa	15360
attgtgaaga	agtgactagt	caaaggagaa	gaggatttca	gctcccaggg	gtggtaaat	15420
gtgggaagat	gactaggaaa	tgtatagtaa	taagggttgc	tatgcagggt	tattttgcca	15480
gtttctggtc	tcctaataag	ggacagggaa	acacctttac	agatggaaat	tcatatcacc	15540
tttccacagg	gaaattttatg	tctgcctta	ggcagttagg	ggaagggcag	agaattcttc	15600
ctgtatctgc	tggtgtctcag	gtgccttcag	ctcaaaataa	tccttatgcc	aaagtagcat	15660
atttgggtgt	ggcatattct	ctgatctctt	tcaacagcat	catctatact	taacaacagc	15720
aaaagttttt	tttaaaaaat	catgtttcaa	gatttgcagt	tggaagacaa	atggacatga	15780
ttgagataaa	tgaagaatat	atatttttta	acaaagaatg	ctgtatat	atgtctctgt	15840
gacattgtgt	tatggaggct	aaggtgttaa	gcattgtgatt	acttttagatg	ccgtatgact	15900
acctgttttt	aagattaaaa	aagaatcaat	aggcagttta	tatgcattggg	agcaagttaa	15960
aaacaacaca	gatgtgatga	aggcgagggtg	aaactgggtcc	gcattctaatt	caggccttct	16020
cctgaaagcc	agtgtgtgca	agataaataa	gtttgtttga	cgaaagcaga	ataactagtt	16080
tgctccttgt	gatgaagata	gttattcaga	aatcattttt	attggctacc	tctgaattaa	16140
taaatgaaaa	gagaaatttt	tttttctgta	ggggatgtct	gatgagttct	taaaaagtgg	16200
atgaacctga	aattatcatg	aacaagcaat	tataatgaac	ttaaaattac	ttaaagagtt	16260
atgaaaaaca	aaaagaaaag	ccgtatgttt	tcttgtgcct	tattttgaag	tgacaaatta	16320
tttgcagggt	acattttag	acggaactaa	tgtgatttaa	aaaatgagta	ctagatttac	16380
agaatgatgc	ctttaaaaaag	tactgtgtgc	actttaatta	ttttatttat	gtttattctg	16440
aaactacctt	tattttgaaa	atgagggtata	gctttgccta	ctggtgacaa	aagtgtaaat	16500
aattcacgtga	acatctgtta	aaaaccagct	tggtgctagg	ctcttggggg	agaaaactga	16560
tcaggccatt	gaggagctca	tagtccctaa	ggggctgggg	acttgtcatt	agggtgtcag	16620
tgtgttctgg	atgctcctga	aggagtgtgg	gcaggtgcgc	accaccatgc	ctggctaata	16680
tttttataat	tatgtagaga	cagggtctgg	ctgtgctgcc	catgtcgggt	ttgaacttct	16740
gggcttaaga	gatcttccct	ccttgcctcc	accgaccccg	cccgcctcact	ccacctcagc	16800

ctccccaag	cactgggatt	gcaggcatgg	gccactatgc	ctgggctgtg	caaaactttt	16860
aaatcagtg	atactcaatg	gtcttgatgc	aattctggct	tggttgtaag	agaatgggga	16920
ttactcaca	agccacgatg	tcacttttaa	ctctgaacag	atcaagctat	tggtattact	16980
catttatgtc	atcgataaac	tttatgaata	aaaactcatt	gtgcaaatat	ttaaatacat	17040
tacatacata	gcactgtgca	gtttctaagg	aaagtaatgg	aaacctttgt	cacatccctg	17100
gcttccagaa	ctttatgtta	tctaagtgca	tttgtctgca	aagttgttgg	gttaattgcc	17160
cctttctttc	ttctcttttt	aagatattaa	taaatagtgt	catgaccaa	agataatcct	17220
tatggacaag	atagatctaa	aaagccttag	ctaatttata	atcttgcata	atccatgatg	17280
acaagatgca	gaaacaaaaa	tgcccagaat	aaaaacttag	caccattagc	agccatttcc	17340
ttttaagtct	ttacaagtat	actcccagtt	tcttgaaaaa	tttattctaa	aatatgtaag	17400
acacacaaaa	cagcagaagg	actaatacag	gtacatcgaa	cacctgtgtg	cctaccgccc	17460
agtttaaaaa	taaaactggaa	tgatgtttct	ctcactctta	cagaataaag	ttttaatcct	17520
tagcatggaa	ttcaaaaagac	tcttgccatt	ccagttcaga	gccacccttc	tggtctcctt	17580
gctcctcagc	cgpgacactg	cccatgtacc	caacaggcct	ccagggttac	tgcttccatt	17640
cgttcttatt	ctpatgaaca	ttttccttca	tctcatctgc	cagaatccta	cctaataata	17700
ctcctgctct	gcagtttaca	gttcttttaa	attaaaaaag	gttgtgtacc	ctttagtgtc	17760
ctgaaaaaag	aaaaaacaaa	tttaaaacct	taaaaaggta	ccatattttc	atagtatttg	17820
cgttatgtct	cattacagtt	cctgtggaca	tgtctgtctc	ttttactaga	ttgattgtgg	17880
gctctttgaa	ggaagatata	tcttatgaac	agtgttttat	atattgttag	caatcaatga	17940
atgcttgcta	tatttttctc	atgaggatat	tgattattct	attttaattt	attaccgtta	18000
acctgtacta	tacataactg	ctttctgtac	ctgagctatt	tatgatctct	gaggctcctg	18060
tgagaaatct	aatttttggt	aatcatggat	ggaaatattc	acaacatcat	tcgtcagttt	18120
cttcacattg	tcttcctttg	tatattacag	atgtttttaa	atatcaaagt	aatgtttttt	18180
tgttttatct	tttagatatt	gctatatgga	gatttggcaa	aaaataaaga	aaatataata	18240
tatttagcaa	atcatcaaaag	cacaggtttg	tatttcattt	gcataaaacc	taggtttttt	18300
tacagatggc	acatgggcat	tcaaaatacc	gttcttatat	ttaaataaag	tggttttttt	18360
aaaacagcaa	ttttctgtgc	agatattaca	cctgttcttg	tatttttggt	attttacttt	18420
ttggaaagtc	agaaacttga	aagctatgaa	ttttcctaaa	cttaccttct	ccctctgttg	18480
gatgtaagta	agctatcttc	ttacttgctt	gctttgtttt	tcctttgtgt	agctctttaa	18540
agagtgtatt	cattcttttt	gtaagtgtgt	ttcttagaag	tagcattggt	gggtcgaagt	18600
gtgtatacat	tttacatttt	tgattgctaa	gctgcagaaa	agctgtattg	gtatgtaagt	18660
actcgtttcc	ttactatgct	cgtcatttct	agtgtctgct	cttcctttcc	ttcttcaa	18720
gggtttggtt	taattctagt	tgctactggt	ccatcagagg	aattgcagag	aactgggtctt	18780
caaaacagtg	cagtatatata	tttaggtgaa	gatacttcta	aaaacctttg	tattttgagg	18840
taattctaga	gtcccaagaa	tttgcaaaaa	gagtacattg	tcagcaatat	ttttcccaat	18900
ggtgacatct	taataatact	gtagcacagt	agcagaatca	ggaaattgtc	attgggtaag	18960
gtacttttta	attctccaaa	taattcagcc	ctccaaaaaa	atcccacttc	ttatgttttc	19020
aaacctgtag	ctacttttga	tgcgtaactc	ctaaattgca	tttttattac	tttaaaaaat	19080
ataataccta	gaagctcaaa	gctggaaaca	gcctgatcaa	tatagtactc	ttaagctaaa	19140
aacaacctga	tcaatatagt	actcttaggg	aaatcactta	tgctgtgggc	ttttttttaa	19200
ttttcttctc	gtcagctgtc	tcttcatgat	tttgtgggtt	ttattactgc	ttataccata	19260
gatgaggtat	agaaagtaaa	agaagttaaa	atgcattttt	ctcaatttag	tgaattaatg	19320
attacattca	gatttatagg	acaaggggtg	aagctacaag	gggttgatag	gaatcttgat	19380
gtatctgagt	attttcccca	actttattac	atgactgggt	cagactattt	tatctaatta	19440
catttcactc	ttggcagaaa	tagcaaaaaca	gtcaaccaat	ggtcaatgct	gctgagaact	19500
ctggcctgtg	cagacatatt	ggctgtttta	cttctaatac	cattctgctt	ttcctgtcct	19560
gctgctgatg	gatgtttctt	ccagggttta	aatatcaaac	aaaagggatc	tggtggccca	19620
gtacagggaa	tggtcttga	tagatttgat	tttctgcat	ttcctttatt	ttgatccagt	19680
gttaatttca	tgtagagttg	tctgtttaac	aggattctct	taaaattcct	tcttcagttt	19740
acctgccagc	ttttctttgt	ccagggttca	gtatgaactc	cactcgatta	atagagctct	19800
ctagtagtga	cttgtggagt	gggttctctg	aacattttctg	gaagtgttgc	tgatagtgat	19860
aatattgata	actagtactg	ttaatttggt	tgcttactac	atgttggtct	ttatatgtat	19920
tccttcagat	taaggacttc	tagaaaacat	ccatgaaaaa	acagattaaa	aaaaacaatt	19980
ctgcatgtat	ttgggactag	aaggtactat	gggaaggata	atcttcatac	tcagaccata	20040
ctgacctgaa	tttcatttat	cagtttagag	aaccacttcc	ccttcccttc	accctacctc	20100
cgagtgcctg	tgactttgta	tcaccgctct	ggcaccacat	cctcatccca	gcaggatttg	20160
ggaaggctgc	tttttgaaag	ccttttaaaa	ttctgtaagt	tgagaaaata	ctaggggaat	20220
gattttaaat	ttcttttagaa	ttacaggctt	tagtcagtat	atgacagagc	cttttcctag	20280
aaaaatgtgc	atataaaaaat	ttgcatgtag	ttttaggggt	tcagagaccc	ctaaagccta	20340
tcctatagag	tggttcattg	tctgattgtg	tttaggtacc	cttctaaaaa	ccttttgaga	20400
tgtttaggaat	cacaacagag	tatctctgaa	aatgtaatta	gcggaaagaa	catttcaaa	20460

actgttggtc	tgcttagact	ttctagtttg	tcttctgcc	ggcttgccg	aataaatgag	20520
tttcttggtc	tgatactcaa	aagaattgac	atttaaatta	gtctctctct	tccttggtt	20580
tcgcttgaca	catccttgte	tctacattct	gtctctgtct	ctgttagctt	atttctctct	20640
cgagtcagca	ggatatagtg	gctgttattt	ctcccccta	tccttcaacg	atctactttt	20700
gacaacactt	tgcccttttt	tttttgagat	ggagtttcac	tcttggtgcc	caggctgggt	20760
gtaatggtgc	aatctcagct	cactgcaacc	tttgccctcc	gggttcaagc	cattttcctg	20820
cctcagcctc	cagagtagct	gggattacag	acatgcacca	ccacgcctgg	ctaattttgt	20880
attttcagta	gagatgggtg	ttcaccatgt	tggtcaggct	ggctctgaac	tcctgacctc	20940
agggtgatctg	cctgcctcgg	cctcccaaag	tgagggtt	acaggcgtga	gccactgtgc	21000
cctgcctgct	atttgccctt	ttaatctcat	gaaatgttct	cttttcttgg	ctgaagtgtc	21060
acttttcttg	ttgaacagca	tgctgtgtga	gtagaatgtt	ataaaaagg	atggactttg	21120
gagtttagaga	gacccaggtt	cctgttcggc	attgcagaaa	tgctgttctg	caataggctg	21180
tgtgtcagtg	ggcaaattac	ttatctctca	gagccttatt	ggtaagggtg	gagttagagc	21240
tcctttcagg	caccttacag	aggctgtctc	ctaactcctg	tagcgtacct	ggctcataga	21300
tggcatttaa	aagtgggtgt	gatgacagtc	atagctcacc	attagcatag	cgctggatcc	21360
atggcaggga	agcgtgtcac	atgcagtatc	tcttggtact	cacagggccc	tcctgaatta	21420
ggaactgctg	tttcatgagg	atagggatga	ggaaattaga	cttgctgccc	ctcactgcct	21480
tcactcctc	tcctccaagt	taatgggaac	tctgactctg	ctttggcttg	attgccatgg	21540
aagattctca	cacagccaaa	tttattgtca	tattgttaa	attatgccag	aacacaaaat	21600
atgaagtatt	tgtaaaagta	atataatctc	agctgtaact	gagatagtca	gaaactgtct	21660
gtaactctgat	gtcctatctg	aaaggtagct	gagaataaac	aagaaataaa	gagaattcag	21720
tagcaaatat	tggtgacaca	aagcttttat	attttgacta	gttaagctag	ttcttaaatg	21780
tttccactaa	aatattcaag	tttaagggca	tagcccagg	cagcttatta	tgaacatgat	21840
gtatttttga	aatcttacac	tttctcttaa	aagtcttgg	gaggggcatg	tgaggccata	21900
atataacccat	aaaaccattt	gttttaaaat	aaaaccattt	tttaaaattc	ttccaaataa	21960
aaaaattatt	gcaggaaaaa	atgctaacc	tggtttttta	ctttgtacgc	caactatatt	22020
tccaagatgt	gctgtagcct	ggtaaccata	cagaaccata	cagaattagt	tctcagaatt	22080
tattgtctgc	ttacttttgc	atttggtaga	ggtataacag	ggtcgattat	atggtttcta	22140
agacatgact	agaaagaaat	atgtttatca	gttattattt	cttccatcta	aattagaagg	22200
ggctagggag	agggttcaa	caggaattta	tatactttag	agaaaagtga	tcattgatag	22260
cccaatagta	tagatatctc	aacccaataa	cacaggttgt	gtctgtctct	gggatcatac	22320
actgtagggg	agaatctttg	caagcaacat	tctacttata	gggagccata	acaaaagtgt	22380
catatgtata	ataattataa	gtcttaagtc	atcaagaaaa	agttaaactg	tgaatgataa	22440
tccttgatta	aaaagagaga	tgtataataa	tggaataag	atttttcttg	gttaattttt	22500
agtattaaaa	tggtctaaatc	ttctttggga	tattctgact	agtatgggtg	attgtctaat	22560
agattttcca	tagctgagag	ctaactcatc	tgtaactctg	ggaaaactgt	cctctttggc	22620
taaaacttta	ttgtaatcc	tctaaatcct	cagcttttat	tttctacaga	cttttttttt	22680
tttttaacat	tttcttctc	tgactcactc	ctttgttct	cattttcatg	gcctgagaac	22740
atgggtgatg	atagaattat	tcctttcaca	gattaacagt	tttcttttcg	agtatcgttg	22800
agctcatgtg	tgtattaact	agagaagtct	cccttacatt	tcatttttat	gttttcttct	22860
tcctcaggag	atagtttgta	gccatttact	ttcaaatcca	agtttctgcg	gttcttaaga	22920
cctgtatcat	ttgtctctg	aatttcactt	catttctctc	ttaaaccatg	tcctctgttt	22980
cccatcttct	gcacccactt	tgccacttcc	tgtttgttta	attggcaagg	gccactctct	23040
gtgttgga	ttttttcttt	ttgaaagctc	aactaacaac	ttctaggaag	ttttttattg	23100
ctactgttat	caattcatac	catcttacc	ttgtttttgc	aaccctttgt	taataacata	23160
tttatttaac	tatagttatt	agcagtctga	gatcatttta	cttggttaca	taaggagcac	23220
atatactctac	ccagcatcat	tgtaaggcat	gtgagacctt	tgtttgattg	ctgtcctaac	23280
ctagtaccga	gtcctaaaaa	ctcattagta	gaagatgaag	tgctcttgcc	ttttgctgaa	23340
catatatata	cacactgaat	atttagtggt	aattcatagt	tgcatgtggc	cattttttgt	23400
ttataatttc	ccctttctca	ttaaaaaac	tttgttttct	agacttttag	atttagagaa	23460
gtcatttttg	ttccatacac	atgctgctgt	tggtattttt	aggtattttg	tgactgtatt	23520
ttatctttga	aataaaaagc	ctttcaagaa	atgcaaaaaa	aaaaagctca	aaaaacagaa	23580
aatgtatatt	ttttaaatat	ctcagataga	tttaaagaaa	ttttaaacat	cctaatacata	23640
gtacttttga	agcccattca	tagtacaacc	tggtgaagag	ctcatgtacg	cgtaactggg	23700
gtcctgtctc	tgagtttgac	tggattgttg	ctgacatctt	ggccatcagg	cagaatgcgc	23760
taggacatgt	gcgctacgtg	ctgaaagaag	ggttaaaaatg	gctgcatttg	tatgggtgtt	23820
actttgtctca	ggttaactgt	ttccatgctt	ttctctctat	atatgtagtt	tataaatttt	23880
tttttttttt	tttgtagaca	gtctcacttt	attgtctcagg	ctgagtgcag	tgggtgtaac	23940
acagctcact	gcagccttga	cctctggggc	tcaagtgaac	ctcctgcctc	tgccctccaa	24000
gtagttggga	ccgtagtgc	caccatcatg	cccggctaaa	ttttctattt	ttttagagaa	24060
tgggggtctc	gctgtgttgc	ccaggctggt	cttggtactca	agcaatctgc	ctgtctcagc	24120

ctacaaaaat	gctggattat	agggtgtgaac	tgccatcccc	aaccctataa	aaatgttata	24180
ttttaaaatt	taacaatata	cttcatgtga	atgtatgggt	tttaaaatgg	gtttaatagt	24240
ttattctcag	ttgaagtaat	tttgtttggc	atttttagtg	gtgtgtattt	atatacgtct	24300
gattatccat	atpcggtttt	ccttcagcat	ctgtggggat	tggttttaga	accaccacag	24360
ataccaaaat	ctaaggtgtt	caagaccctc	atatagaatg	ggatagtatt	tgcatataac	24420
ctgtgcacta	ctttaaatca	tctctagatt	acttataata	tctaatacat	tataaatgcc	24480
atgtaaatgg	ttgtttatact	ttatttttta	tttgtattat	tttaattggt	atattatttt	24540
taatttttat	ttgttcacat	attttttgatc	tgtgatttgt	tgaatctgca	gatgtggaac	24600
tcattggatgt	gaagggccag	ctgcagtaaa	atgaaagagc	aaaaatgcaa	atgtacaaag	24660
ttcaaacaaa	taggaaattt	aaaggcatag	aatttgatag	gcaattacat	taaactgttg	24720
ataacagtaa	ttagtgtatct	gtatgatatt	aaaaaaaaaa	agcaaactgt	atatataaaa	24780
cttactttct	ccagttcttg	aggctagaca	tccaagatca	agggtgtgac	agggttaggt	24840
tctcccaagg	cctctctccc	aggcttgacg	acagcatcct	tcttctctgt	tcctcagggt	24900
gtttttttcc	ctgtgcccac	gcaccctggg	cactgcttcc	tcttcttaga	aggactaggt	24960
gactggatg	actaatcctt	ctacagagac	tgctaaggtc	ccactctgag	gccctttttt	25020
aaccttaatt	accacctcta	agtcctctct	tctgaataca	gtcacagtgg	gaactattag	25080
ggcttttagta	gactgatttg	ggggaacaca	cttctgtccg	taacagtggc	acataaatat	25140
cttttagcagg	attgattttt	taaaatccct	aaagatcgtg	agtattgaca	tgtaaggac	25200
gctttttagt	gactctgtaa	taagtgggtg	gaagaattgg	gagttaaate	catctgatgg	25260
atcaggtttt	ttatttttta	aaatgtgtat	ttaaagaaaga	aagcattttc	attttaactg	25320
ccaacaaaac	taaacttcat	gtgttttcca	atacagtgtc	acatgcagtt	tttttgaatt	25380
atgttgagac	aaggcaattt	tcagctaaat	gttctttaga	agctaattgt	tgaagatatt	25440
aaatatagat	taaattctga	aatgtagttt	tcattctgta	ctttttgcaa	gagaagttgc	25500
ctttttgatg	actctggcca	attgttattt	taaaagtaaa	tgctctttct	cccgatttga	25560
ttgtggcagc	atggaggaat	ctatgtaaag	cgcagtggca	aatttaacga	gaaagagatg	25620
cgaacaaagt	tgacagacta	cgtggacgca	ggaactccag	taagagccta	cccgttttta	25680
tttttcttac	cagctctcag	tttctaaatt	taagaattaa	attaaaatct	aagaattgtt	25740
ttgacaatgt	attttcccat	gtgttaattac	taattcaggg	ttatgctgag	gtaacagaaa	25800
ccctctatgt	acaggtaggc	aggtttttca	gccatcagaa	agattgctgt	aaacaactag	25860
gtcctttgct	ggcagtgga	ccttaaagag	gaataaaaag	agcatttggt	gtcgttcaga	25920
gtctataaat	agaaactaat	gcattttaac	ctgacattta	agctagttta	caagctcatc	25980
ttacttcttg	tcttctttag	tatcagattt	ggtttttaga	gcagcaactg	ttttctgtta	26040
gtgcaaattt	tgaatgtctt	acatgtacag	aaaaaccaaa	aaaggatgaa	tctctacaaa	26100
tgttaaatca	ttcagtgtaa	ataatatttt	ataaaacttt	attccacaaa	agtggggaga	26160
gttcaaatctg	ctttgtatag	aatgctgatt	gctgccaaaag	gcttttcccc	tggttccctc	26220
cggagacaaa	gcaccatgat	caccggggcg	acttgggctt	tctctttcag	tacatgacat	26280
gtgctcagaa	gcttagctcg	tgtgcacagg	ctttcccttt	cctttctggc	tcctccctc	26340
tgtcttccct	cctctcctct	tgccctcccc	tcaccagggg	tcctgggcag	cagctggagc	26400
tcattggtgaa	ggaagaattc	ttcatggtca	gctggcgaag	tgctgtgtgt	gagcattgtt	26460
tattcacatg	cctctcttag	gtgtttttac	attagaacat	tgcatctgtt	ttgggcattg	26520
gttggtgtgac	agaagcagaa	tggaatgaga	tgaacagtga	ccctttatcc	tgttatagct	26580
aacccttgag	aaccaagctt	gggtgtctca	aagggtctgt	ttagtctgaa	acagtgtggt	26640
gaatttgggc	agaattgtgg	tcattgcatg	taggtctcca	aaagacagaa	taagttggta	26700
atatggttta	tgcacttttt	acaaaaaaa	tttaaaaatc	atgaatttat	accttaaaat	26760
gtccatccca	cttctctccc	agctgtccag	tcacccacag	aatggatgac	tgctgtggag	26820
ttcctttctgt	gtcctgctgt	gggcattgta	tatatgaagc	aaatgaagat	agctgccttt	26880
tggtgtatgt	tgccatccca	tgacacagtgg	tcccttgctt	ttttgcccc	atgaatatag	26940
ctgccagtgg	cgctagggct	gaaaaaatca	gctctttaca	cttgctcatgt	gtcttgttta	27000
tgtggctgcc	ttcgtgagtt	tcttcttggt	tttgggttgc	agcagtttaa	gtatcatata	27060
tctgagtgtc	atttaaaaaat	ttttacctgg	attggctctc	tgagcttgga	tctatgattt	27120
gggtgtctgtt	attaattttg	gaaattttct	tgtctttatt	tccttaata	ttattcctac	27180
ccagctcttt	cttctccagt	tatgtttgtg	ttgggttcatt	tctcgtgtgt	ctttagttct	27240
tagatgcatt	attcgttttt	tgttggtttt	tttttaaat	ttttttttta	cgtccctctc	27300
cttttttctt	tttgtgttac	attttggata	atttctgttg	acctaccttt	gagttcatgg	27360
attcttccct	tggctgtgtt	gagttctactg	gtgagccagt	ttaaggcact	cttcatctct	27420
gctactgctg	gtttcattcc	tcacatttcc	ctttgacctt	gtttcatagt	ttccatctct	27480
gtgctagtgt	atctatctga	tcataaagct	tagtccagtt	ttccagttga	acctttatca	27540
ttttattata	cttgacagtc	tcttaaatcc	cctgcttgat	aattccaaca	tctgggccat	27600
atctgagtct	gcpaaattttg	attactttat	ctcttcagat	tgtgctttat	cttgcccttg	27660
tcatacttcc	taagattttg	cctaacgctg	ggcctttttt	gtaagacagg	agaaatggag	27720
gcaagttgtc	ttgatacctg	gaaatggata	gacttgtctt	tctgcttggc	tttttagtgt	27780

gaggagtggga	gtcagtcac	tgaggaggtg	cactgcattt	gggttttgc	catgtgcttt	27840
ttctcacagc	ttcaggtttc	tgtagaactc	attactttgt	ttgtaggttg	gggatgtcct	27900
cccgctagag	cttttctca	gtgtctat	cacactcagc	gttttcacat	agcaccttgg	27960
agtggctctc	ttctttatgc	ctttccccc	tatacttctt	ggatacttgt	tactgaactc	28020
tcgctagttt	ggtggtagaa	ggagagggaa	gggaagtgtc	ttttcattct	tagggagaat	28080
ctcaggggtg	gagccttctc	tgatcctgcc	ttgttctctg	ctgtaagtct	gtgcccagta	28140
tgtattcctg	cctttactaa	gagtttttcc	ctgttctctt	caccagcct	catcgagtat	28200
tcattccgtgc	cccatgggta	gcagggtttt	gttgcccctg	ttcatcagtt	tcaggctgct	28260
gttccatagg	aaaggtagaa	agaaggatgt	gggctgggccc	ctgagccctt	cccacagggc	28320
tgctttttccc	tcccacaagc	ctacatccag	tcttccctga	cgcagtggtg	ttttcttttt	28380
tctttgtctt	gtgagtacac	aggaggtctg	tgggtcagagc	ctgtgaaatg	tgctgcattc	28440
tccttgtgtc	tgtagcccag	gggttcgtct	gttccactgg	ctcatacttg	gctttctgca	28500
aaattgataa	aatttttagc	taaattcttt	ttactgggat	ctgttacatt	ggcccccaac	28560
taaacaacca	cttgcattct	gtttctcctt	tgagttttcc	atctttcctt	agacttttgg	28620
gttagttggg	tgcttgcaa	ccttgcagct	ctctgaaggg	tctaagaaaa	gtcatgaatc	28680
tacagcttgt	cagtgttgtt	gttgtttag	gggtggcagt	agtattcctt	cagcattcta	28740
catacttaat	ggaagcggcc	tcccattttt	ggttaataaa	tttcaaaact	tggaacaatg	28800
ttagatttcc	aaaaacgtca	gaaagaacag	agtgttccctg	tttattcttt	atatagcttt	28860
tttttttttt	tttttttttg	agttggagtc	tcgggtctgtc	accagggctg	gagtgcagtg	28920
gcacgatctt	ggctcactgc	aacctctgcc	tcacgggttc	aagcaatctc	ctgcctcagc	28980
ctcctgagta	gctgggatta	caggcgtgca	ccgccatgcc	cggctaattt	ttgtattttt	29040
agtagagaca	gggtttcacc	atgttggcca	ggctgggtctc	gaactcctga	cctcttgatc	29100
cgccccctc	ggccccccac	agtgtctggga	ttataggtgt	gagccaccac	gcccagcctt	29160
cttcattctag	ctttaacatc	taatgttgac	atctttacata	acatgggata	tatttgcata	29220
aactaagaaa	taaacatttg	taccacacta	ttaattgtac	tacagatttt	tattcagact	29280
ttaccaggtt	ttccactaat	gtcctttttc	tgttctaaaa	tacaatccag	aatagataca	29340
aatccattca	acttcagtgt	tttaaattat	tgtttttcat	tatatgaagt	gctgtgtggg	29400
ttttgtcaaa	tctgttattt	tggttttaat	cttcaagctt	gtctttgttt	ctttaagtga	29460
taaaggcata	atttaaaagg	tgtgttgggt	tatttctagt	cctaaagtct	tgtctgagtc	29520
acttgttttc	tgtgttctt	gcttatggta	ctttctttcc	ttgtttgctt	tgttatcttc	29580
ctttgtctgt	ggctgtgttt	ggttaagtta	tttgtggaaa	tcagttgaag	cctcaggtgg	29640
gagtgtcttt	ctccggagaa	catttctacc	tgttttagct	gggcccccta	aggctcctct	29700
agcgtggggc	ccacccaaac	gagattctga	gttgaagggtg	aactgagcca	ttcaggcagt	29760
gcagccaggg	ttgcagatgc	acgtgagacc	tgctcacctc	tcatttactt	tcaccctgag	29820
agtagagcct	ttgggtgttc	gttcacttgt	ctgattctct	cttcacagtt	ctattagaag	29880
gtccatgggt	tttggtttct	gtgcccttca	tcttatgagt	cttgtaaact	aaagttctgt	29940
tttatgctta	cttctgcttt	actgtgtttg	cttaatttca	gtcttaacat	cttgccaact	30000
cttgggtact	tttaaaataa	tgttatatcc	agctttttta	gttgttttca	gtaggaagg	30060
tgattcaaat	aacctagtct	ggttatgggc	tacgagaata	gcttccctgt	ttttgtggg	30120
caaaattcca	gccttttatg	ttcctagcgc	agtgtggata	acagactggc	aggttcaaga	30180
ggccgtgctg	agcagctttc	actgtaagg	cactgtccca	ggctgggttt	ctaagaatct	30240
ggatgggtgt	ttcatttctt	aatatgtacg	ccctgtgaga	gcggatacat	cttgctcagg	30300
ttcttatgat	tcttttggtt	ctgaagggtg	atgaagtaag	tgacatggta	gaatatgtta	30360
agtcaacttt	cgtgtggctt	actagttctc	atgaatctat	tccatgattg	tatcagttct	30420
tattcagtat	tagtatttaa	gaaatgcaga	attttgtttc	aaaaaatata	tttgtattat	30480
aagtgttgaa	gaaatacatc	tccataatta	ttgctgggac	aatacagtat	tttcttaagg	30540
aacttattgg	ttgtggatgc	aatgaagca	tatttgtgat	aaaaataact	aatagaagtc	30600
attttgttag	actatgagct	agtaaaactt	atggcacaaa	catggagact	taacactttt	30660
tcttccagct	ttcacttaag	ttccttttca	gataggaggc	agcctgggtg	ataagagtat	30720
tggttttgaa	attagattca	ggttttaaat	ccagatcttc	tgtttaatct	ttattttatt	30780
tcaggtagat	tttctggata	acttgctata	gcttatacgt	cagtacttgc	cacttcaatt	30840
ttatgttatg	gagagacggc	ttctttcctt	aaacctcacg	aaccaacctc	tgctagcttc	30900
taagtttttt	cctgccactt	ctttacctct	ctcagccttc	agagaattaa	agggagttag	30960
ggccttgctc	tggattagga	tttgctttta	gggagtgttg	tggctgggtt	gatgttttat	31020
ctagagcact	caaactttct	ccatatcagc	aataaggctg	ttttgctttc	taatcatcca	31080
tgtgttcagt	gaagtacac	ttttaattct	ctttaagaac	ttttcctttg	catccgcaac	31140
ttggctgttt	agtggaaagg	acctagcttt	tgacctacct	tggctttcaa	cataccttcc	31200
tcactaagcc	atcttctagt	attgatgtaa	agtgaagagc	atgcaactct	tcctttcact	31260
ggaacgctta	gcagccattg	tagggttatt	aattggccta	atctcaatat	tgttgtgtct	31320
cagggaatag	ggaaacccaa	ggggcggtag	agagaaagag	agacaggaga	acaggccatc	31380
attggagcag	tcagaacaca	cacgacattt	atcaattaaa	tttgtcatct	tatatgggtg	31440

caattcatgg	cacccccaaa	caattacaat	agtaacatca	gagatcacag	atcacataaa	31500
cagatataat	aatatgaaat	attgtgagat	taccgaaata	tgacacagag	acgtgagggtg	31560
agcacatact	gttggaaaaa	tggcaccaat	agacttgctc	gatgcagggt	tgccataaac	31620
cttcaatggg	aaaaaaatgc	aatttccgtg	aagctcagta	aagcgaagca	tgataaaatg	31680
agatgagcct	gtcactccta	agaatgttcc	tgtacaagtt	ttttgcatct	gttacttacc	31740
ttttcctatt	tgtgaatagt	atcttttttg	agtacgtgtg	tttttttatt	tttatacatt	31800
tatatgtatc	ttttgaagaa	catactttta	agcttaattt	attgattttt	tttctctcat	31860
aatttccact	ttttgtatcc	tatttaagaa	gtccttgcca	aacttaaggt	tgctaagatt	31920
ttctcctttg	ttttcttctg	gaaattttag	agttttgctt	ttacatttag	ttctaggatt	31980
tatttataat	taatgttttc	atatgggtga	agatcgaagt	tcataatttt	ttaatatagg	32040
taaccatcac	tatagaaaag	attatttccc	cccaatgttt	gaaataagta	gactgaatat	32100
agatgggtct	gttatcccta	gatcaatgga	gcatttggtc	tgttatattg	atctatatat	32160
atatatcctt	atgccaatac	catactgtct	taataatgct	tgctttgcag	taagttttta	32220
aatagtgtag	ttgtcttcta	aatttgttct	ttcttttcaa	agttgttttg	gctatttttag	32280
gttttttgca	tttctgtgtg	aattatagaa	ttagctcgac	aatttctacc	caaagtttgt	32340
gggtttttca	ttttgattgt	attgaagata	tagatgaatt	tggaagaat	tgatataaca	32400
ggattgaatc	tttggattca	tgaacgtagc	ctgcatttgt	ttacttaggt	cttctttatt	32460
tatctcagtg	tgtttttagt	tttaatgtac	agatttgcac	atcttttgcc	agatatatcc	32520
ctaagaattt	cagtttttga	tactattgta	gatgacattt	aaaaaaattt	caagtttttg	32580
tttgttgacc	taggcatata	tttgactttt	taataacta	accttgctaa	acttatttat	32640
catctagtaa	cttacaaaat	atattcctta	ggatttctta	cataaacaat	catgtcattg	32700
ttttagaaat	aacagtttta	ctttgtcctt	tttaattctg	atggctttta	tttctttttc	32760
ttgctaaatt	ttctggctag	acctcctagt	acagccttga	ctagaactgg	tgtgagggaa	32820
atcctttcca	tattcctcat	ctttagggaa	aagcactcat	tcttttatcc	attctttagt	32880
tcttagcccc	attgcccttc	ctaaattttt	tctcatcatt	ttccttcac	acaccttggt	32940
ctttttcttt	gcaatcatat	catgatattg	aacgacatgt	ttttatttat	ctgtttaatg	33000
tatttctttt	cctcacttgt	ccatgaaggg	aaggaccata	tgtgttggtt	tcctttgtgc	33060
agttcctgga	acataataag	tatataagaa	atagtttctg	aattagctgt	gaatgaattc	33120
atgccttctt	gctgtctgtc	aatgttcttt	taaattaaac	atctaagaca	gcaaataata	33180
ccacatgagt	tattaacctg	agaaataatc	gttttattta	taaatgactg	agttgaaagc	33240
tgatagccca	cagtaattgc	tttcatggct	ttgaatataa	accttactgt	tacaaaacac	33300
attttcatga	aaatgaatgt	gtgggtgttg	gaactagctt	taatgtttgt	cttctgtttt	33360
ttccttctag	ttgtatataa	ataataagga	attttgtatg	tttttcttaa	ttgtaccac	33420
ttttctacat	tttcttaaca	gatctggtga	atcttcatta	ttaaatataa	ttatacatat	33480
aaattattgt	ttaataataa	tattaattat	taaaaataat	ataaattatt	aaatataaag	33540
atacatataa	tattatctgt	taatttctaa	gttaggtgtg	ggttctgaag	actattatat	33600
gaatgaacaa	aaagcttgca	tatttgcgtg	gaagctgaaa	gtacgaaatt	tttagatacc	33660
attataccag	tatctaaaga	aaaaattcag	taccacatag	gtttttaagt	aggagctgta	33720
tgatcatagg	tcattccagat	gaaggaaggc	ttctgtacca	gacgtacaga	ggtagacagt	33780
gttgctgtag	tactgtctga	gatctggcaa	gaatgaatcc	aataaacgta	gttttctccc	33840
atgagctcct	gttctgttct	ctgtattctg	tttgtatttg	aaaagatttg	gtgtgcataa	33900
cttatttttg	tcttttggtt	gtcaatcaaa	gttatttagt	tagtttttgt	aactcagttc	33960
tcaagctagg	agtttttgct	gtataatttt	aatgtttctg	tttttacttt	cctaagcaga	34020
taagcgtaaa	aapttagact	aattgattac	ttattaaacg	tccagcttga	tattcttctt	34080
tatattattt	tagtttcagt	ttatataaca	aatgaggttt	cttataaata	aaatttaaaa	34140
tgcactaaag	gagctgtgtg	aaataggaat	tctgtgtgaa	gcttttgaat	gtgaacattt	34200
agaacgtttc	acatgggtgg	aatttactat	atgattttca	tcaaagagg	tacttttttag	34260
tgttggtact	taacgatact	gatttctaaa	atttgtattt	ctaaaaatga	cgtattacag	34320
gatctgaaag	ggcaaaaact	cattgaggct	ttgtatgagt	cagcgtttca	tggcctattt	34380
ttaattagtg	aattattagc	atataattag	aaatgttttt	agattcttca	tggctgacct	34440
accaatgaat	gtagcactgc	atttaaaata	tagttcacgt	tatgttcata	cttaattgtt	34500
gcattttgtt	tgtccctctt	gaaacgaagg	tcacatgtaa	ataaatatac	attttctcct	34560
actgtaggaa	atactctgtt	agcattagta	ggtttagctt	tttttaggtt	acaataacaa	34620
aaacaaagct	cacacaaaat	aaaccaaatt	tgctctatgt	cccacagatg	tatcttgtga	34680
tttttccaga	aggtacaagg	tataatccag	agcaaacaaa	agtcctttca	gctagtcagg	34740
caattgtctg	ccaacgtggt	aagtaaaaaa	ttgagtgttt	gaacaaataa	ttttcaaaga	34800
taataacatt	tttagttttt	cttctgtgaa	aagatacttt	tgttttacag	ttgaagggaat	34860
gaatgtattc	attccttgaa	ttagtgtaca	tattatctct	taggaaatga	agtttcttct	34920
ccttaattca	ctttcatgct	attattacat	atatctgaga	aattaagttg	aagtgtctgt	34980
tacgatacat	attcttgtgc	catggattta	tttaaaatct	atctaagtac	atgattatgt	35040
agatggaagc	tttttctaca	gtgtatgggt	tatatgtaat	ggagcttctg	ttttgtaaga	35100



tgacagacct	aagttggagt	ccaaactcgt	acttttatta	gctgtatggt	tgcaacttgg	35160
aagttgtgta	atgttgctga	gcttgcttct	tcctctctta	aaagaacata	tgcccttataa	35220
gtagatctaa	atctgtgtga	ggattagatt	agaaaatatg	tcaagtttct	attggagaag	35280
ttacacaaag	ttgggtccaca	gtgcttggaa	gctgttaatg	tcttcaacaa	tggtaatggt	35340
cttaatatcc	atattttaga	aaattgaata	attggtacac	caataagcta	tgcaatttaa	35400
ccaaattggg	aagtatacag	aaaacagtgg	ctatgctatg	ttcttagagg	tgtctttgaa	35460
gcttgactgt	gatttagtgt	gtgatctcca	tatgttgata	gtcactcact	gagcaaatat	35520
cttggttggt	acattacagc	agggcctatg	acagtgtgtg	ctaattggaac	tttctgcaat	35580
aatggtaaa	ttcttcatct	gttctgtcca	gtgtgctggc	tcctaccaat	gtggtttttg	35640
agcattcaac	atgtgactag	tgcatgaaac	taatttttaa	ttttatttaa	ttttagttaa	35700
attaaaaata	agggggagt	tttacaagg	gcttacaaga	gcagatatgt	cataggata	35760
tgacatcatt	tgtaacagta	cttttaaaaa	atgccagttt	gtttttaaac	acatgtccta	35820
ttaagtaagg	agtgtctcag	aataggagg	ttcagttggt	ctccccatct	gccagctctc	35880
ttttgacttt	cattgtcttc	tctgtcta	agacatgacg	ttctgtcatt	tcagttgctc	35940
ttttgcaatg	ccattgtctc	ttttgccctt	ttcacattta	ttaaacagaa	caaaacaaaa	36000
accactctcg	aatctgtagt	ctacctttgt	tgtaagcact	ttttccagta	ctcactctgc	36060
cctcaatttg	ttttgtctg	atttgaatt	ctctccctag	acttctgtgg	ggctgtctc	36120
cattatcctc	ccaactctct	ggcgattact	tcctagcctc	ctttccagcc	tctttctgct	36180
tcatttctcc	ctgtacatg	tgttatttcc	agtgtcagg	tttgggtgtt	gattaatttc	36240
actttttggt	tctcatgggt	gccttccctc	aaatccatgg	cttttagccat	cgtttccctg	36300
actgtgatg	actcgcaaaa	gcttctctcc	ctccatgtct	ctctgcctaa	ctctggaccc	36360
atttgtacaa	ttgtccatta	gagagcttcg	cttgactggc	ccaaaaggat	gtctcaaac	36420
cagcatattg	aagatagaat	ttatccttcc	atgcatacac	tcataatttct	tgtcttggt	36480
actccatcat	tcagtttttt	tgccctaagt	ttattcacaa	aaagaacaaa	ttgatagcag	36540
ttgcatacct	cttataggaa	acttagacat	gtgggaagaa	gctgttcaga	tggggtcctg	36600
cagaagtgc	ggcactgtgg	taatatttaa	acttttctca	gctgttcgaa	gggttttgtt	36660
ttactaatt	ttccttagac	ttgttttagg	tatttgggtt	tctaattggt	ataagggatg	36720
tggaattaaa	tgtatcttaa	tctgccacct	ggacccatta	aagtaagccc	ctatggtggt	36780
tttttttttt	aattgccatg	gttaaaacca	tagttgctag	cgaaggtgac	atacttaagc	36840
tttttgaact	ctcttaaaag	aaaacagaaa	tttaatgatg	tgtctataat	ggcaaaccag	36900
atacctagaa	ttcccatggt	attcatagg	tgaataacac	tggtcgattgt	agagatttga	36960
gagttctttc	aaaacaggag	aacaaaggga	ataagctaca	aagcaatttt	tttctttgta	37020
gacttaactg	aataaaaaat	atttttatgt	ctcaaacatc	atatgaacaa	atttagttgg	37080
caaattggcaa	gctaataata	ttttataata	taggatatta	atatacttaa	tattacaaaa	37140
gtgcttcata	attagaaaag	acataaaacta	gaaaaatggg	aaaagggcat	gaataagaaa	37200
ttcaagagat	acaaatgacc	cacacacttg	aacaaatgtt	tattctttct	cataatcaaa	37260
gaagtagaaa	ttaaatgaat	actttgaagc	caacttctga	gaaagcatag	caaacaagaa	37320
agctagtgtc	cagctttgtg	tggtaacggc	actctcgctc	ttaagaaggt	gtgtttgctc	37380
cctgtggctg	ctctcaggca	gggccacaaa	cttggtggct	taaaacacca	cagatttctt	37440
ctcttacatt	tgagaagtct	gaaatgggtc	ttactcagct	gaaatcaagg	tgttggcagg	37500
gctgcagtcc	tttgtggagg	cttgggggga	tcttgttctc	ctgtacgggg	tctgtgtgct	37560
gggttcgggt	cctgtgcttg	gtctggggtc	ctgtgcttgg	ttcgaggtcc	tgtgtgtggg	37620
ccagtgtctc	gctttttacca	ccttgaagtt	catctggaaa	tggtcactgg	tcgcccacac	37680
catatagctg	actctggttc	tccctcctcc	tcactcgctc	taaacctgtg	tttttggtg	37740
atttctaata	tctcttttct	tggcccttct	gcagcttgca	gggccttctg	cagctcttgt	37800
ctgccccagc	ccgggggtct	gccccccca	gtgctgggct	gttctgttcc	tgccctgctt	37860
ttcctcagcc	cttggcaacc	ctgtttgttt	tctcccttcc	ttagcagttg	agaacatcgt	37920
aagatcaatg	ctgactgcct	tctgcagcca	agccaggcca	tttcatttca	gccgagccaa	37980
gtctgtgtgg	agcagttctt	ttatttttct	ccttttgact	acctcatggt	tttcacggat	38040
ttttgttctc	ttcacattca	aggatttttt	gctttcagaa	agttatatatt	ctctggaaag	38100
agtgcaccca	atataccctt	tgatttcaaa	atcttaatgt	ggagtctctt	gacttggatt	38160
tctttggaag	aaactgctga	agctgccatg	tctaagaaga	aaactttgga	gaaaaatttt	38220
cttcttagac	atggcaacgt	caacagtttc	taagctcttg	attccgtcta	ccctgtctcc	38280
atcgttgcc	cagtcactct	ccttacttct	ctgcaggggt	ttctcccagc	ttgcaaatgt	38340
actccaattc	tgaataaact	aagtctatag	ctgtgcaaa	agaagtctgg	gccccttgct	38400
ttcttgtgtt	tgactccatc	cactctccag	aaatgaatcc	cacttctcac	ttaaccactg	38460
acctccaaag	catcgtatca	tttgtgtcag	ttgtcatatt	tgttaaacttt	cacataactt	38520
ttgacattat	ttataccttt	ataaccagga	aataatttta	actttattgt	agaaataaac	38580
aatggagtat	aatttttctt	gttgaagata	aatatcacct	cctcttctct	taaacatctc	38640
ttccctttgt	tttgttatta	cattggtttc	cccccttttt	ttatttctct	gggtgtcgta	38700
ttccctgtta	ttattttttac	cttttttttt	ttaatgtgga	tgtttccgga	gtctgtattt	38760

cttgccctttt	catcttctgc	cctttattat	tctcagccac	tgccattact	tcagttatcc	38820
attcccatgg	tttccacatg	cttagcttcg	gttgattctt	gccattttac	agaccatatt	38880
tccaactact	tctagaatgt	tttggtccct	cagcctcagt	atgcccattt	tgaactcatg	38940
ttctctctcc	cccttctttc	ttccttcttt	ctttcgctct	ctctcccttc	cttcttttct	39000
ttccctccct	ccctttcttc	cttccctcac	tcgttctctc	ttgcttgctt	gctttctctc	39060
ctctctctct	tttctttctg	ctttctttct	attcttctcc	ctccctctct	tccttctctc	39120
ccccactccc	caacttccag	gctaaagcag	tcctcctgag	tagttaggac	tacagacata	39180
cacgtgccac	cgggccccagc	tccgtgttct	ctttgtttcc	ctgcctctcg	ctcttccact	39240
tatcttttga	tggcaggtgg	gtgcacgcag	gcattgctctg	catgtcttcc	tcttggccat	39300
tcccttctta	gttatgggtg	ggctttatct	acgcgttctg	gagcagaagc	ctagtccaaa	39360
agctattttt	ttaaaacatt	catgataatt	catttctctt	tatgttttaa	aaatactagc	39420
tttctgtctt	tatttctctta	ctaacttact	tggatgccag	taattagttg	ttttagttaa	39480
caccacagag	tgatattttg	aaacttttga	cttcataaag	ttggatgagc	tccagttaga	39540
aagaaggaag	tgttaactag	tttaactgac	aaataaatgc	ttcccagctt	gggtgtgcgag	39600
tgagattttt	gttgcaagtt	tgtgaatcaa	tttaactgcc	cctgccctgg	ggactaaagt	39660
cagatacgtg	cttgtgggaa	tctttgtctt	tcccacacca	ccctgcattt	taaaaacctt	39720
tgtgtgggac	agtcccacca	tgtaatagct	gttcttctct	actcagctac	tttccctcca	39780
gagaggccag	tagaaaatct	agactagttt	tttatagctt	attttcatgt	cacttattga	39840
gagctactgt	tttctgttaa	attgtcagta	aatattttaa	tcaaggaaaa	gggagggaat	39900
aggaaggaga	gaagaacaaa	tccttaaccc	tagtggaaac	ctaataaatg	ggatttgttc	39960
tggataattg	cagtagtccc	ccagctaaag	aaccttttaa	aaatatgtca	gatataccca	40020
agaggattga	aatcgtatgt	tcatacaaaa	gcttgttcac	ctgcagcctt	catatgcaat	40080
tcctatgaat	gttcataagca	gcattattca	taatagccaa	agtatggatg	caaccctaat	40140
gtccatgaag	caattaatag	gtaaacaaaa	tgtgatctgt	tcacacagtg	gaatactaac	40200
tattcagcca	taaaaaggaa	tgaagcactg	agtcctgcag	ccacacagat	gaacctcaga	40260
tccatgctga	gcgaaagaag	ccagaaacag	gaggccatgt	gctgtgtgac	tgtatttcta	40320
ggaaatcttg	agtcaccatg	ggcaagatgc	tatcaccttt	gttcagtggc	cagaagcgag	40380
ggcactaata	tttacccttg	ccgggttcta	ctagattgaa	gcgtttccgc	taggccataa	40440
acttccaaca	cggtgacttg	tacatgtaga	tatttgatca	atatatagca	aatgaatatt	40500
gatttaaaca	gaaaaaggca	agtgaagtg	ctttctaaac	ttagagccct	aaatatatga	40560
ggttgtggaa	ttaatagatt	ctgttgtgtg	tgtttgaggg	aatttaaaaa	taatttagat	40620
gttaaacagt	atattgtgga	ggtgttttgt	aactaattaa	tgacggcact	gaattgactt	40680
ctaggccttg	cagtattaaa	acatgtgcta	acaccacgaa	taaaggcaac	tcacgttgct	40740
tttgattgca	tgaagaatta	tttagatgca	atttatgatg	ttacgggtgg	ttatgaaggg	40800
aaagacgatg	gagggcagcg	aagagagtca	ccgaccatga	cgggtaagtg	tgttcacgca	40860
cctgaaatgc	ctgtacacgg	tatatacagt	gcacatgttt	atgtagaatt	cagttttaca	40920
aagtaggtta	agtgtacttt	tttctctcat	tacatttacc	cggatatattt	ttcaagatgt	40980
tattaagatg	taacagtgga	gatttcatta	gtcctgcaaa	gtgtgggtatt	tcttggctgt	41040
cgtgtgagtc	ctgtggactc	accaattatc	attaatccag	cctctttcta	ctcaaagttc	41100
acacttaaaa	ggaaagctct	gtaaaaggga	ggaagacgtg	aagaaggagc	acgcctggca	41160
gtactgagtg	caggttatta	gtcagtgtcg	cccttttctg	gtatttttctg	taaaaatttt	41220
attaaatttg	gctgtcattg	tgacaagaag	aaatgcagtt	aagtgtgacc	tttttttttc	41280
cccaaacatg	ttaggtttta	agaacctttg	agctattgtc	agatataacc	agaaaaaaat	41340
agaattttta	gtgagcagga	taacttagtt	aaactaacca	aacatagttg	tagctgttag	41400
agaaatgtaa	acatggaaat	aggcaaacag	ggaagtgtgt	ggagtttctg	tttcttttct	41460
aaaatatctg	tttgagctgg	ggttgagaga	gaacactagg	cttcatgggg	tttttttggt	41520
tttctgtttt	tgttttgaga	caagagtttc	gctctgtcgc	ccaggctgga	gtgcagtggc	41580
gcaatcttgg	ctcactgcaa	cctccgcctc	ccacgttcac	acgattctcc	tgcttagctc	41640
tcctgagtag	ctggaactac	atgcgtgtgc	caccatgcat	gactaatatt	tgtattttta	41700
gtagatatgg	gatttcacct	tgttggccag	gctggctctca	aactccttac	ctcaggtgat	41760
ccacgcacct	cggcctccca	aatgagcttt	gtgtttttac	ctcatcagct	gtttgggggt	41820
gagccactat	gtatgtcagt	gtgcttgtat	cagtaggatc	tactgagggc	agatgttcaa	41880
aatatgagcc	tccagcacgt	tttcatgga	aaccttcacc	tgaagcattc	gtctgaagtt	41940
gatgtgcctt	ggaaattttta	tagagtaata	tttttaacta	caacaaaaca	tttataaaaag	42000
tagacattat	tgaagcattc	agaagtgagc	aaggatagaa	attattctgc	ccaaccttac	42060
acgtaggcct	tctagacgta	gtactgtgca	ccgttacatt	atctaacact	gtctgtgtgt	42120
catcttttga	tgttagggat	ttttccaaag	ttatagtgtg	ttatagtgtg	caaatgatta	42180
gtctgtttaa	taatgataag	atgaggggtca	ctcaggtttt	aaaagaaaag	ctctttgact	42240
gaaagagaga	gcagctgtct	actgcagaaa	gttagggagg	gaggctggag	gagtgaggcc	42300
caggggctag	ctagtataaa	aattgggttat	ggtcgaagga	aaaaaaaatg	taacatattt	42360
atatctgaaa	gatgattgtt	ctcataattg	tatataacac	agagtaattg	taaagtagaa	42420

aactaagggtg	tttttcattt	tagatgtaaa	tgtttagaat	atgtaatgca	tcagttttaa	42480
aattaaaact	gtacgaaatg	cacagtga	cgtcttcctt	gctttccacc	ctgctacctg	42540
gccttccctt	ctccttccta	gcgataacca	gttttcttaa	tttggtgtgc	gttgtagtgc	42600
caaatttaaag	tatatcttct	tattctacca	tccttccctt	cttacagaaa	agtggcatat	42660
taatatTTTT	ctctttttaa	ctatcgaagg	agttacttac	ctatttttgc	atttcaaaac	42720
agacagttca	tcaagattgt	cgttggttta	ttaaacaatag	tttaagatta	aacaagtgtt	42780
tataaccaat	gaaaaacaga	tagactcccc	ataataacct	tgtttaaagt	ctgctacttt	42840
tatcatgtcc	cctcctgtct	aagaacccct	tggttcagca	gagctcatgg	gtaaggccag	42900
cctctgttgc	ctgccatcgg	aggaatgcgt	tcagcccggt	atctctgcct	tgccctcgct	42960
tcctcctgtg	ctgtgccgtg	aagcctcgcc	cgtgggtgaag	ctggctgact	gagtcctcct	43020
gcaccccatg	catattcagt	agttgaaggc	ttgtgtgtgc	caatcctgct	ttccacagga	43080
aaccaccctc	tcttttgttg	ccctcatcca	aggctactgt	tctcccagag	tgacaggcgg	43140
cacctttccc	agcatagcac	tgtgccttct	cctgccccctg	ctcttgcaat	actgctgtgg	43200
cactgatggc	gtgtgttaca	gtgctggcac	ttagcacagg	gctctgcctt	tctctcttcc	43260
cagccgcac	ataagtgcct	tgaggaagcc	aaaaccttct	gtgagttgca	ttgcctgggt	43320
tccaacctcc	cactgccctg	cttatcctct	gctacatgtg	agctgactgt	ggctttgggg	43380
tggtcactgc	ctatgtgtat	tcattacaaa	ttgtctcctt	ttgaaagatt	gacctttctg	43440
acttaccagg	ataccataaa	gaaaaataaa	tcttatcact	tcagtcaagg	ataaagtatt	43500
tctgaattaa	aggaaaaata	caccagagta	aaatcaagac	tgaaagacaa	actgggaaat	43560
tatttgcaac	ctagatcata	gaaaaggggt	catttccctc	ttgcgtaaag	tgcaacttaca	43620
aattgataag	aagatgactg	ataactagaa	agaaaaatgg	gtaaagaaca	acaatagaca	43680
tttcacattt	aapctcatcc	atgataagg	aagtgcacaa	gaaaaactaca	ggggatacct	43740
tttttttttt	ttaatccatt	agattggcaa	acatcccaag	gtttgatcat	aggctcagtg	43800
ggtgagattt	aagtattatc	aggcattttt	atactttgct	gttaggaatg	caatgtagta	43860
caaacccttg	tagaagttgc	tttggaatg	tctctcagat	gtacaaatgc	attcacattt	43920
tagatttagc	atcccgctt	tctgagacat	tattcaacat	gtatacgtgt	gcacataaga	43980
tataataata	acacgttttt	ccttctagtg	tggtgctttt	aacctgtagc	ttgaaaaaac	44040
tctgctttca	ttgttttttt	ttgttttctg	tcactggctc	agccctgctt	tcaattgttt	44100
atatgaattg	atgggtgttc	tggtctggtt	ataacttact	ttagttaaag	agtcacttta	44160
aatttatatga	catctgatat	aagttgtgtt	aggtagaaaa	ttctgttaact	tggaatactg	44220
taagtacttt	gtggccacat	ttcattagta	ttaaatatta	tctctatata	tagtaggcta	44280
tttaatatcc	atattttatg	atgcaattaa	gaaataattt	ttttctgaag	ttggttagatt	44340
gttgatagtc	catggcccag	tgtttctcaa	agcattctgg	gggatcactg	ttgttcagaa	44400
ttagctgcag	tgattgttga	acatgcaggg	cctctgctcc	actccacgtt	gctaccagga	44460
cgctctgcag	gtgagagctg	ggaagctgta	gaagctgcag	tgctaacaaa	tgctacagga	44520
attctttagc	tcaccttcat	gaggtcttat	gttgaggaga	ggcagccagt	agtgctccct	44580
gtccttcccg	ttttatggtg	taagtttcat	tttaaggagg	gtataaatca	aagcccacct	44640
gggcattctc	tcatggttca	ctgcttcttg	taatcatgga	agatgtcatt	gcggcagaga	44700
cgaacacagt	tagtttgatt	actattgatt	tttttttaaa	tatttttctg	aagtggctgt	44760
tgtaatgtaa	taaattgtgt	gcttaaggac	aacctttggt	attctatttg	agttattgtg	44820
atgatcctag	tttaagtttt	tctaccagta	ttttcatatt	acaacatatt	tactttccat	44880
ttctattaat	atattttatat	ttaaagtatg	gaggccgggc	acagtggctc	acgcgtgtaa	44940
tcccagcatt	ttgggatgct	gaggcgggtg	gatcacaagg	tcaggagttc	tagaccagcg	45000
tgaccaacac	ggtgaaatcc	catctctact	aaaaatacaa	aaattagccg	ggcacagtgg	45060
taggcacctg	taattccagc	tactcaggag	gctgaggtag	gagaatcact	tgaatccggg	45120
aggcagcagt	tgcatgtgag	taagatcgtg	ccactggact	ctagcctggc	tgacagagca	45180
agaatccgcc	tataaaaaaa	gggatcaggg	aagaggggat	tacagataac	ccaaagaaga	45240
aggaaaaatc	tccacaagtt	cacctgtcca	gcggttaacc	caatttggtg	attttccctt	45300
aacaatttgg	atattttcct	ttaaatcctc	ttttttataa	tgtctatatg	ttggagagag	45360
tatgtgcctt	tapgtatttt	ttaaagtatg	gatttctgtg	tgtgtctata	tctcctgttc	45420
ttcatatttt	cttgtgtgtt	ataaacagct	gtacatgtca	gtatatatac	ttccgttaact	45480
ttttttttaa	ggctatatag	tgttcattga	tgtagtttaa	cagcagttat	ctccccggct	45540
tcactctgtt	ggatgtggg	tcctgtgtgt	tgcttccaga	gcaaattggg	cttgggtttg	45600
cagcaagtag	acctgtgacc	tgtagcaata	gttggaagac	tttctctatt	accaagcgt	45660
atcagtatac	tttagtgctt	actagaaatt	tatgggtaga	aaaacaataa	tatcttagag	45720
tattttttcc	tagattccct	aaggtgctat	agggtgattt	ttactcatgt	aacatgaact	45780
atgcttcaac	taagatagtt	tttgcaaatg	tgatatataa	agtactttat	taaacctata	45840
ggaagtattt	ataccactta	tttccctcct	tcagtgttag	aacctcctaa	atggcatttg	45900
acattgaact	gctttccact	ttgtcgcagt	ctcctctcat	tgctccctacc	tggtgctctg	45960
accttaggga	cttggtctgt	atagccccac	catggctacg	ctgggccttg	gtcgtctctg	46020
agacttagtt	tcttcatctt	acaaggagat	aataacagcc	cctgctgcg	tagaattgca	46080

gagatcaaat	gaaataatta	acatactcaa	aagcatgccg	taaacacatt	ctgagcacat	46140
gtacgttttta	gaaaaaacia	aaggaccat	gcacatttgc	gagtgtttt	gtctcagcag	46200
cactgcctct	tcttccaaag	ctgacgtctt	agtagaggcc	ctgccacgtc	ctgagcactg	46260
tactccacga	agcattctat	ttctgacatt	cgaaatgcag	tctgttccat	cttccttaca	46320
atctgtatgc	cagcacttga	aataccgggt	atctgcagt	ttgaccaggt	gattacttaa	46380
ttatggaaat	gttgagggtg	agatctagat	aattcagtga	aggcaggaaa	attggtgtcg	46440
gaatctgtct	ttttatgtgt	cagaaataga	aataagatag	ggtgagaagt	aatttgtggc	46500
taaaacacta	taatatgtaa	cacatagtgc	atatgtgtg	ccaagcactc	ctgtagggtg	46560
ttgaaatctt	ctattattat	tatccctact	ttatagactt	gcacccttag	gcacagagag	46620
gcggacagtt	gtccaagggt	accccagagg	tggagatcca	ggctacctga	ctccaccatg	46680
tgtgtctctc	cctagggcac	agttgtgtcg	ctaaaaatac	tttttaagca	gttctttgat	46740
tattcagatg	atagtactgt	aggaaaatta	agacaaaaat	aataaaaaat	taaaatcttt	46800
attttagtgt	tttgcacatg	tattattaaa	gccagtttac	tcctggaagt	gtgtaagaat	46860
acagggtatt	ttgatcacc	taaatgtgc	atgttactaa	gagctcgaca	ctgaagtcaa	46920
gaagagcagt	tgccagagag	acttagcaaa	aacgggaagt	gtgtgggggt	gaaggagcaa	46980
agacaagtct	tcctcggacg	gtggagtgtg	gaattcatca	tttctcagaa	caagctcttg	47040
aacgcatttt	caatttgagg	ccaaagggtc	cagcctccca	ctcggcatac	ctccctacct	47100
tagtcagctc	ttaaatctta	ggaatatttc	tttgttcttc	aaggaaactta	aatatgttaa	47160
cattcttacc	tgtccacagg	gagcccccta	caaagaaggg	agtttctagt	ctccgttctt	47220
tcttggaaata	aataatagcc	tcataccttg	tgcaatcgag	gctgaaaaag	actgtctcct	47280
tttttcaaat	aagcaagtct	tagaaactac	agttgtttac	aggggtcatg	gctattccac	47340
agtaataatt	ttggttcttt	taccaattat	ataatatgtt	aaaatatggc	aagtatcagg	47400
aaagcaagga	gtggcaatga	ttagaaacca	atggccaagt	tagagaggag	gggcaattgc	47460
tcccccaagt	ttgttgtggc	tgtgtagcag	tcagtgcaga	gaagctgtgt	gtcaggcgag	47520
aagcaaaagt	gaggattatc	aggcgccctg	gagtgcccag	ctgtgtgcca	ggtcaggagg	47580
tgccatcggt	agccagacca	gcttcctctc	ggccctgtg	gagctcgag	tctggtgggg	47640
aggcagcagt	caccatgggt	acagggtgaca	cactaggatg	gggctgggtg	tggtaggcat	47700
ttgcccgtcc	cttcagagag	gtgagtatgg	acttagagga	ggctccagct	tcctattcct	47760
gggctgtcta	tagcactaaa	agttgtcaca	tgaaaaataa	catttggtac	tattgattta	47820
acttaatgac	ttatgtaatt	gtagttgact	tagaaattat	aacatgctct	tctacttcag	47880
cttgaaccct	ccaaccacca	gtttataatc	cttttttttt	aacttttgtt	tatttttctt	47940
aaggaatctg	tactttttct	tcatttttaca	actttttttg	tcctgttacc	ttattttcat	48000
ttttacttta	tatgaccatg	agttctaaaa	tagtaaaaaa	aaagaattat	ttttgttctt	48060
tgttagaatt	tctctgcaaa	gaatgtccaa	aaattcatat	tcacattgat	cgtatcgaca	48120
aaaaagatgt	cccagaagaa	caagaacata	tgagaagatg	gctgcatgaa	cgtttcgaaa	48180
tcaaagataa	gtgagtaaca	acagttccag	cacttccgga	acttccggtc	aactgatttt	48240
cagtatagtc	aacaatttga	aaccaatgta	aatgggtata	ttgtctcaag	aatacatttt	48300
ataaattcaa	atcaaatttt	atgcatgtct	gatcgtgttt	taaactttac	ttgtacaaat	48360
cagtctaaaa	gaacttggtt	cagtgggccc	atctacttgc	attgatagta	tttcttggac	48420
aatactacgt	gataacatag	caaattaaat	taaaaacaac	aacaaacaca	caaaaaaact	48480
ttccagtgtc	agatgcccgg	acctacctgt	caggtcacat	aaagtgggtg	tactgtgtga	48540
ggcttggtcg	ttgggccagt	gtgcgcagaa	aagcaaggga	ggggtagagg	actatgcgga	48600
cggtgcagggt	gacatgatgc	tgttatattt	gttggaataa	gaagggggca	gttgacagcg	48660
ttatatccaa	agtgtcttct	gtggttaatt	atattcagaa	atttttagcca	attgttttat	48720
tctctaaata	tgtactttct	gtccaagaaa	ctatcattgt	tcttcttttc	cttgttttac	48780
agtacagtgt	ttttaattaa	ccctcctggg	ttaactttac	caggtgaaaa	tgattaaaag	48840
tgtaaatagg	taacaatgaa	actttaagct	tctatttttc	attgactctt	aactgtacat	48900
gatgtaatgt	attcagcgag	ccattcagga	ccactttggc	ccatggaaga	aattttaaaag	48960
taagatctac	atgtattgac	atgaaaatat	gttctcagaa	aaaagactaa	tgtatttaat	49020
gtcctactta	ttttataagt	atttagaata	cctctgggaca	ttttaaaaca	atgattattg	49080
ctaggggtgtg	tgatttataa	agcaatagaa	gcgctttccc	tttctgtttg	tgtttttagat	49140
tattatatcg	ggtatgttct	gctatcataa	ctttacaaat	cttatgtaat	atgggaaaat	49200
gagttaacta	tgctgttttc	cttcttttac	ctgcctttct	aattctgtgg	gaataaaggc	49260
gtttttgaga	cagcccagggt	gtagttagca	gtccatatcc	atggattcca	cattcatgga	49320
ttccaccaag	cacagaccaa	aaatactcag	aaaaaaagg	ggctggctgt	ggtggctcat	49380
gcatgtaatc	ccagcacttt	gggaggctaa	ggcaggcaaa	ttgcttgagc	ccagaagttc	49440
aagacagcct	gggcaacatg	gcaaaaccct	gtctctacag	aaaatacaaa	aattagccag	49500
gcgtgcacct	gtagtcccag	ctactcagga	ggccgagggtg	cgaggatcac	ctgagcctgg	49560
aagggttgaga	ctgcagttag	ctatcattgt	gccaaactcca	gcctggtaac	agagtgcctt	49620
ttttcaaaaa	aaaaaaaaaa	aaaggatttg	ggaggatatg	catatgttat	attcaaatat	49680
atgccatttt	attcatatat	cagggacttg	agcatccttt	gatcttggtc	tctgccgggt	49740

atcctgggac	cagccccctg	tcgatacaga	gggaccgctg	tctaagaacc	gctggtccta	49800
tctttgactt	ctggcggaat	aggagctcca	tgtaaaaagg	aggagaagct	gcagcgggtt	49860
attagccatt	tgtgagtcag	gtcactgtaa	aactttatca	aaagtttaaa	agacaaaaag	49920
cacccata	aaatgcctta	aaaccacctg	ttgaaatatt	acataataca	ttcatgtata	49980
ctaatacatag	agcatattaa	agataattta	gaagactaga	aacttctatt	aaaccaagtt	50040
tctggatgtt	tccgtattca	tccttatttt	ccagggaacct	gcataacttt	tccagcgtgt	50100
aatagctacc	tgattgatat	tttttgaatt	gaaataactga	agtgaactaa	atctaacttt	50160
tttccattct	ggpcatagga	tgcttataga	attttatgag	tcaccagatc	cagaaagaag	50220
aaaaagattt	ccgggaaaa	gtgttaattc	caaattaagt	atcaagaaga	ctttaccatc	50280
aatgttgatc	ttaagtgggt	tgactgcagg	catgcttatg	accgatgctg	gaagggaagct	50340
gtatgtgaac	acctggatat	atggaacctt	acttggctgc	ctgtgggtta	ctattaaagc	50400
atagacaagt	agctgtctcc	agacagtggg	atgtgttaca	ttgtctattt	ttggcggctg	50460
cacatgacat	caaattgttt	cctgaattta	ttaaggagtg	taaataaagc	cttgttgatt	50520
gaagattgga	taatagaatt	tgtgacgaaa	gctgatatgc	aatggctctg	ggcaaacata	50580
cctggttgta	caactttagc	atcggggctg	ctggaagggt	aaaagctaaa	tggagtcttc	50640
cctgctctgt	ccatttccta	tgaactaatg	acaacttgag	aaggctggga	ggattgtgta	50700
ttttgcaagt	cagatggctg	catttttgag	cattaatttg	cagcgtattt	cactttttct	50760
gttattttca	atttattaca	acttgacagc	tccaagctct	tattactaaa	gtatttagta	50820
tcttgcagct	agttaatatt	tcactctttg	cttatttcta	caagtcagtg	aaataaattg	50880
tattttagga	gtgtcaggat	gttcaaagga	aagggtaaaa	agtgttcacg	gggaaaaagc	50940
tctgttttagc	acatgatttt	attgtattgc	gttattagct	gattttactc	attttatatt	51000
tgcaaaataa	atttctaata	tttattgaaa	ttgtttaatt	tgacacacct	gtacacacag	51060
aaaatgggat	aaaatatgag	aacgaagttt	aaaattgtga	ctctgattca	ttatagcaga	51120
actttaaatt	tcccagcttt	ttgaagattt	aagctacgct	attagtactt	ccctttgtct	51180
gtgccataag	tgcttgaaaa	cgtaaggtt	ttctgttttg	ttttgttttt	ttaatatcaa	51240
aagagtcggg	gtgaaccttg	gttggacccc	aagttcacaa	gatttttaag	gtgatgagag	51300
cctgcagaca	ttctgcctag	atttactagc	gtgtgccttt	tgctgtcttc	tctttgattt	51360
cacagaatat	tcattcagaa	gtcgcgtttc	tgtagtgtgg	tggattccca	ctgggctctg	51420
gtccttccct	tggatcccgt	cagtggtgct	gtcagcggc	ttgcacgtag	acttgctagg	51480
aagaaatgca	gagccagcct	gtgctgcccc	ctttcagagt	tgaactcttt	aagcccttgt	51540
gagtgggctt	caccagctac	tgacagggca	ttttgcattt	gtctgtgtca	agaagttcac	51600
cttctcaagc	cagtgaaaata	cagacttaat	tcgtcatgac	tgaacgaatt	tgtttatttc	51660
ccattaggtt	tagtggagct	acacattaat	atgtatcgcc	ttagagcaag	agctgtgttc	51720
caggaaccag	atcacgattt	ttagccatgg	aacaatatat	cccatgggag	aagacctttc	51780
agtgtgaact	gttctatttt	tgtgttataa	tttaaacttc	gatttctcta	tagtccctta	51840
agttgacatt	ctgtcttact	gctactggat	ttttgtgca	gaaatatatc	agtggcccac	51900
attaaacata	ccagttggat	catgataagc	aaaaagaaa	aaataatgat	taagggaaaa	51960
ttaagtgact	gtgttacact	gcttctccca	tgccagagaa	taaactcttt	caagcatcat	52020
ctttgaagag	tcgtgtgggt	tgaattgggt	tgtgtacatt	agaatgtatg	cacacatcca	52080
tggacactca	ggatatagtt	ggcctaataa	tcggggcatg	ggtaaaactt	atgaaaattt	52140
cctcatgctg	aatgtgaatt	ttctcttacc	tgtaaagtaa	aatttagatc	aattccattgt	52200
ctttgttaag	taacgggatt	taatatattt	tgatatataa	gggtatgttc	taaatttgaa	52260
ctttgagagg	caatactgtt	ggaattatgt	ggattctaac	tcattttaac	aaggtagcct	52320
gacctgcata	agatcacttg	aatgttaggt	ttcatagaac	tatactaata	ttctcacaaa	52380
aggtctataa	aatacagctg	ttgaaaaaaa	ttttgtatca	aatgttttgg	aaaattagaa	52440
gcttctcctt	aacctgtatt	gatactgact	tgaattattt	tctaaaatta	agagccgtat	52500
acctacctgt	aagtcttttc	acatatcatt	taaacttttg	tttgtattat	tactgattta	52560
cagcttaggt	attaattttt	ctttataaga	atgccgtcga	tgtgcattgt	tttatgtttt	52620
tcagaaaagg	gtgtgttttg	atgaaagtaa	aaaaaaaaat	aaaatctttc	actgtctcta	52680
atggctgtgc	tgtttaacat	tttttgaccc	taaaattcac	caacagctct	ccagtacata	52740
aataggctt	aatgactggc	cctgcattct	tcacaatat	tttccctaag	ctttgagcaa	52800
agttttaaaa	aaatacacta	aaataatcaa	aactgttaag	cagtataatta	gtttgggttat	52860
ataaattcat	ctgcaattta	taagatgcat	ggccgatgtt	aatttgcttg	gcaattctgt	52920
aatcattaag	tgatctcagt	gaaacatgtc	aatgcctta	aattaactaa	gttggtgaat	52980
aaaagtgcg	atctggctaa	ctcttacacc	atacactactg	atagtttttc	atatgtttca	53040
tttccatgtg	atttttaaaa	tttagagtgg	caacaatttt	gcttaatatg	ggttacataa	53100
gctttatttt	ttcctttgtt	cataattata	ttctttgaat	aggtctgtgt	caatcaagtg	53160
atctaactag	actgacata	gatagaagga	ataaaggcca	agttcaagac	cagcctgggc	53220
aacatatcga	gaacctgtct	acaaaaaaat	taaaaaaaat	tagccaggca	tggtggcgta	53280
cactgagtag	ttgtctccag	ctactcggga	gggtgaggtg	ggaggatcgc	ttcagcccag	53340
gaggttgaga	ttgcagtgag	ccatggacat	accactgcac	tacagcctag	gtaacagcac	53400

gagaccccaa	ctcttagaaa	atgaaaagga	aatatagaaa	tataaaattt	gcttattata	53460
gacacacagt	aactcccaga	tatgtaccac	aaaaaatgtg	aaaagagaga	gaaatgtcta	53520
ccaagcagt	atlttgtgtg	tataattgca	agcgcatagt	aaaataattt	taaccttaat	53580
ttgtttttag	tagtgttttag	attgaaagatt	gagtgaata	ttttcttggc	agatattccg	53640
tatctgggtg	aaagctacaa	tgcaatgtcg	ttgtagtttt	gcatggcttg	ctttataaac	53700
aagatttttt	ctccctcctt	ttgggccagt	tttcattacg	agtaactcac	actttttgat	53760
taaagaactt	gaaattacgt	tatcacttag	tataattgac	attatataga	gactatgtaa	53820
catgcaatca	ttggaatcaa	aattagtact	ttgggtcaaaa	tatttacaac	attcacatac	53880
ttgtcaaata	ttcatgtaat	taactgaatt	taaaaccttc	aactattatg	aagtgtctcg	53940
ctgtacaatc	gctaattttac	tcagtttaga	gtagctacaa	ctcttcgata	ctatcatcaa	54000
tatttgacat	cttttccaat	ttgtgtatga	aaagtaaata	tattctctgta	gcaactgggg	54060
agtcataat	gaggtcaaag	acatatacct	tgttattata	atatgtatac	tataataata	54120
gctgggttat	ctgagcaggg	gaaaagggtta	tttttaggaa	aaccacttca	aatagaaagc	54180
tgaagtactt	ctaataatac	gagggaaagta	taatatgtgg	aacaaactct	caacaaaatg	54240
tttattgatg	ttgatgaaac	agatcagttt	ttccatccgg	attattattg	gttcatgatt	54300
ttatatgtga	atatgtaaga	tatgttctgc	aattttataa	atgttcatgt	ctttttttaa	54360
aaaaggtgct	attgaaattc	tgtgtctcca	gcaggcaaga	atacttgact	aactcttttt	54420
gtctctttat	ggtattttca	gaataaagtc	tgacttgtgt	ttttgagatt	attggtgcct	54480
cattaattca	gcaataaagg	aaaatatgca	tctcaaaaat	tggtgataaa	aagttatttc	54540
ttgtatatgt	gtataaagtt	acatgttgtg	tatatatgtt	gtattgcaa	atacggctat	54600
taaatatact	gtcatatttt	aaaggttcag	ttttagtgta	tagtaaacaa	gcagtgcact	54660
aagcctcttg	cgggcatcat	ctcatctcac	tgtcatcaca	aaccccatgc	cacagcgtag	54720
cttgaccact	aaaagtaatg	catctgcaag	catactgcca	ggttttggat	agtttgtacc	54780
aacagttacc	ttatcaaggt	aaatcccaga	ctctaaaaga	gttggtgctg	tgtcactaca	54840
tgcataaact	taataaaatt	tcctgccggg	cgcggtggct	cacgcctgta	atcccagcag	54900
tttgggaggg	caggcgaagt	ggatcacttg	aggtcaggag	tttgagacca	gcctggccaa	54960
cgtggtgaaa	ccctgtctct	actaaaaata	caaaaattag	ccaggcgtgt	ggtggcaggc	55020
acctgtaatc	ccagctactt	gggaggatga	ggcaggagaa	tcatttgaat	cctgcaggcg	55080
gaggttgcag	tgagccaaga	tggtcgtcatt	gcactccagc	ctgggcgaca	agagcgagac	55140
tccgtattaa	aaaaaaaaaa	aaaaaaaaaa	aaaaattcct	ctcctgtttg	agctttccct	55200
tacctgtaaa	gaggggagaa	tatgtattta	cttcaagag	ttcagggaaa	tgactctcac	55260
tagtttgaga	ttctaggtat	aaaaatacat	tcttatataa	ttttaacacc	aatgtgagag	55320
attattattc	ttgctaaacc	aattcagttt	tatttgctgt	ctaaaatgtg	tgaataagta	55380
attgtccatt	attttctgaa	gtgttttggg	actcaacaca	tgattgtgag	gaggatttgt	55440
tgctaaacat	ctttctggtt	attcaagctc	gtgtatactg	tgtctgtgtg	agacatgcag	55500
agttactttc	tgtctgggtc	acaggtcagt	tcttgatagt	tttcggacaa	ttaaccagtt	55560
ttcatttgcc	catgaccacc	tttattcttt	ttcctcaact	gcacccatct	tttataaggt	55620
ctttcagttt	attgcagaga	agatggtgga	gaaaagccgg	aattcccacc	caccgctgcc	55680
atcccatgt	tttatcattg	gctagagtgg	aaaatagcag	taactactgt	gagagatcat	55740
ttgtttatat	aatggaaaca	aagatgagga	aagaacctgg	cttagatcag	agaactgatg	55800
tatttagatt	cttttttttt	ttttttttta	gacggagtgt	tgtctgtgtg	cccagactgg	55860
agtacagtgg	ctcaatctcg	gctcactgca	acctccattt	ccctggttca	agcaattatc	55920
ctgcctcagc	ctcccaagta	tttgggatta	caggcgtgtt	ccaccacacc	tggttaattt	55980
tttgtatttt	tagtagagac	gggttttcgc	catgttggcc	aggctggtct	cgaatctctg	56040
acctcagatg	atccacccgc	cttggcctcc	caaagtgtcg	ggattacagg	cgcgagccac	56100
cgcgctggc	ccaatgtatt	tggattctta	aagaacactt	tcaaattaaa	tatcagttga	56160
agagaactag	aactaaagaa	tttctgtgtc	aaactgttta	gcaaattgaa	gtagaagctg	56220
ggagatgtgt	cctggaatga	atgaatacat	cagtaaaata	ccatacgtat	gttatgatgt	56280
tattgtttcc	ttgccttggt	tgatttggtt	ttactgtgaa	ataattttca	atatagaatt	56340
gtgatcgttg	gaatttggtc	atctactaga	aaatgagaaa	gaagttaata	gctatcttcc	56400
ttaaagattt	ctgaggttgg	gattaaggta	gtgttcccaa	ggtgttctaa	aacggcagcg	56460
agagctgtgc	actcacttca	caaatttgaa	ttcctgtctc	gtgttagggc	ctgtgctagg	56520
<210>	180					
<211>	2000					
<212>	DNA					
<213>	Homo sapiens					
<400>	180					
gtggatctgt	gactgttcgc	aggaagagag	gagcgggagc	aggacagaca	ataactgata	60
gtcaggagct	gggtttggag	ataaagaggg	aacaagagaa	agttaagttc	tgtgttttca	120
tggtcaaacat	tgacaaaaag	tttacaactt	cgtgactaac	agtaatctgg	ggtgattcac	180
aacaaattta	capataaaca	catatttact	gactttatac	acagcaatcc	taacgtgaac	240

acagaacctg	ctttatcttt	tcgcacactg	ttctagtgtg	gagatgtctg	gtctcagtta	300
aagaaagcat	aaggagcatt	agttgtgcac	actgtccaca	cccgtagctt	ttttccacca	360
gtactaaacc	tagtgcttct	tacagtacag	ggcaatgaca	gccacagaaa	gagagaagct	420
ccttttactg	tgtaatgctt	cctgctggcc	ttcaaatact	tgttacttga	gagatctcca	480
ttcacctggc	tttgccccca	aaggtcacat	tctaccaatg	atgttgttat	ttgatgttaa	540
tcatgtataa	agaaagtagc	taccatcctg	gccctgatta	gaacttccca	ctgaaatacc	600
gtcctgccta	aaggtagcac	aggtttccat	tatgggtggg	gtggggaggg	ggcgggaata	660
tatatatata	tatatatata	tatatatata	gtaaagcatt	cggcattctt	ttaaagtaca	720
actatccttg	aaaagggtta	catattaaac	cattttttacc	acagccaaag	gggaggagaa	780
agatccaaaa	gtcctgtgga	tctgctttaa	catcaataaa	acagttatcc	acccttcgta	840
gcttttagtg	aaggctacaa	aagtatgctt	tttatggatt	acacatgtgc	acgcaactac	900
tttaattact	acagaaaaaa	acgaggctcc	ttattaaaaa	aaaatcagaa	acaagtccaa	960
cagactctga	ggaaatgaag	caagagtga	ttctgaaaag	gtctaataaa	cagtatggaa	1020
atatacctgt	gggattgttc	ttcagctatg	cataaacatg	taattatcat	cattactgtg	1080
atggggaaaa	acacggacc	taattctgaa	acacccctgt	agcgagagac	gggcaggagg	1140
ggctgctgcy	cactcagagc	ggaggctgag	gaggcggtc	ccccttgcaa	aggactggca	1200
gtgagcagat	ggggacactc	gagctgcccc	gcgacctggg	ccgagctgcc	tacaacctgg	1260
gccaggtg	ctgcaagaat	tagacctccg	ataacgttaa	caccactttt	ctcactgctc	1320
taattgtgtg	catcccggcg	cccaggggct	tgtgagcagc	aggtgctgct	tccaggcagc	1380
tccagcgacc	cttaaacctg	accgcgcgca	cgtccggccc	gaggagagcag	aacaagaggg	1440
acccggaacc	tccctccggc	agcaccacc	ttcaaccagt	tccgtcagtc	gccaccacct	1500
cccttcccg	gtccgcagcc	ggcccagctg	gggagcatgc	gcagtggccg	gagccgggtt	1560
gcccgcgcca	cagcaggtag	ctgtactgca	actgtcggcc	caaaccaacc	aatcaagaga	1620
cgtgttattg	ccgcccaggt	ggaactatgg	caacgggcga	ccaatcagaa	ggcgcggtgt	1680
tgcgcgggag	ccpccctgcc	cggcaggggg	atgtggcgat	gggtgagggg	catgggggtg	1740
gagcatccct	gagccatcga	tccgggaggg	ccgcgggttc	ccttgctttg	ccgcggggag	1800
cggcgcacgc	agccccgcac	tcgcctaccc	ggccccgggc	ggcggcgcgg	cccatgcggc	1860
tgggggcgga	ggctggggagc	gggtggcggg	cgcgccggcc	cgggcccggg	cggtgattgg	1920
ccgcctgctg	gpcgcgactg	aggccccggg	ggcgggcggg	gagcgagggc	ggagctcgct	1980
gccgcccagc	tgagaagatg					2000
<210>	181					
<211>	1901					
<212>	DNA					
<213>	Homo sapiens					
<400>	181					
taaaggttca	gtttgtagt	atagtaaa	agcagtgcac	taagcctctt	gcgggcatca	60
tctcatctca	ctgtcatcac	aaaccccatg	ccacagcgta	gcttgaccac	taaaagtaat	120
gcatctgcaa	gcatactgcc	aggttttggg	tagtttgtac	caacagttac	cttatcaagg	180
taaatcccag	actctaaaag	agttggtgct	gtgtcactac	atgcataact	ttaaataaat	240
ttcctgccc	gcgcggtggc	tcacgcctgt	aatcccagca	gtttgggagg	ccgaggcaag	300
tggatcactt	gaggtcagga	gtttgagacc	agcctggcca	acgtggtgaa	accctgtctc	360
tactaaaaat	acaaaaatta	gccaggcggt	tggtggcagg	cacctgtaat	cccagctact	420
tgggaggatg	aggcaggaga	atcatttgaa	tcctgcaggc	ggaggttgca	gtgagccaag	480
atggcgctcat	tgactccag	cctgggcgac	aagagcgaga	ctccgtatta	aaaaaaaaaa	540
aaaaaaaaaa	aaaaaatccc	tctcctgttt	gagctttccc	ttacctgtaa	agaggggaga	600
atatgtattt	acttcaaaga	gttcagggaa	atgactctca	ctagtttgag	attctaggta	660
taaaaatata	ttcttatata	attttaacac	caatgtgaga	gattattatt	cttgctaaac	720
caattcagtt	ttatttgctg	tctaaaatgt	gtgaataagt	aattgtccat	tattttctga	780
agtgttttgg	aactcaacac	atgattgtga	ggaggatttg	ttgctaaaca	tctttctggt	840
tattcaagct	cgtgtatact	gtgctctgtt	gagacatgca	gagttacttt	ctgtctgggt	900
cacaggctcag	ttcttgatag	tttccggaca	attaaccagt	tttcatttgc	ccatgaccac	960
ctttattctt	tttctcaac	tcacccatc	ttttataagg	tctttcagtt	tattgcagag	1020
aagatgggtg	agaaaagccg	gaattccac	ccaccgctgc	catcccatg	ttttatcatt	1080
ggctagagtg	gaaaatagca	gtaactactg	tgagagatca	tttgtttata	taatggaaac	1140
aaagatgagg	aaagaacctg	gcttagatca	gagaactgat	gtatttagat	tctttttttt	1200
ttttttttta	agacggagtg	ttgctctgtt	gcccagactg	gagtacagtg	gctcaatctc	1260
ggctcactgc	aacctccatt	tccctgggtc	aagcaattat	cctgcctcag	cctcccaagt	1320
atttgggatt	acaggcggtg	tccaccacac	ctggctaatt	ttttgtattt	ttagtagaga	1380
cgggggttgc	ccatgttggc	caggctggtc	tcgaaatcct	gacctcagat	gatccaccgc	1440
ccttggcctc	ccaaagtgct	gggattacag	gcgcgagcca	ccgcgcctgg	cccaatgtat	1500
ttggattctt	aaagaacact	ttcaaattaa	atatcagttg	aagagaacta	gaactaaaga	1560

atttctgtgt caaactgttt agcaaagtga agtagaagct gggagatgtg tcttggaaatg 1620  
 aatgaatata tcagtaaaat accatacgtg tgttatgatg ttattgtttc cttgccttgg 1680  
 ttgatttggg tttactgtga aataattttc aatatagaat tgtgatcgtt ggaatttggg 1740  
 catctactag aaaatgagaa agaagttaat agtatcttc cttaaagatt tctgaggttg 1800  
 ggattaagggt agtgttccca aggtgttcta aaacggcagc gagagctgtg cactcacttc 1860  
 acaaatttga attcctgtct tgtgttaggc gctgtgctag g 1901  
 <210> 182  
 <211> 4550  
 <212> DNA  
 <213> Mus musculus  
 <220>  
 <221> exon  
 <222> 2259..2488  
 <223> exon1  
 <400> 182  
 tacagctatc tgtgtgtatg ggtgcâtgcc atggaccacc tgtagaâgtc acaggacaat 60  
 ctccaattca ctttctcctt ccaccacatg ggctctgtaa tcactaaggt ttgtcaatcc 120  
 tgagtacaga tgttcagAAC catcttactg tctcctctct tctgataaaa catgaggttg 180  
 ccccagagac gtttttagaca gggttataat ctgataaggg aaaagccaca tgccttttcc 240  
 ttacaaatgt aatttctaca gacattccta gaaaattgaa actttatggt tgggaaagga 300  
 gagggggccc tcaggtacct tgtttttctg ttgacaaaag ttgactctta acattgtcaa 360  
 gtaaattgtc ccacaaatgg atcatctgac tatttgcaga atgtcatagg ccaacagaga 420  
 gagaaccctt gaatttccag agaccttcag gttggctcag tcccttcttt ttgatgtgt 480  
 acctcaattc ctgtcttctt gaactcttgt ttgccaatct gaactctacag tctatctgtc 540  
 aaacaattcc tttgtctgga ctggtctgct gaactgacag tgaattgtct tgacagtcc 600  
 tttgcctgcc cttttacctc tgcatcttca ttaaactgga cagtttgtca tatctgtgac 660  
 ccaccaacag ctgcttttcc cctaaagctg ggtttgtggt tcatgttatc gtgacagaca 720  
 ctcttatagc cctgtcagtt ctccagcact ggcttcccaa ggctttttaa actcctttct 780  
 tctttctaac tctttgtagt cactgtaacc tatatatgca tatgtaaaca gagatatact 840  
 tacagagtga tgtatgtgtg atctgagagt taatattagt aattaagact gcaataaaag 900  
 aacctgtgtt tcccttagca agggctacag agtaaaagtg gcctctctgg tgccagcgaa 960  
 gccactgtac atctggaat ttattgtcat tcaatacatt ctgatatcgt gtaaactcct 1020  
 aagcacgtcc atctgacata gtgtgcta at gacaggagtc acctgtatgc cttatgaagc 1080  
 gcatctcaga ggtgatggga aagaacatg gggcaaaaga tgaagggaaa tccaaggcaa 1140  
 ggaagcagag acacaggcgt cagtgtgtgt gaaagggaga aaactagggg cagaataagt 1200  
 gaccttaggg tcaacttagag aaaccaacac acacacacac accacatat ttaaacgta 1260  
 ctttatacag atctgagcgt gcgcactgac ctgtttcctt ctataccttc ttgtatagaa 1320  
 ttatctggtc tccactagtt agggcagtg aaggacctgg gcccttgat aagtttttgc 1380  
 tgttacttaa ctattctagt tttctggagg gaagagaact tatggatcct acatgtatag 1440  
 ggaaatactt tcttacacat tgaaaagaag aaatgtagga tattaggaaa acgcacagta 1500  
 gaaacaagtt aaagagcaag aggttattaa agggcaaaag ttaaggcttt gaaagattta 1560  
 atacaaggag gtgacagtc cgtgaaaggt gaaccaaggg tacaggagac ggaccagcc 1620  
 tcattctgca acagccaaga ggagggaagg tgtgttctt atgcacgtgg gggcacgggt 1680  
 ggccctccgg cagcggaaga cgtgcagtt gtccataacc tgcggcatcg agctcctcct 1740  
 gtgtccacg acttagtcgg ctacgcgtg tcttgacgga agcatcctcg tgtctccacg 1800  
 cagctctcgc acgccagcac aggccaaaac ccaccacctc acttcttccc gggctcatcc 1860  
 ccagccagca ttgcagtcg agcatgcgtc gtgacgaggg caagggaccg agccaatcag 1920  
 aacacgtatt acgcccataa gtcggccaat caggaggcgc cttattaccc gggagccttg 1980  
 cttcaccccg cctccccgct gacaagcacg ggtcgcgcgg agcaaagcga gcaccccgag 2040  
 gcgagtgccg ccggcaagcc gaggcgtgcc ctttccaagg cggcgagcag aggcgcgtcac 2100  
 tgtccccgcc gggccccggg cccccgcggc ccatgctggg gggcgagcca gggcgagggt 2160  
 cggcggcgcg gccggccccg cgcagtgatt ggcggggcgg cggcggtggc tgaggctctg 2220  
 gtgcgcgcgc gggcaacgca ggcggagtcg cggctggcga gccgagagga tgcgtctgtc 2280  
 cctggtgctc cacacgtact ctatgcgcta cctgtctccc agcgtcctgt tgcgtgggtc 2340  
 ggcgcccacc tacctgctgg cctggacgct gtggcggtg ctctccgcgc tgatgcccgc 2400  
 ccgcctgtac cagcgcgtgg acgaccggt ttaactgcgtc taccagaaca tgggtgctctt 2460  
 cttctctcag aactacaccg gggctccaggt gaggcgcggc cgcgcagggc tgcgtgcgag 2520  
 cctccccgc ggccggggcg gcgcttgcaa cccggcgcaa cactcgagc ccggcgagca 2580  
 cgtgcgcag ctacggcct cccgcgcggg ggggaagtgt ctggttctca cttcggggtt 2640  
 ccttctggaa cgtcctgctg aggtctgagt tgttcccggt tccgccccac ccccgccccg 2700  
 ggccggctgt tactgcccac ctacgtgcct gccaaagtag ggcactgagt ccgaggtggt 2760



gatgctggga	ctggcttcat	ttgcacttcc	gaggtctttt	agattagcaa	gacctctagg	2820
cgctgaccaa	agtgcagcgt	gtgaaggacg	actcctgcct	tgggttcctc	ccgggtgaaa	2880
gcgagggcct	agggaggaaa	tgaatacatt	ggttacaata	ggagcctcac	tgctgataca	2940
gttctcttca	gcttggactg	ggcttcaatg	tgggctgata	tcttgtcaga	ttgctttctt	3000
cctgctactg	tttctttctt	tctttccanc	cctccctccc	ccccccccc	cgccccgtgg	3060
agattgaact	ctgaaaacaa	taaagagtag	aaagctctcc	taatgtgaat	tcgttatatg	3120
acatcccata	aaaacctaca	gttgactctc	ctttttgggt	ttcagtttca	aagaagagct	3180
ctgtttgggt	tctcccagat	gtatctatga	ctttccccc	catttctcag	ttcttttcat	3240
tctgtgttag	gggggtactt	tggcgactgg	atcccttact	gagttttgcy	ccagttggag	3300
attatgtctg	aggtaggga	ttaagacctc	tctgaatcac	tatcttttta	aatgttttcc	3360
tagggaatag	gaaaatcact	gttgacatc	aaggtttctg	aaaaattgac	ttttagaata	3420
ggatttcatt	cagaattttt	aggaaccccc	acactgatgg	tttcaaacct	ccctcttact	3480
ttactaagtt	tgtcaagtga	atgtatggtc	taatcgtgga	taagtattta	atttcactag	3540
cagaagggac	aagacagcgg	ggagcacaac	ttaaagttgc	tgaccttgca	catgacaagt	3600
acccttcaga	cgctcaggga	cctctactca	agtgccacct	atattcttgc	tgagagagcg	3660
ttaggatgag	tcagaatgaa	gcaaagttag	tgagtttatt	gattgggaga	gaggacacgc	3720
acttgagggg	agtcaagtgc	aaaccttatt	acccccacc	caggctacag	cagctgtttt	3780
ctaagtatt	ttagggcttt	taagttaacg	ccttaaaact	aagattaagg	agaagagaag	3840
gaaaaaatg	agttcttcta	ttctttccaa	taatgagctc	taaaaaaaaa	agaagcaaac	3900
caggatctca	cactgtatgc	ttgggtggga	ggaactctat	gtagacctca	caggcctcaa	3960
gttcacagag	atctgcctgc	ctctgtctcc	agagtgttag	gactaaaggc	atgtaccgcc	4020
atgtctggat	taaactcttt	tagtttatatg	aaatttaaaa	cggattcatg	gcgggtactga	4080
acagtttaca	tatgagggag	aaatgtgggt	aggcagtaat	atggatcaaa	ataaaatcaa	4140
agtaattagc	tgatcactgg	tcacaagagt	ttgagatgtg	agcttgtctt	ctgccttagg	4200
tcaccagcta	tagggataat	cttttgtttg	ttttttgttg	tttttgtttg	tttggttttt	4260
tggttttttg	agacaggggt	tctctgtgta	gtcctggctg	tcctggaact	cactctgtag	4320
gccaggctgg	cctcgaactc	agaactccac	ctgcctctgc	ctcccaagtg	ctgggatgaa	4380
aggcgtgctg	caccacttgc	ctataatctt	acttgtaatg	gttttagaat	atgtgcacag	4440
tggagagcag	tgttcaagca	gctgtatcca	accaattnca	cttaaagagg	gagaggggtga	4500
gggtgagggc	ctccttttgc	tattcaaaa	cagattgtgt	ggacattgca		4550
<210>	183					
<211>	37950					
<212>	DNA					
<213>	Mus musculus					
<220>						
<221>	exon					
<222>	5259..5328					
<223>	exon2					
<221>	exon					
<222>	12675..12791					
<223>	exon3					
<221>	exon					
<222>	14621..14710					
<223>	exon4					
<221>	exon					
<222>	19822..19912					
<223>	exon5					
<221>	exon					
<222>	21789..21950					
<223>	exon6					
<221>	exon					
<222>	23387..23510					
<223>	exon7					
<221>	exon					
<222>	25520..26016					
<223>	exon8					
<400>	183					
tgagtgaga	ggcctgggta	taattccttt	ttctttgtca	cactgtagca	gttctgcttc	60
tcagcctcag	ttgagactgg	aatacatttg	tcagtctgtt	ctgaagactt	taatgggtga	120
tctttactgt	caccttgact	ggatttagaa	tcgccttgga	gatgtgctta	tggtatgtct	180
gggaggatgt	tttcagaaa	tgtttactga	aggtgtcctc	atctgatagg	gtggggccct	240

ggactcaata	aaaaccggca	tttatctggt	tcctgtccag	ggacacagtg	tggccagcta	300
cttcacactc	ttgctgcctt	cctacatcga	aagactgtat	cttttcttaa	aaggcgaggc	360
aaaataaatc	cccaccccca	ggattttttt	tggctgtcct	ggaactcagg	tccacctgca	420
tctgcctcct	gagtgtctgg	tctacaggag	taccaccag	gcctggctaa	aagaaacctt	480
tcttaaatgt	ctpcgtgcag	acactttgac	acaacaataa	gaaaaataat	tcatacaaa	540
accaagttag	ataatatgtt	atTTTTttag	ttcatagagt	agtaggtaat	ccatgggtgag	600
ggaaaaaaa	aaaaanaacc	cttttgaaat	aaggacagta	ttgagaaaca	tgtatgctga	660
ttgtatgagt	tttgtgat	gtaagtttct	actcaattca	catggtaatt	gtgcattctg	720
atcattaatc	aaataattgt	gtttactact	ttaaccttct	tacaaagtat	agttttacta	780
attagtaatt	tagtaaat	attatttatt	taatgaaaca	tttcaaattg	gactcagaaa	840
aagaccacag	tttttgtaca	tattatagta	gaagtcctt	agtaggtgga	aatcctgtgt	900
ttctttacaa	ggatatgtct	agaacacgtt	aaacaaacag	gaggaggtgt	ggctgcaccc	960
gttaggctaa	ccagtcaaca	tgccttttaa	agccatacgt	gttgtgtgtg	agcatttttt	1020
taaatatata	gaaaatcccc	aaaatagcta	gtataataag	cacacatgcc	agtaagcctc	1080
ttattactgt	aaaataactgt	gtaatacttt	gtgcgttctt	ttatgtgact	gcagtaggtc	1140
tgtttacatc	agcatttacc	catacaaggt	ggctactgtc	attcaggcct	tggaaagttt	1200
tcagcttctt	tgaacttgt	gggttactt	ctccagttgt	atgtgctgtc	catggttggc	1260
tagacatggc	acatgactgg	cgcagtgacc	taaagacaga	ttaaagaatt	catttcaaat	1320
gatttcaagc	acagagttta	aattgtacat	gcactcttag	tcaatgtcct	tgtagctgca	1380
ttttgggtga	attggaggga	catggactag	ttagctgttt	cttttacttg	atgacagttt	1440
tgaatgaaaa	gcatgttaga	tacaaaataa	tttaactgtt	gacccccccc	cacacacaca	1500
cacacactct	ttttccctgt	atgccttcac	tcctcgaatt	ggtttttatc	anatagctgt	1560
tccaggtgta	ccaggcatat	atgaagtagt	ttatgtagtt	tcatttatgc	cagtgcacac	1620
ctgtgaggag	caattagtac	attatacttt	atagttaatg	agatagatat	agagaaagct	1680
gatacatcac	atctatttct	tgtgtaagat	ggagggtgg	gattcaaatg	taagtcttag	1740
tccgtgtctt	atactttgca	gcctgtggtc	ttagtgtatg	tcagtgtatg	aagggaacc	1800
tctttttcag	actgtgtctg	caagaccctg	gaattaaatg	accaaaggca	tactaactgt	1860
agagaaactg	cccattttat	tgatgcta	atTTTTacat	ggtaggagga	aacttggaag	1920
aatgagaag	ccctactcag	ggggattttt	caggtgagat	tatgtagtga	ctagtgtaaa	1980
agaagctact	ataaggata	ccaagtatgg	gaaataagtg	ctgcacacct	cagggtggtc	2040
acacagactc	agactgacag	ctcaggtcct	ctcgtggaag	aaggggacgt	ctttgaaagt	2100
gagaaattca	tgtcttcttt	atagaaagtt	ctactccagc	tctaggccga	agactggggg	2160
cagagagctt	ttttgtgccc	tgcgatttct	taactgtttt	tattcaaaat	cattcttatg	2220
ccaaaagggc	atatttgggg	ttgagttcct	tcagcgatat	catttatatt	tgagaccaat	2280
gagacgtttt	tctcactatg	tattgtattt	caactttcat	gctaaacctt	ctggacttta	2340
ctgagataaa	tcaagaacat	acctttaaac	tttgtagtct	ttctttgcca	ctgtgaccaa	2400
aacacatgac	tgggagaaat	ggtttatttt	gcctcccagt	cttggtgatt	ttagtccctc	2460
acagcagaga	agcctggtag	agcagctcac	tgagtggcaa	caggactgtg	ctcatgtgat	2520
catggaccag	gaagtgtaga	acaaagccag	aactaggggc	cctagtaacc	tacttctgcc	2580
agctaggccc	cacttcctga	aggttccacc	tcctccctcc	ccctcccaa	aaagaaaata	2640
aaaatagtag	caccaactgg	acagcaaaca	cccaaacaat	aagccagtga	ggagggaagg	2700
agggaggggg	gaggaaggag	ggagagaggg	gggaaggaga	aggagagaga	gaaggagaga	2760
gagagggagg	gagggagaga	gagagagaga	gagagagaga	acacaccanc	tctcaagtat	2820
nnnnctnna	nnncttgaa	caaattaaaa	actatgtaac	cagaaaatta	ttttaagagt	2880
atttgatttg	tctgtattta	attaattaac	tgaaaaaaaa	aggaaaatta	attcctttct	2940
ctaaagactt	taaatgaacc	atTTTTttag	tgtatgtgtg	tgtgtgtcta	ggtcaaaaga	3000
caggtctcag	gagttgatat	ttgaccatgt	gaaccctagg	gatcgaactt	ttagctcatc	3060
agctttggca	gcaggaatct	ttatacactg	catcatctca	ctggccttag	ataagttttg	3120
aaaaaaagtg	gaagaatcta	aagttacttg	gataattata	taaaatataa	gtctgaggtt	3180
gggtcaaga	cggaaagaca	tcagtaagga	gccaagagaa	cagccccaga	gagaatctga	3240
gaatcaaggg	tgtgggcaaa	catacagtga	tctacacttt	ttatctgtaa	atagtttgta	3300
aattcttaac	cctttaaaaa	aattccctaa	cccctactct	gcagatcaga	ctgccctgga	3360
actcattgta	gaagcctaag	gtggcctcgc	attcacacaa	tccttctgac	acagctcccc	3420
aagtgtagg	attacaagga	taagctactg	tgtctgcctt	cttagctttt	taaattttta	3480
aagcactaca	acpttgtaac	tagttttata	cttgtaata	aaataacagt	aactgggact	3540
ggggagatgg	ctpaggctgc	ttttccagag	ggacccaggt	tcgattccca	ggagccacac	3600
aatggcttac	aatcatctgt	agctatagtt	ccagggtatc	tgatgccctt	ttctgattcc	3660
tgcaggcccc	aggcaagcat	gggtgtaca	aacacacatg	caggcaaaac	agatgtcttt	3720
ctgcattgct	cttcaccttt	tatattgagg	caaggtctct	cacttgaacc	cagatctcac	3780
tgattggcta	gtgtaaactaa	ctggcttgct	caagggaatcc	ctgtctacac	ttcactaaag	3840
ctcttgctgt	gtagcccagg	ctagcctcaa	attcctaate	ctcctgcttt	agccacttaa	3900

gtgtccggga	tttcaggcat	gcaccactac	atctggttga	aaggatatctt	tggatgggtt	3960
gggtattatt	agcttggagg	ctagcgtatgt	ctccaaagaa	ccttacataa	tttacatatg	4020
catacatata	taacatacata	aacatacata	catacacaca	tgcatacatt	taactggatc	4080
tgtagtctct	gtaaagtgtg	cccagtggtc	cttaaatagt	tctactttat	ttctaagagt	4140
gatcatgata	atgagctggc	atcccaatat	taactctgca	cagatcaact	aaccattggt	4200
tacttatttg	tttgatttat	gtcgtttgta	aacttgatta	gaagtaatta	gacacgtgtg	4260
aacacactgt	gcagttgtct	tagagcagta	gcagcagcct	tagtgtgagt	ccttgacatc	4320
taactttata	ttatgtagct	acattgcttt	tacacagggc	ttattaatgc	cctttcaggt	4380
tttgttatca	ttatgggtatt	tgtgagtatt	tcaccaccag	gaaggcaaac	cctttcggca	4440
tagtagactc	caaagcgttc	actaacttat	gcataattct	tagtgacaaa	gtataagaac	4500
aactgctaag	taaaagtgat	ccaaaggaag	ggaacagatg	agcaacagca	gtcattttgct	4560
cccttcaagt	ccctttctaa	tcactggata	cttaacactt	acaaaacaga	acagtacagt	4620
gaaaacctgg	cctcgtccct	tggtttaaaa	acagagtgtg	gccaggaggg	tacagtggca	4680
catgatgtag	tgaccctgac	ttgatgttcc	agctcacagg	gacctctggc	ctccctgttg	4740
tactccctcg	gctccttagc	ctgcattttt	gttcagcaca	caagtggact	aacacaagac	4800
tccagcacac	agctttcatt	tggctcttatg	ctcagaaacc	ccttctctctc	attfgctaaa	4860
atatatccaa	cggacacagt	agctctccat	tttacaatc	cttaaaatat	agagtatgtt	4920
cataatattt	gcgttatttc	ccattatgtt	attatgtata	tgtccatctc	ttccgctgga	4980
ctgtagagcc	ttatatatta	ttagtgtatca	ataacttctt	gtggattttt	cttacaggga	5040
tatcaatcat	tctgtcttac	tttatgtatca	ttaacctgtc	ctacatttaa	gtactttata	5100
agtatgagct	atttatagtg	tgtggggctc	ccaagagaac	tctgatgttt	gttagtcatg	5160
gatagagcta	gttacattgt	gtccttgtct	gtttccttctc	acattctttt	ttttttttta	5220
attgtcaaag	tcattgattct	ttttgttttc	tcttttagat	attgctatat	ggagatttgc	5280
caaaaaataa	agaaaatgta	atatatctag	cgaatcatca	aagcacaggt	ttgtatttca	5340
tttgatgaaa	tttgggtttt	tctagaaatg	gtaaatgagc	attaatatgt	acacacacat	5400
acacacaaac	acacatatgt	acacacacat	atgtttttaa	gacaggattt	catgtgacct	5460
agaatggcct	catactctct	gagtagctga	gaatgatttt	aagcttgtga	cacacctgcc	5520
ttcatctcca	aggtacagga	attgcagggtg	ccttctttgt	atgcttagct	tatgtgggtg	5580
tgggcatcaa	acccaaggct	tcattgcaaac	taggcaagca	ctgtcccatc	tgagctacat	5640
ccccagccca	tcaaaatgat	attttaaggt	tattttattt	atgagtttta	tcacgtgtgt	5700
gtgtattctg	ctgtgggttt	gagcttgtga	gtacacatgc	ctgtggaggc	cagaagaaga	5760
acaccagatc	ttccctggag	cttgagttgc	aggtagcgat	tagttatcct	ggatggattt	5820
ggggaactaa	actgggggtgc	tttgaaagag	aaatatgtac	tcttaactgc	tgggctttgt	5880
ctccagcctt	taaaatatta	atcttatata	tttaagtaaa	ctaagctagc	ttttgttttt	5940
aacataaatt	tgctgtggat	tttgaatctg	gcttgcaatt	ttattttact	tttttggtgg	6000
ggagggtaa	gttagtaatg	tgaagggtat	ggtttttctc	caggcttctc	ttcttcccta	6060
tttctcaaaa	taacccttta	gtttatttgg	ttttctgtct	ctacgttata	ttttctagaa	6120
tatatatata	tatatacaca	cacacacaca	cacacatata	tatacatata	tatatacaca	6180
cacacatata	tagtgggatt	ggtggaccca	agtgtatata	ttttaattct	taattctaca	6240
tctagcnttt	tatttactct	gagacatatt	cttgcctctg	tgccagggt	ggactcaaac	6300
tcaaaattct	cctgcctccg	tctctggagt	gttggatcac	agttgtatgc	tgccactgtt	6360
tgcactctaa	ctgtgtagtt	gttaagctgc	tatttggtat	gctgggtgct	gactgctcat	6420
ttcctggaca	cagtgtctgt	ataattagct	gtagtctctg	ttgctcttta	atcctggtag	6480
cttaattcta	acttgcattt	tttccgtcag	agaaggcaca	agactagttt	ccagtataga	6540
actgtattta	cttccaatca	ggtcaaatat	atatatatatt	gtttgtttgt	ttgtttgttt	6600
tgttttgttt	tgttttgttt	tttttgagac	agggtttctc	tgtatagccc	tggctgtcct	6660
ggaactcact	ctgtagacca	gactgacctc	gaactcagaa	atccacctgc	ctctgcctcc	6720
tgagtgtctg	gattaaaggc	gtgcgccacc	atgccggcg	ggtcaaatat	ttaaaactac	6780
ctctttttga	atgattattg	atgaattgtg	aattatgctg	ctagtctctg	ggtcccttca	6840
tgagggtcct	cccatgagg	cacctcacag	gtacagggcc	cagccaagag	gccaaatgat	6900
tgccataagt	gtgctttcat	tctccaaata	attcacctca	tccaaaatcc	cacttccat	6960
attttctaaa	ccacggccac	gttgtgtgca	tccgttctat	ggttcgtttg	attgctttga	7020
caagctcagg	ccatagcatc	cgaagcttaa	cagaacccaa	ccgttaggat	attctgaatg	7080
gaaggctact	tgtgtctgat	gcttttattt	tctctccacc	ccattttggt	tgggattgta	7140
gctcttgata	ccgcaccccc	aacacacaca	cagacacccg	gaccatagct	aacacagagg	7200
taaggagct	gaacacggtt	tttctctgtg	tgacacactg	gaaaactgat	gaagctccaa	7260
agcttgatag	ggattttgat	atcagatgta	ttatcctggg	tcattgttcgt	gagactggct	7320
cagactctgc	atttgacttt	gagtcgaacc	acagagtggc	ccagagtggc	tctctgcttc	7380
tagccagcct	gtctgacaca	ctggctggcc	ctctccacca	caatctgacc	tcatgctggt	7440
gagggaatt	tttacctggc	tggcccccag	gacagggcat	gggccatggc	agcatcctgt	7500
cgcagttctg	ttgttcagag	gtggatccca	ctgcaggaaa	ctagagtttc	ccatcaatgt	7560

ctttcttctc	agttttatgg	aaataacctt	tccttaatgg	aactgtaatc	ctacaaggac	7620
aggtgggacc	atgtactcct	gctctcctgt	gggtctcaca	gtaacagggc	ggctgaaggc	7680
atggctgtgg	ttgcctttgc	cttccttttag	agcaggett	caggaaagca	ctgtgagtaa	7740
gcagcaggta	acagttccta	cttgggtgtt	gagatctgaa	gataacgtgg	caagcaagag	7800
gctaagccag	ttcctttctt	agtttagaga	gacatttctt	gactctgcct	ccaggtgacc	7860
tcctctaggt	ctgataacaa	ccagcctttg	gggtttgaga	acattcttgt	tttttgtttt	7920
tggcaacttg	aaaaaagtac	ccagagcctt	tgcatgttaa	gcaagaactc	ttctgcctag	7980
acaagcaact	gtgcaacaca	cacagacaca	agcaagcagc	tcaacctaa	catgctgcat	8040
tccaaatgcc	tttaagaagc	ccacaagttg	ggaacatacc	agaggaatga	ctgaactgcg	8100
ttagaattgc	atgttagtct	acgtgatgga	agaaggcatg	gagtcttctc	taataaaact	8160
gctcatggaa	gactttgcat	gaattccagg	cttcagagg	gttcaaacca	gcaacccaag	8220
gtggatttgg	ttcctgggga	tgacatattt	aggtaagag	agcttaagaa	tcagtcctcg	8280
aaagtgtaat	ttacagaaag	tgttctggga	ctgctgttgg	ttcagcctgt	tcactctcac	8340
tctgttcact	ctggcctgat	gctcatgaag	agttgacctt	taccttagtc	tcagttgctg	8400
cttttacttg	gtatcctgct	gtgtctacat	ccttgctgct	tctaacttgt	ctgctttcaa	8460
cactgttcgc	ttttaaatt	cattcctttt	catttctgt	ctgataagag	atatggttca	8520
gtagatggct	gtaaagctga	gggctctgca	ttctgggaga	ccctgctcat	gtgtgctgct	8580
tctgtgtgtt	ctctgccagt	gggcagagca	cccaggctct	ctgagcctgc	tggttgggtc	8640
tgaatgatata	ctattgatag	aagcttnang	nnnnnnnnna	nnnnntncnn	nnnaaaancc	8700
agtgacgtgc	ctggcatatg	ggtgacattt	agcgactactg	tgaatgatgg	tggggatagc	8760
ccagtgtctg	attcatgggc	ataaacctt	atgtgtcatc	ccataacact	aagaattggt	8820
gagacacgac	tgtacagacg	aggaaagtaa	ggtttatttg	cctgttctcc	tctcttcaag	8880
ttactgaaaa	catggcttat	ggctagataa	gcctcatggc	tagatcactg	ctaaagagtc	8940
ctgacaacca	aattaaatta	ttcttgccaa	aacacaaaat	acagatatcc	atgtgaatgt	9000
aatacattcc	cttacatatt	ttaatccagc	ggtatctcgg	aagcagtggt	cataagtaga	9060
gnacagtatt	gaatctatat	tgtattcttt	gactgtactt	atTTTTTTTT	atttgggata	9120
aggtcttacc	atataacccc	agctggcctt	gaacttacca	tatagaccag	tctggccttg	9180
aactcacaga	gatttgcctg	cctcagcctc	ctccaactct	gagattaaag	gccaaacttg	9240
ccaccattcc	tggcttgtaa	gttacttctt	aagtgttgtt	actaaaattt	ttaaatttaa	9300
ggttatgggt	taggatagtt	tatcttggat	gcatgacata	tttaaaaata	tttatatatt	9360
ctcttaatat	tttttagaag	gtataccgga	ctccaaaata	taactgctta	tttaaatata	9420
aaaacagatt	tattgttcca	taagcttaaa	aattaaagtc	tcaggaaaag	ataccaaact	9480
tggcttttac	ctatttatct	catttatacc	cagaatattc	actggccagc	aaactctgta	9540
gagcagtgat	tctcaacctg	tgggtcacaa	cctatcctgc	ttatcagata	gttacattat	9600
gaattgtaac	agcagcaaaa	tcacagttac	gcaatatcaa	caaaaataatt	ttatgggtga	9660
gggtcaccat	aacgtgagga	actgcattaa	agggtcacag	atttaggcag	gttgagagct	9720
atagccacat	agagccttta	cagggttcat	tctcggtgtt	tctatagaaa	acgtttatat	9780
tagataactt	tctcacagac	ttggttatat	ttccaagaga	tagctgtttt	ataatcccta	9840
ctctaaaaca	attaagattt	ttctagaaag	ttgattattc	acgtgtaaag	agataaaatt	9900
ctaggatatt	tcatttgtat	atgcattatg	aaaaaaattt	aaatgggtcaa	gaattatgcg	9960
atagctgtgg	aaaagtgtcc	catttttaaca	cactttgaac	tccaggcttt	atactgcagt	10020
ttgtttgttg	ttctctcccc	gccccatccc	caactttctt	tcatgctagg	acagaaacca	10080
gggccatgca	cgtgattaga	atatactcct	tcactgagct	gcaccccccc	ccccccccag	10140
ttcttttatt	ttatttttta	ttttgcgaca	ttgtctcact	aagttacctg	ggtaggcctt	10200
gaactcatga	tccttctgct	tcagtctccg	aagttagctg	gatttagaggc	ctgtgctgtc	10260
agcctggatg	taagagtttg	ttgttgattt	aaattagata	ttgtctcctc	taattaaactc	10320
catttgttgt	cattttcatg	gccctgagta	cacttcaaca	gcattccctgt	tcatatcctt	10380
gaattcttct	ctaaatccta	cgagaccttt	cctgatcttt	cattttctgt	tcacatggaa	10440
ggtacctgca	gccatttatt	ctcaagtctt	caaataattct	gcttctcagg	acactttctt	10500
atttctttgt	atttcaccat	agaagttttg	catgacctca	ggcatataag	aacaatataa	10560
tcaaacactg	actgataata	aagagtggag	agatttttat	atTTTTTTTg	tttttgtttt	10620
ttgtttgttt	gtttgggtgt	ttgggtgggt	ggttgggtgg	tttttcgaaa	cagggtttct	10680
ctgtgtagcc	ctggctatcc	tggaactcac	ctgttagacc	atgctggtct	cgagtggagt	10740
ttatttttaa	atacatagaa	tttttagcagt	tattaaagat	aaaaggcagt	ctacatactg	10800
tgagatggat	aggtttgtat	agaagaactt	gactttggct	gaatatttga	tactatagga	10860
tgtagagcat	ttccttgttt	ttcagaattc	atcaggattt	tgatttttga	gatgccagtg	10920
ctaagaatgt	tgtttccaca	aacacattgt	acaatatggc	agaattgtgt	ttagtgtcat	10980
ttcagaaatt	gtttgaaact	ccatttcta	tctaggtcaa	aattcatttc	atggaaactca	11040
agccagtttt	tataaatcaa	gcatttcta	gtaatacaat	caaaagtgca	ctagttttgt	11100
acttacatgc	taaggaatgg	cactgatgaa	atattcacct	actttctgta	acagcagaaa	11160
gctctatgta	tacgaaatgt	acttcactta	atggcacggt	attacatata	tgctagcatg	11220

tgcaagtgaga	agcacgcatg	ttgcatactc	aaaacagaag	acgcaggggc	agctgcacaa	11280
ggcagcggtg	ggcagagaca	cttattcatt	catatgtatg	tggttttgaa	ttaagttttg	11340
ctcatttgctt	atttataaac	ttttattcac	aatttttttg	accactaaaa	tcagtttgca	11400
accacagatt	tcaaaagctg	caaaataaga	cctacataac	tacctcgac	aattgttaat	11460
cacacgagac	ccttgtttgg	gtattgtaag	aattgaacac	tgtatccagg	aagtcattag	11520
taaaaaccta	atgtggtgcc	ttgcttttta	aagatttatt	tacttattta	atataatgc	11580
cctatcatat	atatactgc	atgctagaag	agggcatcag	atgtcagtag	agatgggtgt	11640
gagccacgat	gtggttgctg	ggaattgaac	gcaggacctc	tggaagagca	gccagtgcct	11700
ttaaccactg	agpcatctct	ccagctctta	atgtagtgc	ttttgtcaga	cattgtgtat	11760
atgggatatt	tagctacagt	tgtttcactt	gccatttttt	tccttataat	tttccattct	11820
tcatttaaaa	agaaatatct	cttatttttt	ttacctgtaa	ttaaataatta	ttcaacagtt	11880
atttagtatt	tgggtgttgg	gttacttagt	atttttagtc	ttttaaacat	ttgttctttc	11940
ttttcttgag	tatgtttgag	tccctgcata	tatgcctgtg	cactgtgcat	gtgcctgggtg	12000
cacttggggc	tcattggagg	cctcatatcc	gttggaactg	gagtttagagg	cagagctggc	12060
ttatgggtac	cagacctggg	cctctgcga	aagcagcaac	tgaatcctta	accactgaac	12120
aaaatctctt	cagcccatg	tattttgtac	ctttgtgttt	tatccttgaa	ataaatggcc	12180
ttttaagaaa	tgagaaaagc	ctttaatccc	agcagaggta	agtggatcac	tgagttcaag	12240
gccagcttgt	ccacatagtt	ccaggagagc	cagggctaca	cagaggaaaa	aaaaaatnca	12300
aaaaacagga	aaaacacaca	cctccttgat	ttaagggttt	tttgtttgtt	ggttgttttt	12360
tttttttgag	ttttggggag	ggggtatatt	ttttaatgtg	tctgtagtgt	gctttgtttt	12420
aagcatttta	atcatacttt	atttttaaaa	aaactaaaag	cttttttaag	gctaggtctt	12480
gctatgtggc	cctagtgttc	ctgggacttg	ctctgtacac	cgggttgact	ctgagcctgt	12540
gcgccttctg	cctctgcctc	catagttaga	ttctcaggac	atgttacaaa	gactgtgctg	12600
tgaagatgag	tttttgttcc	tgggagggaa	gggtggagct	gacttgtgag	gtactgactt	12660
gggtctgcct	tacagttgac	tggattgttg	cggacatgct	ggctgccaga	caggatgccc	12720
taggacatgt	gcgctacgta	ctgaaagaca	agttaaaatg	gcttccgctg	tatgggttct	12780
actttgtctc	ggtaaaactt	gtctttgccc	ttttatttca	aaactaacac	catttaatga	12840
aactatatct	gatttttttg	tttatgtgtt	tgttttatgg	taccctgat	tgaacatggg	12900
gtcatatgtg	tgtactgag	tgacagcctt	agttcagaca	tttttttaag	cgactttttac	12960
tagtattttt	atttagaatt	ctatatgtgt	gcacatgcat	atgtgtgctt	gtgtgcacac	13020
gtggatgcat	gtgaggtcga	aggacaattt	ctagtacaag	tgtgagtgtc	acttttttagg	13080
caccttccac	ctttattttg	agacagctct	ctagaccttt	gctgagttgc	ccaggctagc	13140
cggccagtga	gccctgggca	tctaccggtc	tctgctcctt	tacctttact	taggttacia	13200
gtgtgtgctg	ctacgcccag	ctgtttacta	gattctaggg	atccaaatgt	gggtcctcgt	13260
aacttgtgag	acaagtactt	tccaaactga	gccacctccc	tagctcttct	tcacgggttc	13320
tgatgggtgtg	tgtctagatg	gctgggtgtc	cgtatattta	agtccagtag	cagaaatata	13380
aatacctagg	agtcocaaatg	aaagctacaa	gtgcagaatt	gacaatcggt	aatgttcgga	13440
aattgattca	aaagtagtta	gtgagtgaac	gacaggagct	aaaagcagac	tctgagctca	13500
gagtgtgaag	tgtggagaaa	tgtgttttct	cacagttctg	aaggctgaaa	gtctccccc	13560
ggtcaggatg	tgggtgggtac	tgctgtctcc	caacacccac	ctctttggat	tatagactgc	13620
agccttctcc	ctgtgttctg	agccggcctt	tcccacatgt	ggacatcctt	ggtgggtgtt	13680
ccaccagcag	ggcctcagct	agtgccttta	tttcaactaa	ctgtaatgat	tttcttaag	13740
accctgtctc	catcacagct	cactgtggaa	gctgaagctt	caatgtaaga	gttaaggggg	13800
gagggggaaa	tttagtccat	aatgggtgtc	caccaatctc	tgtagctgag	tccatgattc	13860
agttctttta	aggctctgag	tgtagacatt	atcttaatta	ttttgccc	ttatgtatta	13920
tctttaattt	attttatgta	actgaatgcc	tgtgtatata	tgtttctggt	tcctagtcca	13980
tatttttaatt	ccttaaagga	tggaggtgta	gacttttgtc	tttttaattt	tctatccttc	14040
cctcctggcc	tctgtgggcc	tcttttacgt	atttattatt	tttaatttat	tttatgtgtt	14100
tgagtgtttt	gtatctatgc	attcctgggg	cccatggagg	tcagaaaaac	acattaggtg	14160
gcctacaact	gagttatggg	tggtttgtga	ccatgggggtg	ctgggacttg	atcaccagtc	14220
tctgagaaga	ctcgtgtctg	ctgagccttc	tctccagctc	tgggagtggtg	gatattttta	14280
ggatactttt	aattgacttg	gtgaatgaca	gtagaaaatc	aatgagttag	gatccatcgg	14340
aaaaagcttt	tgaactaaat	cttttaagaa	gaaaaatatt	taagtgtctaa	caaaattaaa	14400
tgtgtatttt	ccatgatgca	gttttacttg	ggctctgtag	aaataggatt	ttcaggtaca	14460
tattgtatat	atagttggca	atatttaaat	actaactgtc	gcttgagttc	tgaaatgtag	14520
ttttatgttt	tttactcatt	aggagtacag	ttgccttaat	aactacggag	attagttatt	14580
aaagaataat	tgctcttctt	ttttcttttc	tgtgtaccag	catggaggaa	tttatgtaaa	14640
acgaagtgcc	aaatttaatg	ataaagaaat	gagaagcaag	ctgcagagct	atgtgaacgc	14700
aggaacaccg	gtaagtgcgc	cgccttttat	tctcgaaggc	agggttaagaa	gttaagttct	14760
taagtcatct	tgaaaatata	ttaccccatg	tggagcaatg	gaactgggtc	ggggttttgt	14820
tgagataaag	tgctcctctg	cctgtaggta	agattgtctc	agggtattgt	aaggtttctc	14880

ctgagtaaca	gtcagcatgg	gctcgggacg	ggcaagggca	ggccttagtg	tgcagaggat	14940
ggagctcact	gaagcccca	agagtttagt	ttcacatgag	attcagttct	agaagaagtt	15000
aaattgcttt	ctttctgtgt	aaatttggat	ttttattgta	gaaattaaag	tttgttttct	15060
tttaaaacaa	acacaaaccc	agagcaaaga	gtctcctagt	gaagagtcac	tccgtgtcag	15120
tattttacac	aactgttttt	ctgtaaaggg	ggaaaaagaa	ttcaaactct	ctctttcaag	15180
aatgctgact	gctgccaact	gcctctcccc	gtggccccct	tctgtataga	caggcatagc	15240
tatggtgagg	acttggggcg	ctcttgtcct	tctcctctct	ctgcttctct	accttttctc	15300
tcgtgcccct	cacttaccag	gccctgggaa	gctacacacc	aggcaacagt	gaccagggcc	15360
tcggcctggg	cttcgaccaa	ttactagagc	agaaacagca	gcagctgcag	tggtgttttg	15420
tgctgtgcac	tgtattaggt	tggtgttttc	tcacctttgg	gttttgtgat	gttttgcata	15480
agtcctggta	ccattctagt	ttttacattc	tgggtagata	gagtttatct	aaggctctca	15540
ggcatatgaa	tggaagagct	cctctttaca	gccattcgtg	tagcatgcat	aactgctcct	15600
ctgtattctc	tctagtgtct	ttttttttgt	gtgtgaatct	gatgtcttgt	tattcaccta	15660
caatgtggag	taatggtcat	aaacatataa	agtacttatg	cctttatctg	ccaaattgta	15720
tttaactttt	cagcttttaa	tataactttt	tatataataa	ttaattttatt	ttaaaaaaa	15780
ttgaatacca	gcctgttata	gtggcatatg	cctgtgttcc	tagcactcag	gagacaaagg	15840
cagaagtgtg	agaacttcag	actcatactc	agctatatac	aagaccccaa	atttgtgcta	15900
gattctgcag	tacagccatg	agtgtcccca	tcttagaggg	agatcgctca	tccttgtgct	15960
gttctttaag	tcttaccctg	caaccactgt	taagtacact	cttgctcaca	gtcctttaga	16020
atctcacact	ctttctcttt	acagacacca	tgctattgcc	cactttatta	tttatctgat	16080
gtctacaaag	attatgaaag	agaaacttgt	atgcattctg	tgtaaagtac	ttgacacaaa	16140
taatagtatt	caagaatgac	ttcttaaagt	aacactgaat	gaatagtttg	ttctaatttt	16200
tttgatcaac	aaatcaaaaa	atatttagat	taaataatct	agatacaaa	cataatacca	16260
catgaatcat	taaagttagt	aatcaatctt	ataagttagt	gacctaaaa	ctcatagaca	16320
ttaataattg	ctttcattgc	ttagatataa	actttattga	ttaatacgtt	ctcatgaaag	16380
tggttcttgg	aaggttctgg	aaacgaaaa	atttttctta	ctgctttttt	cttctagtaa	16440
ctgattgaat	ttttctgcag	ttccataaag	catctggtca	attgctatta	tccaatatga	16500
ggatatataa	caaagtattg	attttttaaa	ttggcgggtg	taagacaaga	ctgggcgtgt	16560
gaatgagggg	gtctctgttt	cttgtccctt	ctcttgggtt	cttttctttt	tggtgggttt	16620
ccttctccag	ctgtctatgt	atgggttctg	attatcttat	tatatcttat	ttgtttattt	16680
ttcattgtta	cttcttagaa	gccaacatgt	tatatcacct	ccactccac	cattaggtgt	16740
ctcacaaata	ccccaaagta	aacaaccaca	tcatgtcatg	ttctgtatac	ttccataagt	16800
gttggttaact	ctactgactc	ttgtgagcag	gcccaattgg	ctttatccct	agctgggtga	16860
cctgggttcc	tcccaaacac	cataccgtcc	atcaaactga	gtccttttcc	aagcacacac	16920
cagatactgc	tcacttgagg	actcttctca	tcacctaag	gactgcctgc	tcctcggcag	16980
aaagggcctc	tagtcccata	cccttacgcc	ctcaccaatg	ccttaggaac	atgtgtctca	17040
tgccccgtgt	ggtcatttcc	gtttacagta	gggaaatttg	cctgataact	tgacgacac	17100
ctataaagag	gccttgcttg	ctctcatatt	tagctggaga	agataatgta	ctcaccaact	17160
ccactctatg	caaccagctc	tgctctgccc	atgccagtc	gacgtgaatc	ttacacctgg	17220
attcagattg	atgaatctac	aacatcacc	actccatgct	tccttctaaa	tcagcagttc	17280
tagcctgaat	gacagatgct	accctaagtct	catctagtta	gccctgtccg	gagtaacctc	17340
gaccttgagg	attagaccag	gatgcacatc	ctgcaccagt	tcctttgtgc	cacctgactt	17400
catcccacc	gggcatagc	ccatgctcag	gctccaccct	ccatgcacaa	agctggcttt	17460
tccagcttcc	ttcacctgta	tcagacacaa	atagcaaaag	gggtccacgt	gcctagggtc	17520
catcacaaga	ccatgtgcgg	tagtttgga	aacagtctcc	acttgaggct	cagatagntt	17580
ggaatcttgg	ctctcatgta	gttgtagtga	ttagatcagt	ttaggaagta	tgacctttnt	17640
ggaaagaaca	tataactggg	aggggctttg	agatgtaaag	gccccacaca	attcctagtt	17700
caaactntac	tgctgtctca	aagcttgagg	catgaactct	cactgttctc	gatgtcatgg	17760
ttcctgtctg	cttcacaaat	tccctatcat	taggggtccc	tttcttctct	ggaattttta	17820
gataaataaa	cacttctttt	aaaacaacaa	gaacaacaaa	tctgacnctg	ataatggatt	17880
ttaaggcgct	ttctctggat	aagaaaaaaa	aaagaatata	tttgcatagg	tgctgtatta	17940
cttttgtcat	tggtataacc	tgactggaag	caacttaaa	gaagaagaat	gtatcttgat	18000
ttgtagatta	agagcaccat	gactaagaag	gcatagcagc	acaggtgcac	cagcaagaac	18060
ataggctgct	agctcagatc	tctgtagata	tgggaacagg	gcaggaagct	agtagtctat	18120
aaacctcagg	accatcccca	tggagttcct	tgtcttccag	tgatgtcctg	tgtcttaaa	18180
tttcacagtt	ccpacagcag	cacctgccgt	ctgggaacca	acctgtgggt	gatattttac	18240
aacgtgatag	gcataatttt	tctctagccc	tgtaggttta	tagccatcct	atacttcagt	18300
ttatctagtc	capctcagtc	tgtgtgtctt	atagttccaa	cacttcaaaa	ctacaaagtc	18360
ttaagggcca	tgggctcggg	tttattagag	cagtaacacc	tctactagct	ttctgtgtta	18420
cccactcctc	tttaggtctg	gttgaaatcc	taataggaag	cagcttgaga	ggaggggtta	18480
ttgtggccca	tactttgttg	gtacattcta	tcatgcaagg	gtggcactgt	gatacagccc	18540

aggccatccg	aggatggtac	tgttggctta	catctgggtg	ggacaggaaa	tggttaattct	18600
caaggccac	ctgcttggtg	acttctttca	gttaagcccc	atactctaaa	tcctctacaa	18660
cctcccaaca	taatgccacc	agctggggat	cagctgttga	cagtgtctgc	ccaggggagc	18720
agtttaaate	cagaccagg	gacctgaaaa	cagagaactg	cagaggggct	gtgggacttt	18780
ataccagctt	tcagacaaa	tcacggcatt	tctttgtgag	cttgggttcat	aaacaaatat	18840
atattctcct	ataggctcct	ttagtgggtg	tttcatatcc	acaaatttgt	tcagaaaaac	18900
actgtgtttt	atgctagctg	tgtaggagat	aataccgctg	ggagtcactt	gagcatggat	18960
aagtgcata	gttcgtcctc	atgagtcctt	gtcctgtttc	tgtattatgt	ttacttgatg	19020
agtttagttt	gtcagttggc	caccaattaa	aaagtatcat	tttatttttt	ttacaatact	19080
cagttctcaa	gttaggagtt	ttgttattat	atggcctcaa	tattcacatt	ttaacctttc	19140
caggagttaa	gtataaaaac	ttatatcaac	tgttgactta	gtaaatatct	attacagata	19200
ctatattctt	cttagtttat	atcatgaata	tgaggttgct	taaagtaagt	gatgtaaaat	19260
acactagggg	atgcttataa	aatggaatgt	tgtgagtttt	ttgaaacacg	agtactaaat	19320
tcataagttt	ttaaatagtt	acactgttag	cttcagtagt	gctagatata	tgtctataat	19380
ggctgaagag	tggagcttgg	atattataag	tgtactctgt	atattcatgc	agacatatag	19440
cagattccac	tagtatgtgt	ggttaatatg	tgctaataaa	aatttaatac	aaaagtcag	19500
ttttattact	gggaaccaga	gggttggtt	gtgctgattt	taagtcagtg	actattagca	19560
tattctaaga	aacagtttta	ggattttaaa	gattggcttt	accataaatg	tagagctatg	19620
ttttactata	atccatatta	tggtcggcct	taattcaatc	tctgcagttt	ggttactctg	19680
ctcaaagtga	aggtcattta	taaatgatac	acattttctc	accataggaa	atactacctg	19740
gccaataaca	gagttagaat	tgctaaattg	atggtacca	caatggactc	aacacaaact	19800
aaagtttatt	taigccca	gatgtatctt	gtgattttcc	cagaggggac	aaggtataat	19860
gcaacataca	caaaactcct	ttcagccagt	caggcatttg	ctgctcagcg	gggttaagtaa	19920
agattttaact	gtattcagaa	aaacactttt	ttagaagag	tgatctttgt	ttccttcaga	19980
gtcatactaa	agaatatgcg	tttcttgtaa	gagctaagt	agagaatac	cgatcttcta	20040
cagagttagg	tatattctta	ttagtctgtg	cttgagaggt	tagagacgca	ggcttgcctat	20100
ggcacatttc	ccatgctgtg	aattgagtta	aaaatgtagg	taaatgatat	ccccagaaa	20160
gtatactttt	ggagtgactc	agtataaagc	ctggtgttat	aacataaaca	cgcacgtgcg	20220
gatgtatatg	tagcacatat	gtaaaacacag	gtatatgcat	tgtataaaga	aagtggagggt	20280
cggggccac	tgcaggcaag	tcttttagtg	atgctgagct	aatgctgaga	ggtagaaaga	20340
ccaagaaggc	tggagttgct	cattcggcaa	aggtcagagc	tcactgtgtg	ccataactcg	20400
agtgttctgt	ctcccttttg	atacagtttt	cttgttttta	attattagtt	tttacaatta	20460
tcccataaaa	tgtgggctca	ttgtggctcat	cgttttcata	aagtccttca	agtatacacc	20520
cagcaagtat	ctaaatacac	tgggaagaat	cagtcagctg	atggccttgaa	gtttcaggac	20580
atctagtgcc	acatcatgct	tcagaaccga	cctgcactta	gtcaggggtca	tattcatgcc	20640
acgtgaagac	gagaggaggc	catgccgtct	gacttaggat	ggaaatttcc	ttcagacaaa	20700
cagcaacggg	ctaggtctta	gttataggca	tagtgtctgt	ggttatacta	ggcagacatt	20760
agtggactgg	gtgttagaag	gtacagacag	gcaagaattt	gctgtagatt	tgtttccctc	20820
atgtgttgac	accacatcta	acctgctttt	tgagcttcta	gtcctaataa	tctcataaaa	20880
atactggttg	aaccagaaat	ggtgttgcaa	agctatgac	ccagctcctg	ggatctaggg	20940
tgggaggatc	ataaatttga	ggccagcttg	ggtctgtcct	agagaaaaaa	gaaaaataaa	21000
aagtctggtc	aaggttaacat	ggagcctgga	gttttcacag	gggtgattctg	taaaggtcct	21060
gagacaagat	ggcctctagt	ggcgaatgac	ttagctgaca	agaaaacttt	cccagcttgg	21120
ttgacttttc	agacttcata	caagtttgtg	aataaattac	actccttctg	cccttgggac	21180
tgaactcaga	tatgtggttg	tgggaatggc	tttctttccc	acaccacctt	gcattttaaa	21240
aattcttctg	tagacagtcc	caccatcctg	tagctgttct	tccttatgtc	gccactttcc	21300
ctggagagag	gcagtgcaga	cttcaaccg	cttctcccta	gtcgtgtgtc	atagcacatc	21360
gaaagacctg	gtgcttcctg	tgaaattgta	agtacatcct	ggagtccagg	agaggaggaa	21420
gccgaacaga	gtggaggga	tgctgagttc	gttcctaaga	aagactgcgt	gcttagcaag	21480
atgctgctgc	tctcctgtcg	tgtctttctt	gtcagaactt	atcaaagaga	aggctcgcag	21540
tgggtcataa	tcttcccaag	gaccagcctt	cccagcttct	cgcagcatat	ctcattcatg	21600
tagatgttta	atggatatgt	gtcaatgggg	ttgacctaag	tgagatggca	atgtatgtga	21660
gcattctagg	tgtgaggtta	tgccattaaa	ctttaatttc	cgtctatttg	tggtagttga	21720
taagtaattt	agatgttgac	tttcatgtat	tcctaattat	gaccacattg	aatctacctg	21780
ctttctaggg	cttgcagtat	taaaacacgt	actgacacca	agaataaagg	ccactcacgt	21840
tgcttttgat	tctatgaaga	gtcattttaga	tgcaatttat	gatgtcacag	tggtttatga	21900
agggaaatgag	aaaggttcag	gaaaatactc	aaatccacca	tccatgactg	gtaagtcctg	21960
atttccatag	aagctgaata	gtacatggta	caggtaagat	aaactcctgt	ttgttcgctt	22020
tgcttagctt	ggttcagttt	ggttttcagt	agaggggtcc	actatgaagc	tctggctggc	22080
cgggaactca	ctatgtagac	caagctggcc	ttgggctcca	ctacaccag	caccaatcac	22140
ccactcttat	cttttatgct	ttttgttttt	gctttgagct	ttctttataa	catgtttggg	22200

aaggacattg	tcattattta	caagaagaaa	tatggctctt	tcccaacatg	ctagaattta	22260
aagactcaga	actcttgcc	ttgtcagtga	caaagtga	atggctgtga	agtgcagtgg	22320
ctttgagtga	gaatagttca	ggtaactata	gccacagact	caacatttga	acatgggaac	22380
aggtgagaac	ggagtgtgg	aagattctgg	ccccttcag	agaattcatt	ttagagagag	22440
atgagagtag	taagggaag	agaagagaga	gacgtggtat	tttgctgcag	actaaagaga	22500
tctcttataa	tcgcagtact	aaggaggaag	aagcagaaga	tgatgactac	agggccaggc	22560
tgaacaatct	agtaaaatcc	taagtcagga	agtcagggct	gaggtgcagc	tcagtaggag	22620
agttgtgtgc	tgccctacac	aaggcctgga	tttagctccc	agtagcaacg	aagggaggcg	22680
aggggtgggca	aaatcgaaca	cttactcttg	gagactccct	ttatgaatat	taccacactc	22740
cagtaaatac	tctccagaga	tttcagatga	gattctgctt	cctggtaaac	aggaggccaa	22800
gaatattatg	tcacactgaa	catgggatgg	aagacatggt	ctgaggaatg	tctgcactcc	22860
agtgtgatga	agacttgaag	tttagggaca	ttttccctcc	ctggccccac	tcaccccatc	22920
tgtattgagt	attcccctag	tgctcatctt	tatttgtatg	ttacttttca	ggaaggggaa	22980
gcagattgat	attcaaaccc	agccagtttt	cttaataact	ttgtggatgg	gattggcttt	23040
gacagtaaat	gaggaaatgt	aaaatgtaaa	agattctaatt	ttttaattatt	ttaaaggtga	23100
ggttttctgt	tagtacgcag	agtgagaggt	ttcttactga	tgctctgcgt	cctagaggaa	23160
ggatggctac	ttctccaagg	cttgctgtta	gaagtcagt	acatgggctt	aacaagagat	23220
atgtgcta	gaggttttaa	tttcagctta	atactgcaaa	tcataagtgc	atagctttat	23280
tggttttaaat	tcttttagtc	ttaatgtttc	atttttacca	taagttactt	tgtataatca	23340
caaattctaa	actagtaaga	cgtgaaat	tcttctctt	tgtagagtt	tctctgcaaa	23400
cagtgcctaa	aacttcatat	tcactttgat	cgtatagaca	gaaatgaagt	tccagaggaa	23460
caagaacaca	tgaaaaagt	gcttcatgag	cgctttgaga	taaaagatag	gtaagtggta	23520
agagctccag	catttagaaa	gtgcagttca	accaaatttt	actctcagat	cctgcttgaa	23580
aggagtcttt	ttatcttcat	tatttagtaa	atactaata	tacctgcata	gacaagacca	23640
catatactta	aatgtagcat	gtttcatggt	gcgttaccct	tgtttaacaa	ttagtttaa	23700
cactctacat	cagtttgcc	gttgatttct	gtaccatgac	aactcaacac	agcgtgcgt	23760
ttattccaaa	gtcgatagca	cagcaaaagt	gaaactaaag	tctgtattgt	ttcaagaatg	23820
ctttttgtga	actcgggtta	aatcttattc	tatctttcgt	tgttcacatt	gtacattttc	23880
atgagtcact	ataaaaatca	tgacatgggt	gcctacctgc	agtgtttgct	ggacagttag	23940
ctgctgtgtg	ataagagcct	ttcctcttca	gctacacggg	ggacacgagg	ctttggggtt	24000
caagactgaa	gcacgggtga	gcacaacacc	tttgtgtgt	gggaagggaag	ggaattgttc	24060
ttttcataat	gaaattgtcc	cccttcttga	gttagtagaa	agtattacaa	ggatagagag	24120
ttgaaatgaa	gctttatatt	agatttatgc	cttggtgtgt	cacgtgtttc	tacctgacat	24180
aacttttcaa	cccagccgct	caggattatt	ttgatgatgg	gaacaatgta	agaaggccta	24240
tgatctggta	actcactgtt	gtagctctgt	ggaagcggct	cacaggcagt	agggacgctt	24300
ctgtgctttt	gtgcctgtcc	tgctgttaga	atcttacaga	ggaggatgaa	tgaatgaccc	24360
tttttatttc	tcttgtctgc	ttttctaatt	ttatgggaat	aagaactttt	ggtaggtctc	24420
tgctactggc	ctcttgttgt	gaagagacac	cttgagcaaa	gcaactcttc	tgagagaaag	24480
catttagttg	gggaattcct	tacagcttca	gaggttgagt	tcgttttcat	catgctgagg	24540
acaggagggc	actcaggcag	gagaagtagt	tgagagccac	attctgatct	acaggcagag	24600
agagacagac	tgagcctggc	atgggttctt	ggaacctcaa	agcctctcat	ccctacccca	24660
tctcccgacc	cttatacaca	cttcctccaa	caaggctaca	ccttctaata	cttcttaaag	24720
agtcaccaca	ctcagcgact	aagcattcag	atatgtgaac	ctgtcggagc	ctttcttact	24780
cagatcacct	caggaggaaa	actcctatgc	tataagaatt	tcttttcttt	cgcactcttg	24840
aaagcttggt	tttgtgtgat	tagatcctgg	cctcacacat	gctcggcaat	cattttactg	24900
ttgagctcca	gcctcagccg	ttttcattgg	cttatgggat	gcgagccatg	ggagagaagc	24960
tagaaggcct	tttgttttat	gagtcgggtt	ggtggaacca	cttacagatg	gaagatttac	25020
aaacaaaaat	gaagctgggg	ccatcaaggc	tcagcactcg	ctgctcttcc	agagagttca	25080
ggttcatttc	tcagtaacca	catgggtggc	ttgtaaatgt	aacttcatat	tcaatgaccc	25140
tgacaccctc	ttctggcctc	tgtgggcacc	agacacaatc	atgggtataca	gacacacaca	25200
ctagccaaca	cccatctaca	taaaagtata	taaacatatc	tttatcttaa	aaatccccga	25260
agtcctcatt	aaatatctta	gatccccgcc	gtgttttgat	ttttgtttcc	cacgtgggtga	25320
ggatataata	tcattgtccaa	actgtaagga	gtgaatgcc	tcccgctgct	ctcggacacc	25380
tctgcactca	tccaagtttt	ctaaggagct	gtacttgctc	agcaagtact	caatacctaa	25440
taaatgggtt	atgtttgttt	caacaccaaa	aatgtccaaa	actgaaagat	caattctgtt	25500
gttttccttc	tggccatagg	ttgctcatag	agttctatga	ttcaccagat	ccagaaagaa	25560
gaaacaaatt	tcctgggaaa	agtgttcatt	ccagactaag	tgtgaagaag	actttacctt	25620
cagtgttgat	cttggggagt	ttgactgcgg	tcactgtgat	gacggagtc	ggaaggaaac	25680
tgtacatggg	caactgggtg	tatggaaccc	tccttggtcg	cctgtgggtt	gttattaaag	25740
cataagcaag	tagcaggctg	cagtcacagt	ctcttattga	tggctacaca	ttgtatcaca	25800
aattaaataa	ggagttttct	tggtgttggt	tttttgggtt	tggtttgttc		25860



tgttttaagc cttgatgatt gaacactgga taaagtagag tttgtgacca cagccaacat 25920  
gcatttgatt tggggcaaac acatgtggct tttcaggtgc tgggggttgc ggagacatgg 25980  
aagctaagtg gagtttatgc tgtttttttt ttttttttaa tgttttcacg aattaatgtc 26040  
cacttgtaaa gattattgga tactttctgt aattcagaag gttgtatttt aacactagtt 26100  
tgcagtatgt ttcgctatat tggttatcct ccatttgact acttggcagc tcagactcct 26160  
aataactaaag tattttacat tttgaagcta tgtgatactg gttttttgtt gttgttggtg 26220  
ttgttaattt ctgaaagtca atgaaagaca ctgtaatgat gcgttaagat gttccaagaa 26280  
aaaggtgaga attattcatg gcaaaaaaga tctgtctagt gtatattttt attatattgc 26340  
tctatttagc taattttcct tatatttgca aaataatgaa catttttaat atttattaaa 26400  
atgcttgatt tgcatacccc cgattctaca gagaataatg tgtaaagtgt cagaatagac 26460  
ttgaagctct gctgtgactc agtctccttt gtcagagctt ctagtagccc agctactgag 26520  
ctgctttgtt agtacctcca gcacctgagc cgtaagtac ttataaatgc aagggaccgc 26580  
ttatcttcat atcggaatag acatgaacag agctctaagg cgatgaaagt ctgccagcat 26640  
cctctctgtc ctgcacagtg ccttctgcct ggctccattt gctttggcac tgcgttcgat 26700  
ctagagtgtg ggtgctcact gcttatttca gccctggctc tgtggttttg tgtcctccag 26760  
tgggtgctgt cactgttggg gtgcaggttg tgcgtccctg actcagaggg gcagctccct 26820  
ggctcctgag ggtgagcctt cttggctact acagaagtat tgtgcgtttg tgtatggcaa 26880  
gaaccatcag gattggataa atgtgttatt tctctttgat ttccatggag ccacactggt 26940  
ggtacatgtc cctgtgtaac agagctacct ttcaggagca catcatactg tctgtagtc 27000  
cggcacggtg tgtcctgtga gaagaggct tctaactgtg gatttgcctg gtttctatgt 27060  
tgtgatttaa gctgattgac ctactagtca tccaaggtaa cattctgca aatttcatac 27120  
agatttttgt cacaataa ctataccaat gatctagttg aaatagacca attgaatcac 27180  
aataaataat tttttttaat tgagggaata tttgcttctt gttttttcaa agccagaaaa 27240  
cgagccattt caaacatctt tgaagagtca tgtgctgtca cttgttttct atgtgttagt 27300  
gtctatattc atgtatggat acacatgaac atgtatatc atacacacac gccaatagaa 27360  
tataacagcc taaaaacaat ccagcttgtg tatcatgtta ctgtgctgaa ttgtaatgg 27420  
ttttacttac aaagtggagc taaaatcgat ttcatgtctt tgttaaatac gtttttttca 27480  
gcaatcctat tagagcttat tttgaccaga tcaaaataag tacaagttca gagactttaa 27540  
atatggctga ggtctagagc gatagctcag tagttaggaa cacatgccac tctttcaagg 27600  
gcttcagttc ccagcactca tatggaggct cacagaaggc tggaattcca gcttcagga 27660  
attggacaca tctcttagct tccatggatc tgtctgtctg tctctccctt ctctctctct 27720  
ctctctctct ctctctctct ctctctctct ctccctctt aaatatcatg gatatgctgt 27780  
gcattttaa ttttaagacac agaaccattg gaattacatg gattatagct gattctcttt 27840  
gaacagggca cagtgttctg cgtaagatct cttgatcatt agcactggac tcaactctct 27900  
cacaagtagc ctatcaaatg tggatttaga aaatacattg tgtcaaaatc tttgaaagat 27960  
gagaagaatc tcctaacaat gtttattttg acttgacatc actatttctt gaaaattaac 28020  
tgtctatgat tcttttca tagtgtaaga tcttacttgt atcaccatca gcttgacgt 28080  
taggggctgc agttgttctc cttcataaga ctgccatccg tgtgcatgct tttatgtttt 28140  
tcagaaagga tgttgggatg aaagtaagaa aacaaagtct cttcttgtct ctcagtctg 28200  
tgatcactag catttcacaa ctcagggtat catccatttt ccagcagata aaagggttag 28260  
cgattaaccc tgcattctga gtttagaaag ctacaatatt ttttaaatat tgagcaatga 28320  
ttttaaaaaa atacattgga ataccccaa tttgtaagca atccaaaagt tggagtgat 28380  
aagctaattt gcctacttta aaggatgtga cctcaccaca ggaaacctgt aggtattact 28440  
taacaaggct ttacatgaaa atgccaccgt ggccatttct taaacactgg tggcttcttc 28500  
cagatttcat ttctatgttt gtttgtttgt tgtttttttt ttacttagat tgctgtgagg 28560  
tttttttttt ataacaaata tacatttttt tctttgtcac attacatgct ttgtcaatca 28620  
aatgacctaa ctagggtggc tattaagaaa actacatatt gaaatctgcc aaaatgtcgg 28680  
cataaacaac ctggctccta attgtgtacc agatctacat ttgaaagaac agaaatgtct 28740  
cacaagacaa taaggctcata tgtaaaacac taaataaact ttaacctcaa caattgtttc 28800  
tgaagtgttg agattaaaga ctgagtgttt gcggaacgtt gacatgtcca tggccaggct 28860  
agtttctcgt tttctttttg tcttaagact aaacattggc tggcttaaaa tattaccagt 28920  
tctatatagt ttacattata gacagaatat ataacattta agtattagta tgaaaatcag 28980  
tactttgggt agactaatat ttggaatatc cagatgattt gatattcatg aggtaaagta 29040  
agtattgtg tgaactgact aacttaaaat ctcttattca tatatcatgg ataacagctg 29100  
ggagtgtgta cacatggctg ccatccaggc actcggaaaa tccaggtttt gagaaagaga 29160  
gtgttttcca agtcagcctg gtctatatag caagcttcag actagccagg actgtgtacc 29220  
aagatcttct tcacgccacc cacacaaaca agagaagtta tatagagaat tgcttgggat 29280  
ttagttacaa catttttgtt aggtatttcat ttaattggga ggggtgggg gaggtagcag 29340  
tttgcatttt cagagaatgg gttccattcc cagcatccac agggcagtaa ctgaagtaac 29400  
tgtagttcca gggatccaa caccctcata tggacacaca cgcaggcaaa agaccagcat 29460  
gcataccatt aaaatgaatt attaaaattt ttttaaaaaa gacttttgata tatttttagt 29520

ttgtgtatgc	tgaggtggat	ctgttgctct	tggcaactg	agtagtaata	cgtggagttt	29580
tataagaggc	cagagatctc	actttcatat	aataacctt	ggataataac	ctagttgcta	29640
cctaagcaaa	gaagttcctt	ggaagagcta	catcacagaa	aattggatc	agagcatata	29700
ctcagtcct	ggttatgcca	tgagacaggc	cagtggttta	cccagactag	tgttttgtca	29760
tatctgaata	aaataaataac	aaagtcgtat	tctacaactc	taaatectca	tatagaaaat	29820
aaaataggaa	tgcggttgac	acaatttggt	gcctgaagtg	atgaggaaat	taagttacaa	29880
agtaatcaag	aagtactcag	ttactgtgtc	ttcttatgta	gcttcagaat	aagttctgat	29940
agtgttactg	ggtttcctgg	tgctttataa	tccagtacaa	gggaaaatat	gaacctcatt	30000
tttttttata	aaaatttggt	gacaaaaaag	ttacattggt	ttataaaagt	ttttgtgttt	30060
tatgtgtata	tggtgtgttg	attgccagat	atagctataa	agctacatta	tttcagatgg	30120
gtattttgtt	tgagatttgc	tggtgtgtcg	ctgtatgctg	actgcttctg	cacctcttgt	30180
aggcaccagc	ctgtctgcag	gcagaccacc	caccccatgt	tggttccttc	tgctctctgt	30240
tagccatcca	accttgagca	gctttaaata	aagagtgtcc	caggtttatc	acctgtagaa	30300
cattgtctacc	acctgtagaa	cattgtctacg	atgttagcca	caaagaattg	ttgagagcat	30360
gtgaactact	gcagatgatg	cagcagatgc	agagcataga	aagaatgtag	ctgggtgctgg	30420
ctcgggttcc	tgctctctgt	cttcatagat	gaccttcagg	tcagatgggg	tttgataaaa	30480
gtctagggat	gggatgccat	ctggctcctt	cctgtttctc	aaggagacag	tggtgtgtgg	30540
aagtggatgc	tcgtagtatt	ggttcagatt	gtaggttttg	tttaagattc	agatcacgtg	30600
gtgtagagag	atacaacaag	agaaagaaca	tctccattat	aagtcacatg	gcctctcaac	30660
tacagctcaa	tagaagctgg	atcttcaggg	caggtagtgg	gctcagtcct	cacactttgg	30720
gcacaggtgg	aggggatggc	atggcagctg	cagtaatgtc	tgaggggttc	cttctggcat	30780
tccgcactat	ggtacgtttt	agactaagtt	cgagacaatg	cctatcggag	tgccccctta	30840
tccgtgtagt	ggggagtgtg	gaagtctgtg	gagaggctag	tggtgtgact	ccatggagtg	30900
tggtttccca	tatgtaaaag	tgactcagc	tgagctctat	ttgtcgaaag	tgtaggggta	30960
catctgttca	ttattttctg	acatatatta	gagctcagaa	ttgtcaggaa	tgttcatgat	31020
taaacattgt	tccttgtcat	ttggaacat	gcctatattg	gcttctgtcc	tatgcctgca	31080
cgagttaact	ttctgtctgg	gccaggagtc	gttttgattt	ttacaggcat	ctaacatatt	31140
tctgtttgtc	tgataccccc	ttccttctgt	tttcttatct	gcactgtctc	ttcagttctg	31200
cgcagaaaaga	gcagagaaaag	tccttctgtc	tcccaagacc	tcccatgggc	tttgtgtctc	31260
tgctgaagac	aaaccaactag	tgtggcaact	agtttgatat	cttttctctc	tataatagaa	31320
atgagatcat	ccaagttacg	ctggagaact	gggtgtgtgt	tctgtgtctc	tgatttcccg	31380
ctaccccttt	ctcttagaga	aattcattgt	agtttggtta	gttggaagtc	ctggtaagggt	31440
ggcaggtcct	gtgttatagt	gagggcctta	ttgtctgggt	gtgggactcc	aaagttagtc	31500
tcaagtcccc	cgaggactca	acctagtacc	caatctttcc	aaccaccctt	tctttccaac	31560
cctatgacat	taaattttca	gtgcttcaat	tttgagagaa	cattcagacc	atggtaaat	31620
cacatgagca	ccaagtggga	ttcttagggg	aaattctgaag	gaactgcttt	taaaaaataa	31680
cagttttcct	gatgaggttg	cattggaacc	agctgtttag	aactgaaccc	cctcctcccc	31740
tgtgaatggc	ttgaacaggt	ggtttatttt	ccagaagcaa	catagagctg	gattaatgtg	31800
ttattagggg	actaaattat	ctgttgacta	tactgttaac	ctcaacctca	aaatgtagta	31860
gtggctacca	gaccacccca	acacctctct	gattaaagcc	atcaagagga	cagtagagga	31920
cagaagaatg	ctattttccc	ttcaagacat	ttccagagag	tccctaattg	ttctacttag	31980
ctgtgactgc	ctggaaactta	gtacaggctt	catttagctg	caagggaagc	tgacaagtga	32040
aatcttcagc	tttgacatcc	tcaataaaat	tgaaggtctg	agggatgaaa	ctattctgag	32100
aacagtggca	gtgttcacac	catcttctga	atataaatgc	tgaagattac	taaatggata	32160
tggtgaattt	taccacagtc	caaaccaaca	gagcaggctg	gggatgtggc	tgagttggta	32220
cagtacactt	gcctagcata	gaacagctgc	tgtggttaaca	aacattgtta	caccaaacac	32280
tcggtgtgtg	gcatcaggag	aattagacat	ttcaaggcca	tcctcagctc	tgcatgtagt	32340
tcaaggccac	ctgggggtta	ctgatttcca	ccttactttt	tgaggcagtg	tcactaaacc	32400
tgtagttcac	tgatgaagct	acagtggccc	acaaactcca	gcagccctcc	cagggctgtg	32460
actaccttta	aacaacaaca	aaacaaaag	tgctggagat	ccaaactcaa	gttcttatcc	32520
tcagggcaag	cactttactg	actgggcttc	caggcctgct	ctattgtctg	ggcaagactc	32580
gctataacta	cctacaattc	ctcttcaaca	tctatttgga	ttctcttttt	ctttgttttt	32640
gagacaagggt	ttctctgtgt	atctatccct	ggctgtgtct	gaactcacta	tgtaaaccag	32700
gctgtcctcg	aactcaaaga	ggtctgcctg	cttctgcctc	atgagtgtctg	ggattaaagg	32760
tgtaccacca	capccctacc	tattgggttc	ttaaagggca	ctttccaaaa	ctgatgaagg	32820
agagttaaaa	taaggagaat	ttttgtaaaa	ctaactgtta	atgtgaactg	tgaatgtatg	32880
ccccaaaate	ctacatatct	acaaaatata	caggtttttc	gttatttttag	tcaccttctc	32940
tggttggtct	ggttttactc	ttgagtaatt	tttactaggc	aactttcagg	acagaatgac	33000
agttcttgaa	gtttaatctt	agtagaaaaa	caggcaatag	tgagaaagct	gtgtcacagt	33060
gttggtctgc	aaccactcca	gatctattcc	ctcccaatgt	ctatggagtg	cattgtctgc	33120
ttaggtgctg	aataaggaca	tgccaacatt	tcttatctac	tgtaaggcaa	aattgggtgag	33180

cagtgtcact	aagcctgtag	ttcactgatg	aagctagagt	gcccactagt	gagccactat	33240
cactagttag	cacactagta	atagtgtgaa	agaaagtgtg	ttccactgtc	ccatgaacac	33300
accacacctc	atctcctcta	ccttgacact	ggttcagatt	tcggcagatg	cagtgtgagc	33360
gggtgcttaag	aagtctgaag	gatagctctg	ggggatgggtg	gcacatgaag	gtcacatctt	33420
tttttaatta	ggtgatttaa	atgggtttgt	gcacatgagt	gcaacaccca	tacaggccag	33480
aagggggcat	aagatccccg	gctctagagt	gtcaggctgt	tgtgaactgc	ctggaatagg	33540
tggtgggaac	tgatcttggg	tcccctggaa	gagcattgag	tacatgtcat	cactatctct	33600
ccagcctcac	actcttatcg	gtatgtctca	tttgtggggg	caatttgggg	agtcccacct	33660
ctggatcagg	tactgtgaaa	tatggagtgt	gcggggctgg	atccctgtct	tgccaccagc	33720
cagctgagga	aagctgtcaa	ttgtcttctt	gtgtctgcct	cagtttccta	gaaactagaa	33780
aggaaaaatg	gatggatatca	ctaagttcag	ccttccattg	taaggatcca	ttgaagtagt	33840
tgggtgtgat	tactcacgct	gggtcccttc	cccttctgag	ctgcaagcat	cagctgttgg	33900
accagcaggt	tctgtgtctc	gacaggaagc	agtggtgaaac	tgggctgcaa	gatgatccca	33960
gttctagcat	ttgtgtcaac	cccccttgtc	cacatctgtt	tccacatttt	atttcatcct	34020
tgagcacata	accttttcat	tttgatacat	gcttttcttt	acatagcctg	ggccagcctt	34080
gaactcatgc	tcctcctgcc	ataattctct	aaggactgac	aatgacagtg	gacatgcacc	34140
gccagggtccc	actgtctcta	gtacttttgt	tatgggtctt	ctatgtctgt	ctaattgtgga	34200
atgtgtacat	gcacaccagg	cattgtggga	tgaagttagaa	catgttcatg	cacacaaaga	34260
tcaatcccaa	acagcatctg	cccccccttc	ccctctgccc	cttccccggg	tggatgttag	34320
cattctcttg	tgtagtgtg	ggtcggcccc	ttctctgtct	ctaaacattg	aaacaagggg	34380
aacagaccca	tacataactc	caacacagcg	atctggtcca	agtctggatg	tagaaccctt	34440
ggctctgtct	ttttctcctc	ttcccagagc	tgtcccagtc	gcctcccttt	ctaattggct	34500
gggtgtcatc	taacttgatg	tatatgtttc	ttccttggtc	tgttttatga	ctggcctgtc	34560
ccagtcatta	gtgcatctgg	tgttagaagc	taagctcaac	ttggcctcac	agtctttagt	34620
ttcaggacac	atgggggttat	ggctggatcc	tgtgtcaagc	acatcacttc	tttctgaacc	34680
cacacatctt	aagacatggg	cattgtcaca	tggctgacag	cagtacattc	ttgtatgtag	34740
ttttctctca	agtgttttgg	ttacatggcc	ctaaagccta	ggactgtctg	tcttcaatca	34800
tgtctgccac	ctgtgtccca	gcaagcccaa	gttgtgggtc	ttcctgtcta	atctgcctct	34860
tttatcttta	gcccctctcc	ataagcctct	ttccaactga	gctctggggt	attcagatta	34920
cagctcccta	ctgttttctt	gagaccctaa	gccgtccac	gatgcacagc	agcacttttc	34980
agcagtcccta	tggagatgca	gtgtcgagat	gatatgagct	gtgtgtgata	tgtaaaatgc	35040
atggtagaat	ctgaaaacca	agcacaacaa	aataaaaaaa	aagttcatta	atattttatag	35100
caatcatgta	atgttttgaa	aatacgtgcc	taataaaatt	attttcacct	ttttaaaaat	35160
gtgggtacta	tgatgtttaa	tagcgcatat	tcatgtggca	tatttacctc	aatgggggca	35220
ggaaaacaag	acgacaaagg	gtgggatatt	tctctcaact	ttcctagctg	caacttgtga	35280
taatttgtga	gacacattaa	gcactcaaga	aggggtttgat	tctgtagaca	aaacccgggg	35340
tcagacagtc	cacagcttca	gcatagtgtg	ttcactatat	ctaaagtagg	ttttacttct	35400
ttaagattaa	ttttgttctt	agacagtttc	ccacatgcag	cacaaagcag	ttatgtttta	35460
atttttatagg	ctacaatgac	ggtatttcaa	aaagtaagat	gtgggagtta	caccttgaat	35520
gatcacccaa	ggatgttggg	ccatcagcac	tgaatggccc	agaaccaatc	tacagcaaat	35580
ctgagagagc	agagacatgc	tggtccactg	tgtcaggggc	cttgcccatg	gtgggttgcta	35640
ttgacatctt	gacaggatct	aaaatcacct	aggtgtgtct	tgggcacctc	tggacatata	35700
tgggagtctc	tagactggat	tgcttgaggt	tgatttaaaa	accctaactg	ggcagcacca	35760
ttcattggga	tggggctcctg	gcgtggataa	agattggagg	atgaggggagc	accagggttc	35820
acctctctgc	ctcctgactg	gatgcagcat	gatcagctcc	ctctcctgct	gtgacacaa	35880
ctctgtaccc	tcaaaactgaa	caaaacaagc	tcactcgaaa	gttgctttcg	ttagtccact	35940
tgtcacagct	ttgtcacagt	gatgagaaaa	ataacacaca	ccccccaaat	gaggattgga	36000
tcgaggccct	gggcatgctg	ggtaaactct	tcaccactaa	gctatatctc	taaccctttt	36060
tactttgaat	agggtcttaa	ttgccagat	tggccttaaa	atgataatc	tcctgtcttc	36120
agcttcccaa	gcagctggaa	ttacaaatgc	gagcactagt	atgtatatgt	agtatataaa	36180
taatataattg	tacatatatt	cacatgtgtg	tgtacatgtg	tgtacacaa	tacaaatggg	36240
agccccagaa	tgatgtatac	acatacatat	aagttaggtaa	ttaatatata	catattttaca	36300
catattatac	tagtgtgtat	atatagaata	tataatatgt	attgcacata	tattcacata	36360
tgtgtgtaca	tgtccatgtc	acattgtgat	tgctagttaa	ataccacttt	tcctgtctct	36420
ctaatccaag	tagcaattga	acctgtgaat	tatggttaa	ttcaggggat	tgaaaagcct	36480
gacctccaca	gaatcagcta	acgttagctg	ctactatgat	tccaagcagt	agttactgga	36540
agtgtttgct	cctgtattct	cttgggaagga	agaactagat	tgtagtcat	cagttccag	36600
tcggttatgg	tccccctgtt	ggcctcatag	accagaaagt	agagttgcat	ggtttgattg	36660
taaagtgacc	cacctccac	aggttcacat	gcttgaacac	tgggtcccca	gatgatggag	36720
ctcttggggg	aggttggga	gaccatctag	gaggttagct	tacctggagg	aaatgggctt	36780
ctaggggaca	gacattgaag	atgataggtt	ttccctgttt	ctgccctggg	aatatatcca	36840

```

aattcctcac tcgaatgcc aagccctgag atgagccttc cccttcacag tggactctac 36900
cctcaaaccg tgagccagag tagatccttc gtcctttgct agatgttttg tcacggcagt 36960
gaggtaaagt accgagcaca gcatcattgt tccattttta cggatgggaa aactgagcct 37020
tgaggagatgc ccaaggctgt cagccttgag tgttttaaaag ctgcaaggat tggatgcatt 37080
tgtcctatat cagggaaaca tgatggggat ggagctctggg tgctgaggca ctgagttggc 37140
aagagggaag gcctggtgtt cactcaaagt cattaaggac caattgtgtc tgcaatcctg 37200
tgccctccct tgaacaatag gtaagggtca gtgtgagccc tgattactcc cagcagaagg 37260
atacatctgc tttggagacc aaagtccctt cactgtagaa actaggtcct tcaaggtctc 37320
agataaaaag acaatgggtt ttgtgcta atccacccaa tgggtgtgtg ggtcacacta 37380
ctcactgggc caacatggtg gtctggacag tcacacagga tgaagtgagg aaaggcaagt 37440
ctccccggca cctcccttat cgtcactgca aggccaccct gactaagagt cccctcttca 37500
atgctggccc tacgaatata ttatcactct tcgtgtttac aagatttctc tccttgggaag 37560
gtgtgatgtg gacagtgaag ctctcaacaa cccctactca cctaagacct agacaagaga 37620
gtctgggatg gccatgtatg tactccttca gtaattgaca gccattttct ttgtctagga 37680
agtcttctca caagcttccg caccataacg gtatcggctg tcctgattta cagacatgtc 37740
attgggactg atatctgcaa caaagagcag atctccagtc actcatctca ctccagcttg 37800
ttcacagaac aaaaaggagt tgaaggaggg tcttcatcat tgggtgttcc cttccatgga 37860
gcataggaca ggcatggtgc ccagaacctg cccagcttct agctctccaa gcctcatgct 37920
ttcctgtctc nahaaaaaaa aaaaaaaaaa 37950

```

&lt;210&gt; 184

&lt;211&gt; 1381

&lt;212&gt; DNA

&lt;213&gt; Mus musculus

&lt;400&gt; 184

```

gagccgagag gatgctgctg tccctggtgc tccacacgta ctct atg cgc tac ctg      56
                                     Met Arg Tyr Leu
                                     1
ctc ccc agc gtc ctg ttg ctg ggc tgc gcg ccc acc tac ctg ctg gcc      104
Leu Pro Ser Val Leu Leu Gly Ser Ala Pro Thr Tyr Leu Leu Ala
5                                     10                                     20
tgg acg ctg tgg cgg gtg ctc tcc gcg ctg atg ccc gcc cgc ctg tac      152
Trp Thr Leu Trp Arg Val Leu Ser Ala Leu Met Pro Ala Arg Leu Tyr
25                                     30                                     35
cag cgc gtg gac gac cgg ctt tac tgc gtc tac cag aac atg gtg ctc      200
Gln Arg Val Asp Asp Arg Leu Tyr Cys Val Tyr Gln Asn Met Val Leu
40                                     45                                     50
ttc ttc ttc gag aac tac acc ggg gtc cag ata ttg cta tat gga gat      248
Phe Phe Phe Glu Asn Tyr Thr Gly Val Gln Ile Leu Leu Tyr Gly Asp
55                                     60                                     65
ttg cca aaa aat aaa gaa aat gta ata tat cta gcg aat cat caa agc      296
Leu Pro Lys Asn Lys Glu Asn Val Ile Tyr Leu Ala Asn His Gln Ser
70                                     75                                     80
aca gtt gac tgg att gtt gcg gac atg ctg gct gcc aga cag gat gcc      344
Thr Val Asp Trp Ile Val Ala Asp Met Leu Ala Ala Arg Gln Asp Ala
85                                     90                                     95                                     100
cta gga cat gtg cgc tac gta ctg aaa gac aag tta aaa tgg ctt ccg      392
Leu Gly His Val Arg Tyr Val Leu Lys Asp Lys Leu Lys Trp Leu Pro
105                                     110                                     115
ctg tat ggg ttc tac ttt gct cag cat gga gga att tat gta aaa cga      440
Leu Tyr Gly Phe Tyr Phe Ala Gln His Gly Gly Ile Tyr Val Lys Arg
120                                     125                                     130
agt gcc aaa ttt aat gat aaa gaa atg aga agc aag ctg cag agc tat      488
Ser Ala Lys Phe Asn Asp Lys Glu Met Arg Ser Lys Leu Gln Ser Tyr
135                                     140                                     145
gtg aac gca gga aca ccg atg tat ctt gtg att ttc cca gag gga aca      536
Val Asn Ala Gly Thr Pro Met Tyr Leu Val Ile Phe Pro Glu Gly Thr
150                                     155                                     160
agg tat aat gca aca tac aca aaa ctc ctt tca gcc agt cag gca ttt      584
Arg Tyr Asn Ala Thr Tyr Thr Lys Leu Leu Ser Ala Ser Gln Ala Phe
165                                     170                                     175                                     180
gct gct cag cgg ggc ctt gca gta tta aaa cac gta ctg aca cca aga      632

```

Ala Ala Gln Arg Gly Leu Ala Val Leu Lys His Val Leu Thr Pro Arg  
185 190 195  
ata aag gcc act cac gtt gct ttt gat tct atg aag agt cat tta gat 680  
Ile Lys Ala Thr His Val Ala Phe Asp Ser Met Lys Ser His Leu Asp  
200 205 210  
gca att tat gat gtc aca gtg gtt tat gaa ggg aat gag aaa ggt tca 728  
Ala Ile Tyr Asp Val Thr Val Tyr Glu Gly Asn Glu Lys Gly Ser  
215 220 225  
gga aaa tac tca aat cca cca tcc atg act gag ttt ctc tgc aaa cag 776  
Gly Lys Tyr Ser Asn Pro Pro Ser Met Thr Glu Phe Leu Cys Lys Gln  
230 235 240  
tgc cca aaa ctt cat att cac ttt gat cgt ata gac aga aat gaa gtt 824  
Cys Pro Lys Leu His Ile His Phe Asp Arg Ile Asp Arg Asn Glu Val  
245 250 255 260  
cca gag gaa caa gaa cac atg aaa aag tgg ctt cat gag cgc ttt gag 872  
Pro Glu Glu Gln Glu His Met Lys Lys Trp Leu His Glu Arg Phe Glu  
265 270 275  
ata aaa gat agg ttg ctc ata gag ttc tat gat tca cca gat cca gaa 920  
Ile Lys Asp Arg Leu Leu Ile Glu Phe Tyr Asp Ser Pro Asp Pro Glu  
280 285 290  
aga aga aac aaa ttt cct ggg aaa agt gtt cat tcc aga cta agt gtg 968  
Arg Arg Asn Lys Phe Pro Gly Lys Ser Val His Ser Arg Leu Ser Val  
295 300 305  
aag aag act tta cct tca gtg ttg atc ttg ggg agt ttg act gcg gtc 1016  
Lys Lys Thr Leu Pro Ser Val Leu Ile Leu Gly Ser Leu Thr Ala Val  
310 315 320  
atg ctg atg acg gag tcc gga agg aaa ctg tac atg ggc acc tgg ttg 1064  
Met Leu Met Thr Glu Ser Gly Arg Lys Leu Tyr Met Gly Thr Trp Leu  
325 330 335 340  
tat gga acc ctc ctt ggc tgc ctg tgg ttt gtt att aaa gca taa 1109  
Tyr Gly Thr Leu Leu Gly Cys Leu Trp Phe Val Ile Lys Ala \*  
345 350 355  
gcaagtagca ggctgcagtc acagtctctt attgatggct acacattgta tcacattggt 1169  
tcctgaatta aataaggagt tttcttggtt ttgttttttt tgttttggtt tgttctggtt 1229  
taagccttga tgattgaaca ctggataaag tcgagtcctg tgaccacagc caacatgcat 1289  
ttgatttggg gcaaacacat gtggcttttc aggtgctggg gttgctggag acatggaagc 1349  
taagtggagt ttatgctggt tttttttttt tt 1381  
<210> 185  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-14-107  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-14-107.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-14-107.mis2  
<400> 185  
ctaaacaacc accaaatgca tacagcaacc aggcaaatgc ctgatag 47  
<210> 186  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>

<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-14-317  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-14-317.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-14-317.mis2  
<400> 186  
cataacatgc aaggtgggca agaaaaagag gtgggcacag ctcatga  
<210> 187  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-14-35  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-14-35.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-14-35.mis2  
<400> 187  
atccaacaca gaaaccgcta aaaccaggca gaagctgtct gcagaga  
<210> 188  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-20-149  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-20-149.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-20-149.mis2  
<400> 188  
tttttgctgt gtcttcaaag tgactcttgg tttattgcct gctaagg  
<210> 189  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-20-77  
<221> allele

47

47

47

<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-20-77.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-20-77.mis2  
<400> 189  
tgcaacatga agattctgaa gggactttgt tgtctgagaa cacatct  
<210> 190  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-22-174  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-22-174.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-22-174.mis2  
<400> 190  
ggattgtgca gaagttgcct ttcattgttca aaaatgttaa tttgttt  
<210> 191  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-22-176  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-22-176.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-22-176.mis2  
<400> 191  
attgtgcaga agttgccttt catattcaaa aatgttaatt tgtttgt  
<210> 192  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-26-60  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23

47

47

47

<223> potential microsequencing oligo 4-26-60.mis1  
 <221> primer\_bind  
 <222> 25..47  
 <223> complement potential microsequencing oligo 4-26-60.mis2  
 <400> 192  
 gatgggaaag tgcattcttaa gacagtttagc aggccaagga gcgactt 47  
 <210> 193  
 <211> 47  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> allele  
 <222> 1..47  
 <223> polymorphic fragment 4-26-72  
 <221> allele  
 <222> 24  
 <223> polymorphic base A  
 <221> primer\_bind  
 <222> 1..23  
 <223> potential microsequencing oligo 4-26-72.mis1  
 <221> primer\_bind  
 <222> 25..47  
 <223> complement potential microsequencing oligo 4-26-72.mis2  
 <400> 193  
 catcttaaga cagtttagcag gccaaaggagc gactttaag ggtgagc 47  
 <210> 194  
 <211> 47  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> allele  
 <222> 1..47  
 <223> polymorphic fragment 4-3-130  
 <221> allele  
 <222> 24  
 <223> polymorphic base A  
 <221> primer\_bind  
 <222> 1..23  
 <223> potential microsequencing oligo 4-3-130.mis1  
 <221> primer\_bind  
 <222> 25..47  
 <223> complement potential microsequencing oligo 4-3-130.mis2  
 <400> 194  
 tattgggcct aaaacagtat tctataaagc ttaaattggt attact 47  
 <210> 195  
 <211> 47  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> allele  
 <222> 1..47  
 <223> polymorphic fragment 4-38-63  
 <221> allele  
 <222> 24  
 <223> polymorphic base A  
 <221> primer\_bind  
 <222> 1..23  
 <223> potential microsequencing oligo 4-38-63.mis1  
 <221> primer\_bind  
 <222> 25..47  
 <223> complement potential microsequencing oligo 4-38-63.mis2



<400> 195  
tataagttat aagaaaatca ggcagaggct aaactttttt tttgttt 47  
<210> 196  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-38-83  
<221> allele  
<222> 24  
<223> polymorphic base G  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-38-83.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-38-83.mis2  
<400> 196  
ggcagaggct aaactttttt tttgtttggc aatgctgttg agaatat 47  
<210> 197  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-4-152  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-4-152.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-4-152.mis2  
<400> 197  
tactttccca ttgttcctga cttcgttatc ctatatataa acagaaa 47  
<210> 198  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-4-187  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-4-187.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-4-187.mis2  
<400> 198  
tataaacaga aacatggatg agtaaaaaaa aaaaaaaaaa aaaaaaa 47  
<210> 199  
<211> 47

<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-4-288  
<221> allele  
<222> 24  
<223> polymorphic base G  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-4-288.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-4-288.mis2  
<400> 199

ctgtcatcaa ctaattttca caagtaccta tgttttgatt tcatgta

47

<210> 200  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-42-304  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-42-304.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-42-304.mis2  
<400> 200

attatttaaa actattttatg taaccttatt ttcagggggtt ttaatt

47

<210> 201  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-42-401  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-42-401.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-42-401.mis2  
<400> 201

taagaaagaa ttctgtgttc tggacaaagt ttaaaccac agagcca

47

<210> 202  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele

<222> 1..47  
<223> polymorphic fragment 4-43-328  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-43-328.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-43-328.mis2  
<400> 202  
agaattctgt gttctggcca aagcttaaac ccacagagcc agtttaa  
<210> 203  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-43-70  
<221> allele  
<222> 24  
<223> polymorphic base G  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-43-70.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-43-70.mis2  
<400> 203  
atcgctcca ttattctcaa aaagaccatg ggacacaaca caagaag  
<210> 204  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-50-209  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-50-209.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-50-209.mis2  
<400> 204  
atatagagtg tgcattccctg aactgaaac tgaaggcttt atggttt  
<210> 205  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-50-293  
<221> allele  
<222> 24

47

47

47

```

<223> polymorphic base G
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-50-293.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-50-293.mis2
<400> 205
cctgagtcctc agggggctga caggggacag tttaaacat tgatgaa      47
<210> 206
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-50-323
<221> allele
<222> 24
<223> polymorphic base C
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-50-323.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-50-323.mis2
<400> 206
tttaaacat tgatgaatct ttactactac aaaagggttc gatttag      47
<210> 207
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-50-329
<221> allele
<222> 24
<223> polymorphic base C
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-50-329.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-50-329.mis2
<400> 207
acattgatga atctttatta ctacaaaagg gttcgattta ggctagc      47
<210> 208
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-50-330
<221> allele
<222> 24
<223> polymorphic base A
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-50-330.mis1

```

<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-50-330.mis2  
<400> 208  
cattgatgaa tctttattac tacaaaaggg ttcgatttag gctagcc  
<210> 209  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-52-163  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-52-163.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-52-163.mis2  
<400> 209  
gaacaggata ttcttaacta ccaaagaatt ttacacatct attgttt  
<210> 210  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-52-88  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-52-88.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-52-88.mis2  
<400> 210  
tccatgtcat tattattcaa aagcttaaaa aatacacaag gtgaaaa  
<210> 211  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-53-258  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-53-258.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-53-258.mis2  
<400> 211

47

47

47

gagaaatcat gcagagagaa tgcatcttca ctcaaatttt aacctaa 47  
<210> 212  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-54-283  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-54-283.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-54-283.mis2  
<400> 212  
aagtagtttt tcacactttc tctatgatac aatcgatggc ttaatct 47  
<210> 213  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-54-388  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-54-388.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-54-388.mis2  
<400> 213  
ctctctatcg tatacatctt tacacacgct gcagcgccaa gactcca 47  
<210> 214  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-55-70  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-55-70.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-55-70.mis2  
<400> 214  
tattaagaac ctaggtttta aaaaactctc tategtatac atcttta 47  
<210> 215  
<211> 47  
<212> DNA

```

<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-55-95
<221> allele
<222> 24
<223> polymorphic base A
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-55-95.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-55-95.mis2
<400> 215
ctctctatcg tatacatctt tacacacgct gcagcgccaa gactcca
<210> 216
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-56-159
<221> allele
<222> 24
<223> polymorphic base C
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-56-159.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-56-159.mis2
<400> 216
aagttttcct tctcttctgt agacgtctcc atgttacagt caactat
<210> 217
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-56-213
<221> allele
<222> 24
<223> polymorphic base A
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-56-213.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-56-213.mis2
<400> 217
atggctcatg ttcactctgg ttcaccttca gaggagtttg atatttt
<210> 218
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47

```

<223> polymorphic fragment 4-58-289  
 <221> allele  
 <222> 24  
 <223> polymorphic base G  
 <221> primer\_bind  
 <222> 1..23  
 <223> potential microsequencing oligo 4-58-289.mis1  
 <221> primer\_bind  
 <222> 25..47  
 <223> complement potential microsequencing oligo 4-58-289.mis2  
 <400> 218  
 cataacctgca gcctgctttt ggtgaggggt gactacttta cctgcaa 47  
 <210> 219  
 <211> 47  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> allele  
 <222> 1..47  
 <223> polymorphic fragment 4-58-318  
 <221> allele  
 <222> 24  
 <223> polymorphic base A  
 <221> primer\_bind  
 <222> 1..23  
 <223> potential microsequencing oligo 4-58-318.mis1  
 <221> primer\_bind  
 <222> 25..47  
 <223> complement potential microsequencing oligo 4-58-318.mis2  
 <400> 219  
 tgactacttt acctgcaata tttatttgca agtttatttc ttccttt 47  
 <210> 220  
 <211> 47  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> allele  
 <222> 1..47  
 <223> polymorphic fragment 4-60-266  
 <221> allele  
 <222> 24  
 <223> polymorphic base G  
 <221> primer\_bind  
 <222> 1..23  
 <223> potential microsequencing oligo 4-60-266.mis1  
 <221> primer\_bind  
 <222> 25..47  
 <223> complement potential microsequencing oligo 4-60-266.mis2  
 <400> 220  
 aacaggacca agacactgca ttagataaag tttcagtatt tcttagc 47  
 <210> 221  
 <211> 47  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> allele  
 <222> 1..47  
 <223> polymorphic fragment 4-60-293  
 <221> allele  
 <222> 24  
 <223> polymorphic base C



<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-60-293.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-60-293.mis2  
<400> 221  
aagtttcagt atttcttagc agacgaagcc agcaggaagt cctccta 47  
<210> 222  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-84-241  
<221> allele  
<222> 24  
<223> polymorphic base G  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-84-241.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-84-241.mis2  
<400> 222  
gaaaaaaaa tagtgactgc cacggtgaat aattcagttc ttcagaa 47  
<210> 223  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-84-262  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-84-262.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-84-262.mis2  
<400> 223  
acggtgaata attcagttct tcaaaagcag caacatgata tcatgga 47  
<210> 224  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-86-206  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-86-206.mis1  
<221> primer\_bind

<222> 25..47  
<223> complement potential microsequencing oligo 4-86-206.mis2  
<400> 224  
gtattcaaat caggacacac cacaatggc atctacacgt taacatt 47  
<210> 225  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-86-309  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-86-309.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-86-309.mis2  
<400> 225  
tggctctagg caggccactt tagagagtga ggaaccagag agcagaa 47  
<210> 226  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-88-349  
<221> allele  
<222> 24  
<223> polymorphic base G  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-88-349.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-88-349.mis2  
<400> 226  
gaaactaaaa gacaatattc agtgtgagat tttccaagtt ctttatg 47  
<210> 227  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-89-87  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-89-87.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-89-87.mis2  
<400> 227  
ttcttccctg aacgctgggt tcacatagtt tttgtgtga gaataga 47

<210> 228  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-123-184  
<221> allele  
<222> 24  
<223> polymorphic base G  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-123-184.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-123-184.mis2  
<400> 228  
ccagcccaga acattcacca gctgggccaagagttctgct gggtttt 47  
<210> 229  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-128-202  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-128-202.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-128-202.mis2  
<400> 229  
aatgtctgtt tcttagagaa ctgaaacaca cacacataca tacacac 47  
<210> 230  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-128-275  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-128-275.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-128-275.mis2  
<400> 230  
acaccctac ctcacatgtg tagacaaatg tatgcatata tgtctct 47  
<210> 231  
<211> 47  
<212> DNA  
<213> Homo Sapiens

<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-128-313  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-128-313.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-128-313.mis2  
<400> 231  
tatgtctcta gacagatata cataagattc tatttggcat agaaaaa 47  
<210> 232  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-128-60  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-128-60.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-128-60.mis2  
<400> 232  
gcactgtgac ccaggcgcta ggtccctctt acagtgcac tccgaca 47  
<210> 233  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-12907-295  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-12907-295.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-12907-295.mis2  
<400> 233  
gctatatggc attatatctc cacagggcag acctgatgta caagatg 47  
<210> 234  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-130-58

<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-130-58.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-130-58.mis2  
<400> 234  
aaagcaaaag agcttcaaaa atacttcagg agtgtgcata tggcgag 47  
<210> 235  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-134-362  
<221> allele  
<222> 24  
<223> polymorphic base G  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-134-362.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-134-362.mis2  
<400> 235  
caaaacactc atgttagtta gatgattatt cctattacaa agataag 47  
<210> 236  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-140-130  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-140-130.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-140-130.mis2  
<400> 236  
tgttcaaaag cagctacaga ccacatgtaa acaattgagc atggctg 47  
<210> 237  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1462-238  
<221> allele  
<222> 24  
<223> polymorphic base G  
<221> primer\_bind

<222> 1..23  
 <223> potential microsequencing oligo 99-1462-238.mis1  
 <221> primer\_bind  
 <222> 25..47  
 <223> complement potential microsequencing oligo 99-1462-238.mis2  
 <400> 237  
 ccctttcaag gtttagtaact catgtgctgt gtttctgctt cagaagg 47  
 <210> 238  
 <211> 47  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> allele  
 <222> 1..47  
 <223> polymorphic fragment 99-147-181  
 <221> allele  
 <222> 24  
 <223> polymorphic base A  
 <221> primer\_bind  
 <222> 1..23  
 <223> potential microsequencing oligo 99-147-181.mis1  
 <221> primer\_bind  
 <222> 25..47  
 <223> complement potential microsequencing oligo 99-147-181.mis2  
 <400> 238  
 gtgtcatgaa aaagagcatg ataaaaagaa aaacttaaat ctttata 47  
 <210> 239  
 <211> 47  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> allele  
 <222> 1..47  
 <223> polymorphic fragment 99-1474-156  
 <221> allele  
 <222> 24  
 <223> polymorphic base G  
 <221> primer\_bind  
 <222> 1..23  
 <223> potential microsequencing oligo 99-1474-156.mis1  
 <221> primer\_bind  
 <222> 25..47  
 <223> complement potential microsequencing oligo 99-1474-156.mis2  
 <400> 239  
 cttgtactca taagttaaatt attgataaca agaagaaata tggactt 47  
 <210> 240  
 <211> 47  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> allele  
 <222> 1..47  
 <223> polymorphic fragment 99-1474-359  
 <221> allele  
 <222> 24  
 <223> polymorphic base A  
 <221> primer\_bind  
 <222> 1..23  
 <223> potential microsequencing oligo 99-1474-359.mis1  
 <221> primer\_bind  
 <222> 25..47

<223> complement potential microsequencing oligo 99-1474-359.mis2  
<400> 240  
aaaaaaaatc aaattattgt accaaattcc ctaatatcag atgtgta 47  
<210> 241  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1479-158  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-1479-158.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-1479-158.mis2  
<400> 241  
tttaaaaatc cacttgtaat cgccgctaatt tggagtgtat attcagg 47  
<210> 242  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1479-379  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-1479-379.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-1479-379.mis2  
<400> 242  
gtagagctgt gtactgaggt cagagaagca gctcatggta cagcctt 47  
<210> 243  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-148-129  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-148-129.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-148-129.mis2  
<400> 243  
ttcatatcta tacaataat tttaaattta atacataggg ctgcaaa 47  
<210> 244

<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-148-132  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-148-132.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-148-132.mis2  
<400> 244  
atatctatac aaataatttt gaacttaata catagggctg caaaaca 47  
<210> 245  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-148-139  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-148-139.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-148-139.mis2  
<400> 245  
tacaaataat tttgaattta atacataggg ctgcaaaaca aggttga 47  
<210> 246  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-148-140  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-148-140.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-148-140.mis2  
<400> 246  
acaaataatt ttgaatttaa tacatagggc tgcaaaacaa ggttgat 47  
<210> 247  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>



<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-148-182  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-148-182.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-148-182.mis2  
<400> 247  
ttgatgttga tatgggcaac tgtatgttgg atggtcccaa agcattc 47  
<210> 248  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-148-366  
<221> allele  
<222> 24  
<223> polymorphic base G  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-148-366.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-148-366.mis2  
<400> 248  
tccttgtaa aggtctctcc ctggtgctca cggctgccgc ctcaaag 47  
<210> 249  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-148-76  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-148-76.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-148-76.mis2  
<400> 249  
tgatagaatg ccttcctgaa ttactactct tgatggcttc ataaaac 47  
<210> 250  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1480-290  
<221> allele

<222> 24  
<223> polymorphic base G  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-1480-290.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-1480-290.mis2  
<400> 250  
tgcaccatct tcaccacaac cccgggcaac cactgatcct tttactg 47  
<210> 251  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1481-285  
<221> allele  
<222> 24  
<223> polymorphic base G  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-1481-285.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-1481-285.mis2  
<400> 251  
tcccataacc tggttttgctt ctcgctctaa cctcaagatg gtataaa 47  
<210> 252  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1484-101  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-1484-101.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-1484-101.mis2  
<400> 252  
aaaaagatca aatataagca tgtaactcct ctccttaaaa tctcagt 47  
<210> 253  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1484-328  
<221> allele  
<222> 24  
<223> polymorphic base G  
<221> primer\_bind  
<222> 1..23

<223> potential microsequencing oligo 99-1484-328.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-1484-328.mis2  
<400> 253  
ggacacgtgg tcatgaggag tttgaaggga ttcagttttc agatccc 47  
<210> 254  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1485-251  
<221> allele  
<222> 24  
<223> polymorphic base G  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-1485-251.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-1485-251.mis2  
<400> 254  
gattgccttg atatatgctc ccagagaacc aagaatgtcc ccttttc 47  
<210> 255  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1490-381  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-1490-381.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-1490-381.mis2  
<400> 255  
tgcacagtgg aaataccatg tcacggtacg ctactgtgca tctcttc 47  
<210> 256  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1493-280  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-1493-280.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-1493-280.mis2

<400> 256  
ggatgacaga gtattgttgg aggaatgggg ttggtctgct tgttttt 47  
<210> 257  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-151-94  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-151-94.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-151-94.mis2  
<400> 257  
attgagatca ttgataagga aatattctaa aatttcaaaa tctatat 47  
<210> 258  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-211-291  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-211-291.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-211-291.mis2  
<400> 258  
ctggttatat cagactgacc ttcattgttt caacagggtca atgcctt 47  
<210> 259  
<211> 45  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..45  
<223> polymorphic fragment 99-213-37  
<221> allele  
<222> 23  
<223> polymorphic base T  
<221> primer\_bind  
<222> 1..22  
<223> potential microsequencing oligo 99-213-37.mis1  
<221> primer\_bind  
<222> 24..45  
<223> complement potential microsequencing oligo 99-213-37.mis2  
<400> 259  
gtgcttccgg ctgcaggact gttggaggac tccagtgtct gacag 45  
<210> 260  
<211> 47

<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-221-442  
<221> allele  
<222> 24  
<223> polymorphic base A  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-221-442.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-221-442.mis2  
<400> 260  
tgcctttgta gatatgcatg ggaattccat gacctagcca gacgaat 47  
<210> 261  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-222-109  
<221> allele  
<222> 24  
<223> polymorphic base C  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-222-109.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-222-109.mis2  
<400> 261  
caggtgagga gtgctggatt ggccacgata tgaatttctt cagcagt 47  
<210> 262  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-14-107, variant version of SEQ ID185  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID185  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-14-107.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-14-107.mis2  
<400> 262  
ctaaacaacc accaaatgca tacggcaacc aggcaaatgc ctgatag 47  
<210> 263  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele

```

<222> 1..47
<223> polymorphic fragment 4-14-317, variant version of SEQ ID186
<221> allele
<222> 24
<223> base G ; A in SEQ ID186
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-14-317.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-14-317.mis2
<400> 263
cataacatgc aagggtgggca agagaaagag gtgggcacag ctcatga
<210> 264
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-14-35, variant version of SEQ ID187
<221> allele
<222> 24
<223> base T ; C in SEQ ID187
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-14-35.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-14-35.mis2
<400> 264
atccaacaca gaaaccgcta aaatcaggca gaagctgtct gcagaga
<210> 265
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-20-149, variant version of SEQ ID188
<221> allele
<222> 24
<223> base T ; C in SEQ ID188
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-20-149.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-20-149.mis2
<400> 265
tttttgctgt gtcttcaaag tgattcttgg tttattgcct gctaagg
<210> 266
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-20-77, variant version of SEQ ID189
<221> allele
<222> 24

```

<223> base T ; A in SEQ ID189  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-20-77.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-20-77.mis2  
<400> 266  
tgcaacatga agattctgaa gggctctttgt tgtctgagaa cacatct 47  
<210> 267  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-22-174, variant version of SEQ ID190  
<221> allele  
<222> 24  
<223> base C ; A in SEQ ID190  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-22-174.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-22-174.mis2  
<400> 267  
ggattgtgca gaagttgcct ttcctgttca aaaatgttaa tttgttt 47  
<210> 268  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-22-176, variant version of SEQ ID191  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID191  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-22-176.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-22-176.mis2  
<400> 268  
attgtgcaga agttgccttt catgttcaaa aatgttaatt tgtttgt 47  
<210> 269  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-26-60, variant version of SEQ ID192  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID192  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-26-60.mis1

<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-26-60.mis2  
<400> 269  
gatgggaaag tgcattcttaa gacgggtagc aggccaagga gcgactt 47  
<210> 270  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-26-72, variant version of SEQ ID193  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID193  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-26-72.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-26-72.mis2  
<400> 270  
catcttaaga cagtttagcag gccgaggagc gactttaag ggtgagc 47  
<210> 271  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-3-130, variant version of SEQ ID194  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID194  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-3-130.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-3-130.mis2  
<400> 271  
tattgggcct aaaacagtat tctgtaaagc ttaaattggt attaact 47  
<210> 272  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-38-63, variant version of SEQ ID195  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID195  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-38-63.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-38-63.mis2  
<400> 272



tataagttat aagaaaatca ggcggaggct aaactttttt tttgttt 47  
<210> 273  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-38-83, variant version of SEQ ID196  
<221> allele  
<222> 24  
<223> base T ; G in SEQ ID196  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-38-83.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-38-83.mis2  
<400> 273  
ggcagaggct aaactttttt ttttttggc aatgctgttg agaatat 47  
<210> 274  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-4-152, variant version of SEQ ID197  
<221> allele  
<222> 24  
<223> base T ; C in SEQ ID197  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-4-152.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-4-152.mis2  
<400> 274  
tactttccca ttgttcctga cttgtttatc ctatatataa acagaaa 47  
<210> 275  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-4-187, variant version of SEQ ID198  
<221> allele  
<222> 24  
<223> base T ; A in SEQ ID198  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-4-187.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-4-187.mis2  
<400> 275  
tataaacaga aacatggatg agttaaaaaa aaaaaaaaaa aaaaaaa 47  
<210> 276  
<211> 47  
<212> DNA

<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-4-288, variant version of SEQ ID199  
<221> allele  
<222> 24  
<223> base C ; G in SEQ ID199  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-4-288.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-4-288.mis2  
<400> 276  
ctgtcatcaa ctaattttca caactaccta tgttttgatt tcatgta 47  
<210> 277  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-42-304, variant version of SEQ ID200  
<221> allele  
<222> 24  
<223> base T ; C in SEQ ID200  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-42-304.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-42-304.mis2  
<400> 277  
attatttaaa actatttatg taatcttatt ttcaggggtt ttttaatt 47  
<210> 278  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-42-401, variant version of SEQ ID201  
<221> allele  
<222> 24  
<223> base C ; A in SEQ ID201  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-42-401.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-42-401.mis2  
<400> 278  
taagaaagaa ttctgtgttc tggccaaagt ttaaaccac agagcca 47  
<210> 279  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47

<223> polymorphic fragment 4-43-328, variant version of SEQ ID202  
<221> allele  
<222> 24  
<223> base T ; C in SEQ ID202  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-43-328.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-43-328.mis2  
<400> 279  
agaattctgt gttctggcca aagtttaaac ccacagagcc agttaa  
47  
<210> 280  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-43-70, variant version of SEQ ID203  
<221> allele  
<222> 24  
<223> base C ; G in SEQ ID203  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-43-70.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-43-70.mis2  
<400> 280  
atcgctcca ttattctcaa aaacaccatg ggacacaaca caagaag  
47  
<210> 281  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-50-209, variant version of SEQ ID204  
<221> allele  
<222> 24  
<223> base T ; C in SEQ ID204  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-50-209.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-50-209.mis2  
<400> 281  
atatagagtg tgcattcctg acattgaaac tgaaggcttt atggttt  
47  
<210> 282  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-50-293, variant version of SEQ ID205  
<221> allele  
<222> 24  
<223> base T ; G in SEQ ID205

<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-50-293.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-50-293.mis2  
<400> 282  
cctgagtcctc agggggctga cagtggacag tttaaacat tgatgaa  
<210> 283  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-50-323, variant version of SEQ ID206  
<221> allele  
<222> 24  
<223> base T ; C in SEQ ID206  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-50-323.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-50-323.mis2  
<400> 283  
tttaaacat tgatgaatct ttattactac aaaaggggttc gatttag  
<210> 284  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-50-329, variant version of SEQ ID207  
<221> allele  
<222> 24  
<223> base T ; C in SEQ ID207  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-50-329.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-50-329.mis2  
<400> 284  
acattgatga atctttatta ctataaaagg gttcgattta ggctagc  
<210> 285  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-50-330, variant version of SEQ ID208  
<221> allele  
<222> 24  
<223> base T ; A in SEQ ID208  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-50-330.mis1  
<221> primer\_bind

47

47

47

```

<222> 25..47
<223> complement potential microsequencing oligo 4-50-330.mis2
<400> 285
cattgatgaa tctttattac tactaaaggg ttcgatttag gctagcc
<210> 286
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-52-163, variant version of SEQ ID209
<221> allele
<222> 24
<223> base C ; A in SEQ ID209
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-52-163.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-52-163.mis2
<400> 286
gaacaggata ttcttaacta ccacagaatt ttacacatct attgttt
<210> 287
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-52-88, variant version of SEQ ID210
<221> allele
<222> 24
<223> base T ; C in SEQ ID210
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-52-88.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-52-88.mis2
<400> 287
tccatgtcat tattattcaa aagtttaaaa aatacacaag gtgaaaa
<210> 288
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 4-53-258, variant version of SEQ ID211
<221> allele
<222> 24
<223> base G ; A in SEQ ID211
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 4-53-258.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 4-53-258.mis2
<400> 288
gagaaatcat gcagagagaa tgcgttctca ctcaaatttt aacctaa

```

<210> 289  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-54-283, variant version of SEQ ID212  
<221> allele  
<222> 24  
<223> base T ; A in SEQ ID212  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-54-283.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-54-283.mis2  
<400> 289  
aagtagtttt tcacactttc tctttgatac aatcgatggc ttaatct  
<210> 290  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-54-388, variant version of SEQ ID213  
<221> allele  
<222> 24  
<223> base C ; A in SEQ ID213  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-54-388.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-54-388.mis2  
<400> 290  
ctctctatcg tatacatctt tacccacgct gcagcgccaa gactcca  
<210> 291  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-55-70, variant version of SEQ ID214  
<221> allele  
<222> 24  
<223> base T ; A in SEQ ID214  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-55-70.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-55-70.mis2  
<400> 291  
tattaagaac ctaggtttta aaatactctc tatcgatac atcttta  
<210> 292  
<211> 47  
<212> DNA  
<213> Homo Sapiens

47

47

47

<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-55-95, variant version of SEQ ID215  
<221> allele  
<222> 24  
<223> base C ; A in SEQ ID215  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-55-95.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-55-95.mis2  
<400> 292  
ctctctatcg tatacatctt tacccacgct gcagcgccaa gactcca 47  
<210> 293  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-56-159, variant version of SEQ ID216  
<221> allele  
<222> 24  
<223> base T ; C in SEQ ID216  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-56-159.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-56-159.mis2  
<400> 293  
aagttttcct tctcttctgt agatgtctcc atgttacagt caactat 47  
<210> 294  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-56-213, variant version of SEQ ID217  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID217  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-56-213.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-56-213.mis2  
<400> 294  
atggctcatg ttcactctgg ttgccttca gaggagtttg atatttt 47  
<210> 295  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-58-289, variant version of SEQ ID218

<221> allele  
<222> 24  
<223> base C ; G in SEQ ID218  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-58-289.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-58-289.mis2  
<400> 295  
catacctgca gcctgctttt ggtcaggggt gactacttta cctgcaa 47  
<210> 296  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-58-318, variant version of SEQ ID219  
<221> allele  
<222> 24  
<223> base C ; A in SEQ ID219  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-58-318.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-58-318.mis2  
<400> 296  
tgactacttt acctgcaata tttctttgca agtttatttc ttccttt 47  
<210> 297  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-60-266, variant version of SEQ ID220  
<221> allele  
<222> 24  
<223> base T ; G in SEQ ID220  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-60-266.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-60-266.mis2  
<400> 297  
aacaggacca agacactgca ttatataaag tttcagtatt tcttagc 47  
<210> 298  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-60-293, variant version of SEQ ID221  
<221> allele  
<222> 24  
<223> base T ; C in SEQ ID221  
<221> primer\_bind



<222> 1..23  
<223> potential microsequencing oligo 4-60-293.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-60-293.mis2  
<400> 298  
aagtttcagt atttcttagc agatgaagcc agcaggaagt cctccta  
<210> 299  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-84-241, variant version of SEQ ID222  
<221> allele  
<222> 24  
<223> base T ; G in SEQ ID222  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-84-241.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-84-241.mis2  
<400> 299  
gaaaaaaaa tagtgactgc cactgtgaat aattcagttc ttcagaa  
<210> 300  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-84-262, variant version of SEQ ID223  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID223  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-84-262.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-84-262.mis2  
<400> 300  
acggtgaata attcagttct tcagaagcag caacatgatc tcatgga  
<210> 301  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-86-206, variant version of SEQ ID224  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID224  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-86-206.mis1  
<221> primer\_bind  
<222> 25..47

47

47

47

<223> complement potential microsequencing oligo 4-86-206.mis2  
<400> 301  
gtattcaaat caggacacac cacgaatggc atctacacgt taacatt 47  
<210> 302  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-86-309, variant version of SEQ ID225  
<221> allele  
<222> 24  
<223> base T ; A in SEQ ID225  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-86-309.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-86-309.mis2  
<400> 302  
tggctctagg caggccactt tagtgagtga ggaaccagag agcagaa 47  
<210> 303  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-88-349, variant version of SEQ ID226  
<221> allele  
<222> 24  
<223> base C ; G in SEQ ID226  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-88-349.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-88-349.mis2  
<400> 303  
gaaactaaaa gacaatattc agtctgagat tttccaagtt ctttatg 47  
<210> 304  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 4-89-87, variant version of SEQ ID227  
<221> allele  
<222> 24  
<223> base T ; C in SEQ ID227  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 4-89-87.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 4-89-87.mis2  
<400> 304  
ttcttcctg aacgctgggt tcatatagtt tttgtgttga gaataga 47  
<210> 305

<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-123-184, variant version of SEQ ID228  
<221> allele  
<222> 24  
<223> base C ; G in SEQ ID228  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-123-184.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-123-184.mis2  
<400> 305  
ccagcccaga acattcacca gctcggccaa gagttctgct gggtttt 47  
<210> 306  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-128-202, variant version of SEQ ID229  
<221> allele  
<222> 24  
<223> base C ; A in SEQ ID229  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-128-202.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-128-202.mis2  
<400> 306  
aatgtctggtt tcttagagaa ctgcaacaca cacacataca tacacac 47  
<210> 307  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-128-275, variant version of SEQ ID230  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID230  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-128-275.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-128-275.mis2  
<400> 307  
acaccctac ctcacatgtg taggcaaagt tatgcatata tgtctct 47  
<210> 308  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>

```

<221> allele
<222> 1..47
<223> polymorphic fragment 99-128-313, variant version of SEQ ID231
<221> allele
<222> 24
<223> base G ; A in SEQ ID231
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-128-313.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-128-313.mis2
<400> 308
tatgtctcta gacagatata catgagattc tatttggcat agaaaaa 47
<210> 309
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-128-60, variant version of SEQ ID232
<221> allele
<222> 24
<223> base T ; C in SEQ ID232
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-128-60.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-128-60.mis2
<400> 309
gcactgtgac ccaggcgcta ggttcctctt acagtgcac tccgaca 47
<210> 310
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-12907-295, variant version of SEQ ID233
<221> allele
<222> 24
<223> base G ; A in SEQ ID233
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-12907-295.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-12907-295.mis2
<400> 310
gctatatggc attatatctc cacggggcag acctgatgta caagatg 47
<210> 311
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-130-58, variant version of SEQ ID234
<221> allele

```

```

<222> 24
<223> base T ; C in SEQ ID234
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-130-58.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-130-58.mis2
<400> 311
aaagcaaaag agcttcaaaa atatttcagg agtgtgcata tggcgag 47
<210> 312
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-134-362, variant version of SEQ ID235
<221> allele
<222> 24
<223> base T ; G in SEQ ID235
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-134-362.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-134-362.mis2
<400> 312
caaaacactc atgttagtta gattattatt cctattacaa agataag 47
<210> 313
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-140-130, variant version of SEQ ID236
<221> allele
<222> 24
<223> base T ; C in SEQ ID236
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-140-130.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-140-130.mis2
<400> 313
tgttcaaaag cagctacaga ccatatgtaa acaattgagc atggctg 47
<210> 314
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-1462-238, variant version of SEQ ID237
<221> allele
<222> 24
<223> base C ; G in SEQ ID237
<221> primer_bind
<222> 1..23

```

```

<223> potential microsequencing oligo 99-1462-238.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-1462-238.mis2
<400> 314
ccctttcāag gttagtaact catctgctgt gtttctgctt cagaagg 47
<210> 315
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-147-181, variant version of SEQ ID238
<221> allele
<222> 24
<223> base G ; A in SEQ ID238
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-147-181.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-147-181.mis2
<400> 315
gtgtcatgaa aaagagcatg atagaaagaa aaacttaaāt ctttata 47
<210> 316
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-1474-156, variant version of SEQ ID239
<221> allele
<222> 24
<223> base T ; G in SEQ ID239
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-1474-156.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-1474-156.mis2
<400> 316
cttgactca taagttaaāt atttataaca agaagaaata tggactt 47
<210> 317
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-1474-359, variant version of SEQ ID240
<221> allele
<222> 24
<223> base G ; A in SEQ ID240
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-1474-359.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-1474-359.mis2

```

<400> 317  
aaaaaaaatc aaattattgt accgaattcc ctaatatcag atgtgta 47  
<210> 318  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1479-158, variant version of SEQ ID241  
<221> allele  
<222> 24  
<223> base T ; C in SEQ ID241  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-1479-158.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-1479-158.mis2  
<400> 318  
tttaaaaatc cacttgtaat cgctgctaatt tggagtgtat attcagg 47  
<210> 319  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1479-379, variant version of SEQ ID242  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID242  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-1479-379.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-1479-379.mis2  
<400> 319  
gtagagctgt gtactgaggt caggggaagca gctcatggta cagcctt 47  
<210> 320  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-148-129, variant version of SEQ ID243  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID243  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-148-129.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-148-129.mis2  
<400> 320  
ttcatatccta tacaaataat tttgaattta atacataggg ctgcaaa 47  
<210> 321  
<211> 47

<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-148-132, variant version of SEQ ID244  
<221> allele  
<222> 24  
<223> base T ; C in SEQ ID244  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-148-132.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-148-132.mis2  
<400> 321  
atatctatac aaataatttt gaatttaata catagggctg caaaaca  
<210> 322  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-148-139, variant version of SEQ ID245  
<221> allele  
<222> 24  
<223> base T ; C in SEQ ID245  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-148-139.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-148-139.mis2  
<400> 322  
tacaaataat tttgaattta atatataggc ctgcaaaaca aggttga  
<210> 323  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-148-140, variant version of SEQ ID246  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID246  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-148-140.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-148-140.mis2  
<400> 323  
acaaataatt ttgaatttaa tacgtagggc tgcaaaacaa gggtgat  
<210> 324  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele

47

47

47



<222> 1..47  
<223> polymorphic fragment 99-148-182, variant version of SEQ ID247  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID247  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-148-182.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-148-182.mis2  
<400> 324

ttgatgttga tatgggcaac tgtgtgttgg atgggtcccaa agcattc

47

<210> 325  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>

<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-148-366, variant version of SEQ ID248  
<221> allele  
<222> 24  
<223> base T ; G in SEQ ID248  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-148-366.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-148-366.mis2  
<400> 325

tccttgtaa aggtctctcc ctgttgctca cggctgccgc ctcaaag

47

<210> 326  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>

<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-148-76, variant version of SEQ ID249  
<221> allele  
<222> 24  
<223> base T ; C in SEQ ID249  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-148-76.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-148-76.mis2  
<400> 326

tgatagaatg ccttctgaa ttattactct tgatggcttc ataaaac

47

<210> 327  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>

<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1480-290, variant version of SEQ ID250  
<221> allele  
<222> 24

<223> base T ; G in SEQ ID250  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-1480-290.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-1480-290.mis2  
<400> 327  
tgcaccatct tcaccacaac ccctggcaac cactgatcct tttactg 47  
<210> 328  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1481-285, variant version of SEQ ID251  
<221> allele  
<222> 24  
<223> base T ; G in SEQ ID251  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-1481-285.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-1481-285.mis2  
<400> 328  
tcccataacc tgttttgctt ctctctctaa cctcaagatg gtataaa 47  
<210> 329  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1484-101, variant version of SEQ ID252  
<221> allele  
<222> 24  
<223> base C ; A in SEQ ID252  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-1484-101.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-1484-101.mis2  
<400> 329  
aaaaagatca aatataagca tgtcactcct ctccttaaaa tctcagt 47  
<210> 330  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-1484-328, variant version of SEQ ID253  
<221> allele  
<222> 24  
<223> base C ; G in SEQ ID253  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-1484-328.mis1

```

<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-1484-328.mis2
<400> 330
ggacacgtgg tcatgaggag tttcaaggga ttcagttttc agatccc
<210> 331
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-1485-251, variant version of SEQ ID254
<221> allele
<222> 24
<223> base T ; G in SEQ ID254
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-1485-251.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-1485-251.mis2
<400> 331
gattgccttg atatatgctc ccatagaacc aagaatgtcc ccttttc
<210> 332
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-1490-381, variant version of SEQ ID255
<221> allele
<222> 24
<223> base T ; C in SEQ ID255
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-1490-381.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-1490-381.mis2
<400> 332
tgacacgtgg aaataccatg tcatggtacg ctactgtgca tctcttc
<210> 333
<211> 47
<212> DNA
<213> Homo Sapiens
<220>
<221> allele
<222> 1..47
<223> polymorphic fragment 99-1493-280, variant version of SEQ ID256
<221> allele
<222> 24
<223> base G ; A in SEQ ID256
<221> primer_bind
<222> 1..23
<223> potential microsequencing oligo 99-1493-280.mis1
<221> primer_bind
<222> 25..47
<223> complement potential microsequencing oligo 99-1493-280.mis2
<400> 333

```

ggatgacaga gtattgttgg agggatgggg tttggctgct tgttttt 47  
<210> 334  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-151-94, variant version of SEQ ID257  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID257  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-151-94.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-151-94.mis2  
<400> 334  
attgagatca ttgataagga aatgttctaa aatttcaaaa tctatat 47  
<210> 335  
<211> 47  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..47  
<223> polymorphic fragment 99-211-291, variant version of SEQ ID258  
<221> allele  
<222> 24  
<223> base G ; A in SEQ ID258  
<221> primer\_bind  
<222> 1..23  
<223> potential microsequencing oligo 99-211-291.mis1  
<221> primer\_bind  
<222> 25..47  
<223> complement potential microsequencing oligo 99-211-291.mis2  
<400> 335  
ctgggtatat cagactgacc ttcgtgtttt caacaggtca atgcctt 47  
<210> 336  
<211> 46  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> allele  
<222> 1..46  
<223> polymorphic fragment 99-213-37, variant version of SEQ ID259  
<221> allele  
<222> 23  
<223> base GC ; T in SEQ ID259  
<221> primer\_bind  
<222> 1..22  
<223> potential microsequencing oligo 99-213-37.mis1  
<221> primer\_bind  
<222> 24..46  
<223> complement potential microsequencing oligo 99-213-37.mis2  
<400> 336  
gtgcttcgg ctgcaggact gtgcggagga ctccagtgtc tgacag 46  
<210> 337  
<211> 47  
<212> DNA

<213> Homo Sapiens  
 <220>  
 <221> allele  
 <222> 1..47  
 <223> polymorphic fragment 99-221-442, variant version of SEQ ID260  
 <221> allele  
 <222> 24  
 <223> base C ; A in SEQ ID260  
 <221> primer\_bind  
 <222> 1..23  
 <223> potential microsequencing oligo 99-221-442.mis1  
 <221> primer\_bind  
 <222> 25..47  
 <223> complement potential microsequencing oligo 99-221-442.mis2  
 <400> 337  
 tgcctttgta gatatgcatg ggacttccat gacctagcca gacgaat 47  
 <210> 338  
 <211> 47  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> allele  
 <222> 1..47  
 <223> polymorphic fragment 99-222-109, variant version of SEQ ID261  
 <221> allele  
 <222> 24  
 <223> base T ; C in SEQ ID261  
 <221> primer\_bind  
 <222> 1..23  
 <223> potential microsequencing oligo 99-222-109.mis1  
 <221> primer\_bind  
 <222> 25..47  
 <223> complement potential microsequencing oligo 99-222-109.mis2  
 <400> 338  
 caggtgagga gtgctggatt ggctacgata tgaatttctt cagcagt 47  
 <210> 339  
 <211> 18  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> primer\_bind  
 <222> 1..18  
 <223> upstream amplification primer for SEQ 185, SEQ 262, SEQ 186, SEQ 263,  
 SEQ 187, SEQ 264  
 <400> 339  
 tctaactct catccaac 18  
 <210> 340  
 <211> 19  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> primer\_bind  
 <222> 1..19  
 <223> upstream amplification primer for SEQ 188, SEQ 265, SEQ 189, SEQ 266  
 <400> 340  
 gttatcgtga gactttttc 19  
 <210> 341  
 <211> 18  
 <212> DNA  
 <213> Homo Sapiens  
 <220>

<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 190, SEQ 267, SEQ 191, SEQ 268  
<400> 341  
tgctggtgct gtgataac 18  
<210> 342  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 192, SEQ 269, SEQ 193, SEQ 270  
<400> 342  
tacagccctg taagacac 18  
<210> 343  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> upstream amplification primer for SEQ 194, SEQ 271  
<400> 343  
cagtatgttc aatgcacag 19  
<210> 344  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 195, SEQ 272, SEQ 196, SEQ 273  
<400> 344  
aaaacatcga catgggac 18  
<210> 345  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 197, SEQ 274, SEQ 198, SEQ 275,  
SEQ 199, SEQ 276  
<400> 345  
agcatttcga gtcattgtg 18  
<210> 346  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 200, SEQ 277, SEQ 201, SEQ 278  
<400> 346  
ccctctttcc tcatgtag 18  
<210> 347  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>

<221> primer\_bind  
<222> 1..19  
<223> upstream amplification primer for SEQ 202, SEQ 279, SEQ 203, SEQ 280  
<400> 347  
taactcgtaa acagagaac 19  
<210> 348  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 204, SEQ 281, SEQ 205, SEQ 282,  
SEQ 206, SEQ 283, SEQ 207, SEQ 284, SEQ 208, SEQ 285  
<400> 348  
gcgtattgaa gctctttg 18  
<210> 349  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 209, SEQ 286, SEQ 210, SEQ 287  
<400> 349  
aacacgggga ttttaggc 18  
<210> 350  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> upstream amplification primer for SEQ 211, SEQ 288  
<400> 350  
cacatactaa ggctaattg 19  
<210> 351  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 212, SEQ 289, SEQ 213, SEQ 290  
<400> 351  
gttgctggaa cctatttg 18  
<210> 352  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 214, SEQ 291, SEQ 215, SEQ 292  
<400> 352  
tcgatggcctt aatctacc 18  
<210> 353  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>

```
<221> primer_bind
<222> 1..18
<223> upstream amplification primer for SEQ 216, SEQ 293, SEQ 217, SEQ 294
<400> 353
aaagaggagt aaatgggg                                     18
<210> 354
<211> 18
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..18
<223> upstream amplification primer for SEQ 218, SEQ 295, SEQ 219, SEQ 296
<400> 354
tccccacagc taagagcc                                     18
<210> 355
<211> 18
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..18
<223> upstream amplification primer for SEQ 220, SEQ 297, SEQ 221, SEQ 298
<400> 355
atacctaatt tcaggggg                                     18
<210> 356
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> upstream amplification primer for SEQ 222, SEQ 299, SEQ 223, SEQ 300
<400> 356
ttaacagagt accttgag                                     19
<210> 357
<211> 18
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..18
<223> upstream amplification primer for SEQ 224, SEQ 301, SEQ 225, SEQ 302
<400> 357
gtacagcctt ttgcttac                                     18
<210> 358
<211> 18
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..18
<223> upstream amplification primer for SEQ 226, SEQ 303
<400> 358
aacgtgtcat agaaagcc                                     18
<210> 359
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
```



<222> 1..19  
<223> upstream amplification primer for SEQ 227, SEQ 304  
<400> 359  
gctgatgagt tagataacc 19  
<210> 360  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 228, SEQ 305  
<400> 360  
aaagccagga ctagaagg 18  
<210> 361  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 229, SEQ 306, SEQ 230, SEQ 307,  
SEQ 231, SEQ 308, SEQ 232, SEQ 309  
<400> 361  
gaccagggtt taagttag 18  
<210> 362  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 233, SEQ 310  
<400> 362  
tctgttagga cctgtgag 18  
<210> 363  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> upstream amplification primer for SEQ 234, SEQ 311  
<400> 363  
ccataacagc tagtacaac 19  
<210> 364  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 235, SEQ 312  
<400> 364  
tggaaggtta ctcagaag 18  
<210> 365  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind

<222> 1..19  
<223> upstream amplification primer for SEQ 236, SEQ 313  
<400> 365  
agagcatagt ataaagcag 19  
<210> 366  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> upstream amplification primer for SEQ 237, SEQ 314  
<400> 366  
ctagaagtag cttaaacag 19  
<210> 367  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> upstream amplification primer for SEQ 238, SEQ 315  
<400> 367  
gcagccaatc ttatatattc 19  
<210> 368  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> upstream amplification primer for SEQ 239, SEQ 316, SEQ 240, SEQ 317  
<400> 368  
aaggttgtag agtagaaag 19  
<210> 369  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 241, SEQ 318, SEQ 242, SEQ 319  
<400> 369  
caactgacac tataaccc 18  
<210> 370  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 243, SEQ 320, SEQ 244, SEQ 321,  
SEQ 245, SEQ 322, SEQ 246, SEQ 323, SEQ 247, SEQ 324, SEQ 248, SEQ 325, SEQ  
249, SEQ 326  
<400> 370  
cagtgaggagtg tttatgtg 18  
<210> 371  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>

<221> primer\_bind  
<222> 1..19  
<223> upstream amplification primer for SEQ 250, SEQ 327  
<400> 371  
ttgcacaaaa ggtatagag 19  
<210> 372  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> upstream amplification primer for SEQ 251, SEQ 328  
<400> 372  
aggctcccct ttgagttg 19  
<210> 373  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 252, SEQ 329, SEQ 253, SEQ 330  
<400> 373  
atcctttcta gctgggag 18  
<210> 374  
<211> 20  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..20  
<223> upstream amplification primer for SEQ 254, SEQ 331  
<400> 374  
gtttaagaat gtgtgatggg 20  
<210> 375  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> upstream amplification primer for SEQ 255, SEQ 332  
<400> 375  
aaggcaacag cgttgtgac 19  
<210> 376  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> upstream amplification primer for SEQ 256, SEQ 333  
<400> 376  
ttttgggggt tttcagtg 18  
<210> 377  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind

```

<222> 1..18
<223> upstream amplification primer for SEQ 257, SEQ 334
<400> 377
aacacaacag caaatccc
<210> 378
<211> 18
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..18
<223> upstream amplification primer for SEQ 258, SEQ 335
<400> 378
tccttacttg taaccccc
<210> 379
<211> 20
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..20
<223> upstream amplification primer for SEQ 259, SEQ 336
<400> 379
atactggcag cgtgtgcttc
<210> 380
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> upstream amplification primer for SEQ 260, SEQ 337
<400> 380
ccctttttct tcactgttc
<210> 381
<211> 20
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..20
<223> upstream amplification primer for SEQ 261, SEQ 338
<400> 381
aggggagatg agggaagttg
<210> 382
<211> 20
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..20
<223> downstream amplification primer for SEQ 185, SEQ 262, SEQ 186, SEQ 263,
SEQ 187, SEQ 264
<400> 382
gactgtatcc ttgatgcac
<210> 383
<211> 20
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind

```

<222> 1..20  
<223> downstream amplification primer for SEQ 188, SEQ 265, SEQ 189, SEQ 266  
<400> 383  
gcataattgt gcttgactgg  
<210> 384 20  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> downstream amplification primer for SEQ 190, SEQ 267, SEQ 191, SEQ 268  
<400> 384  
tgctgagagg agcttttg  
<210> 385 18  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> downstream amplification primer for SEQ 192, SEQ 269, SEQ 193, SEQ 270  
<400> 385  
tgaggactgc taggaaag  
<210> 386 18  
<211> 20  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..20  
<223> downstream amplification primer for SEQ 194, SEQ 271  
<400> 386  
acaaaatcag gaacaatggg  
<210> 387 20  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> downstream amplification primer for SEQ 195, SEQ 272, SEQ 196, SEQ 273  
<400> 387  
ttgcattttc cccccaac  
<210> 388 18  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> downstream amplification primer for SEQ 197, SEQ 274, SEQ 198, SEQ 275,  
SEQ 199, SEQ 276  
<400> 388  
accatttgga caatgggg  
<210> 389 18  
<211> 20  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind

<222> 1..20  
<223> downstream amplification primer for SEQ 200, SEQ 277, SEQ 201, SEQ 278  
<400> 389  
gctctttaaac tggctctgtg  
<210> 390 20  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> downstream amplification primer for SEQ 202, SEQ 279, SEQ 203, SEQ 280  
<400> 390  
ggcatgactt cacgtttc  
<210> 391 18  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> downstream amplification primer for SEQ 204, SEQ 281, SEQ 205, SEQ 282,  
SEQ 206, SEQ 283, SEQ 207, SEQ 284, SEQ 208, SEQ 285  
<400> 391  
aggatcttct acagtcac  
<210> 392 18  
<211> 20  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..20  
<223> downstream amplification primer for SEQ 209, SEQ 286, SEQ 210, SEQ 287  
<400> 392  
tggtagcggtt tgaaatcatc  
<210> 393 20  
<211> 20  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..20  
<223> downstream amplification primer for SEQ 211, SEQ 288  
<400> 393  
tataagcaca aataggttcc  
<210> 394 20  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> downstream amplification primer for SEQ 212, SEQ 289, SEQ 213, SEQ 290  
<400> 394  
gaataactga ggggagtg  
<210> 395 18  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind

<222> 1..19  
<223> downstream amplification primer for SEQ 214, SEQ 291, SEQ 215, SEQ 292  
<400> 395  
gtgaatctcc ttccaag  
<210> 396  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> downstream amplification primer for SEQ 216, SEQ 293, SEQ 217, SEQ 294  
<400> 396  
ctaaggtgtt gtagacag  
<210> 397  
<211> 20  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..20  
<223> downstream amplification primer for SEQ 218, SEQ 295, SEQ 219, SEQ 296  
<400> 397  
cacctcgata aatcaagtc  
<210> 398  
<211> 20  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..20  
<223> downstream amplification primer for SEQ 220, SEQ 297, SEQ 221, SEQ 298  
<400> 398  
gttcacttaa ttctgttgag  
<210> 399  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> downstream amplification primer for SEQ 222, SEQ 299, SEQ 223, SEQ 300  
<400> 399  
cgccttttct gaaagggtg  
<210> 400  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> downstream amplification primer for SEQ 224, SEQ 301, SEQ 225, SEQ 302  
<400> 400  
atcttctgca cagcagcg  
<210> 401  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19

```

<223> downstream amplification primer for SEQ 226, SEQ 303
<400> 401
tattttcttag ctcttctgg
<210> 402
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> downstream amplification primer for SEQ 227, SEQ 304
<400> 402
agcaagagtg attgtaaag
<210> 403
<211> 18
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..18
<223> downstream amplification primer for SEQ 228, SEQ 305
<400> 403
tattcagaaa ggagtgagg
<210> 404
<211> 18
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..18
<223> downstream amplification primer for SEQ 229, SEQ 306, SEQ 230, SEQ 307,
SEQ 231, SEQ 308, SEQ 232, SEQ 309
<400> 404
agagcggttct tgcctttc
<210> 405
<211> 20
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..20
<223> downstream amplification primer for SEQ 233, SEQ 310
<400> 405
ggtaacccta aaatgttatc
<210> 406
<211> 21
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..21
<223> downstream amplification primer for SEQ 234, SEQ 311
<400> 406
agaaaccata agggatatatt g
<210> 407
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19

```



<223> downstream amplification primer for SEQ 235, SEQ 312  
<400> 407  
acagtgc aaa gg ttat atc  
<210> 408 19  
<211> 21  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..21  
<223> downstream amplification primer for SEQ 236, SEQ 313  
<400> 408  
gaacaac ctt ga attag ctt g  
<210> 409 21  
<211> 21  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..21  
<223> downstream amplification primer for SEQ 237, SEQ 314  
<400> 409  
gattccagaa gtccatttca g  
<210> 410 21  
<211> 21  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..21  
<223> downstream amplification primer for SEQ 238, SEQ 315  
<400> 410  
aggtaagaat gagcaaaaag g  
<210> 411 21  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> downstream amplification primer for SEQ 239, SEQ 316, SEQ 240, SEQ 317  
<400> 411  
gcttggtgttt gttcaattc  
<210> 412 19  
<211> 18  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..18  
<223> downstream amplification primer for SEQ 241, SEQ 318, SEQ 242, SEQ 319  
<400> 412  
cttgaaatac tcccagcc  
<210> 413 18  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19

<223> downstream amplification primer for SEQ 243, SEQ 320, SEQ 244, SEQ 321,  
 SEQ 245, SEQ 322, SEQ 246, SEQ 323, SEQ 247, SEQ 324, SEQ 248, SEQ 325, SEQ  
 249, SEQ 326  
 <400> 413  
 ccatgaactg agaactttg  
 <210> 414 19  
 <211> 18  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> primer\_bind  
 <222> 1..18  
 <223> downstream amplification primer for SEQ 250, SEQ 327  
 <400> 414  
 ggtgacaggt aaagaaac  
 <210> 415 18  
 <211> 21  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> primer\_bind  
 <222> 1..21  
 <223> downstream amplification primer for SEQ 251, SEQ 328  
 <400> 415  
 attcaggcac agaagtcata c  
 <210> 416 21  
 <211> 21  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> primer\_bind  
 <222> 1..21  
 <223> downstream amplification primer for SEQ 252, SEQ 329, SEQ 253, SEQ 330  
 <400> 416  
 agggcagcac aatgtagtaa g  
 <210> 417 21  
 <211> 18  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> primer\_bind  
 <222> 1..18  
 <223> downstream amplification primer for SEQ 254, SEQ 331  
 <400> 417  
 cctctttatc tccaaacc  
 <210> 418 18  
 <211> 19  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> primer\_bind  
 <222> 1..19  
 <223> downstream amplification primer for SEQ 255, SEQ 332  
 <400> 418  
 gaaaacaatc aagctctgg  
 <210> 419 19  
 <211> 19  
 <212> DNA  
 <213> Homo Sapiens  
 <220>  
 <221> primer\_bind

<222> 1..19  
<223> downstream amplification primer for SEQ 256, SEQ 333  
<400> 419  
ccttttatatc cttggagtc 19  
<210> 420  
<211> 21  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..21  
<223> downstream amplification primer for SEQ 257, SEQ 334  
<400> 420  
tattacacgt tccaactctt c 21  
<210> 421  
<211> 20  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..20  
<223> downstream amplification primer for SEQ 258, SEQ 335  
<400> 421  
ctgtgtttaa gtgactgctg 20  
<210> 422  
<211> 21  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..21  
<223> downstream amplification primer for SEQ 259, SEQ 336  
<400> 422  
ttattgcccc acatgcttga g 21  
<210> 423  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> downstream amplification primer for SEQ 260, SEQ 337  
<400> 423  
tcattcgtct ggctaggtc 19  
<210> 424  
<211> 21  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..21  
<223> downstream amplification primer for SEQ 261, SEQ 338  
<400> 424  
gaaacagact gaagcaagga c 21  
<210> 425  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19

<223> potential microsequencing oligo for 4-14-107.mis1  
<400> 425  
acaaccacca aatgcatac 19  
<210> 426  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-14-317.mis1  
<400> 426  
acatgcaagg tgggcaaga 19  
<210> 427  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-14-35.mis1  
<400> 427  
aacacagaaa ccgctaaaa 19  
<210> 428  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-20-149.mis1  
<400> 428  
tttttgcgtgt gtcttcaaag tga 23  
<210> 429  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-20-77.mis1  
<400> 429  
acatgaagat tctgaaggg 19  
<210> 430  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-22-174.mis1  
<400> 430  
ggattgtgca gaagttgcct ttc 23  
<210> 431  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-22-176.mis1

<400> 431  
tgcagaagtt gcctttcat 19  
<210> 432  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-26-60.mis1  
<400> 432  
ggaaagtgc tcttaagac 19  
<210> 433  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-26-72.mis1  
<400> 433  
ttaagacagt tagcaggcc 19  
<210> 434  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-3-130.mis1  
<400> 434  
gggcctaaaa cagtattct 19  
<210> 435  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-38-63.mis1  
<400> 435  
agttataaga aaatcaggc 19  
<210> 436  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-38-83.mis1  
<400> 436  
gaggctaaac tttttttt 19  
<210> 437  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-4-152.mis1  
<400> 437

ttccattgt tcttgactt 19  
<210> 438  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-4-187.mis1  
<400> 438  
tataaacaga aacatggatg agt 23  
<210> 439  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-4-288.mis1  
<400> 439  
catcaactaa ttttcacaa 19  
<210> 440  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-42-304.mis1  
<400> 440  
tttaaaacta tttatgtaa 19  
<210> 441  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-42-401.mis1  
<400> 441  
taagaaagaa ttctgtgttc tgg 23  
<210> 442  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-43-328.mis1  
<400> 442  
ttctgtgttc tggccaaag 19  
<210> 443  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-43-70.mis1  
<400> 443  
atgcctcca ttattctcaa aaa 23

<210> 444  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-50-209.mis1  
<400> 444  
atatagagtg tgcattccctg aca 23  
<210> 445  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-50-293.mis1  
<400> 445  
cctgagtccc agggggctga cag 23  
<210> 446  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-50-323.mis1  
<400> 446  
tttaaaacat tgatgaatct tta 23  
<210> 447  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-50-329.mis1  
<400> 447  
acattgatga atctttatta cta 23  
<210> 448  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-50-330.mis1  
<400> 448  
gatgaatctt tattactac 19  
<210> 449  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-52-163.mis1  
<400> 449  
gaacaggata ttcttaacta cca 23  
<210> 450

<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-52-88.mis1  
<400> 450  
tccatgtcat tattattcaa aag 23  
<210> 451  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-53-258.mis1  
<400> 451  
aatcatgcag agagaatgc 19  
<210> 452  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-54-283.mis1  
<400> 452  
aagtagtttt tcacactttc tct 23  
<210> 453  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-54-388.mis1  
<400> 453  
ctatcgata catctttac 19  
<210> 454  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-55-70.mis1  
<400> 454  
aagaacctag gttttaaaa 19  
<210> 455  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-55-95.mis1  
<400> 455  
ctctctatcg tatacatctt tac 23  
<210> 456  
<211> 23



```

<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..23
<223> microsequencing oligo for 4-56-159.mis1
<400> 456
aagttttcct tctcttctgt aga                23
<210> 457
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> potential microsequencing oligo for 4-56-213.mis1
<400> 457
ctcatgttca ctctggttc                      19
<210> 458
<211> 23
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..23
<223> microsequencing oligo for 4-58-289.mis1
<400> 458
catacctgca gcctgctttt ggt                23
<210> 459
<211> 23
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..23
<223> microsequencing oligo for 4-58-318.mis1
<400> 459
tgactacttt acctgcaata ttt                23
<210> 460
<211> 23
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..23
<223> microsequencing oligo for 4-60-266.mis1
<400> 460
aacaggacca agacactgca tta                23
<210> 461
<211> 23
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..23
<223> microsequencing oligo for 4-60-293.mis1
<400> 461
aagtttcagt atttcttagc aga                23
<210> 462
<211> 19
<212> DNA

```

<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-84-241.misl  
<400> 462  
aaaaaatagt gactgccac 19  
<210> 463  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-84-262.misl  
<400> 463  
tgaataattc agttcttca 19  
<210> 464  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-86-206.misl  
<400> 464  
tcaaatacagg acacaccac 19  
<210> 465  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-86-309.misl  
<400> 465  
tctaggcagg ccactttag 19  
<210> 466  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-88-349.misl  
<400> 466  
ctaaaagaca atattcagt 19  
<210> 467  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-89-87.misl  
<400> 467  
ttcttcctg aacgctggtt tca 23  
<210> 468  
<211> 19  
<212> DNA  
<213> Homo Sapiens

<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-123-184.mis1  
<400> 468  
cccagaacat tcaccagct 19  
<210> 469  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-128-202.mis1  
<400> 469  
tctgtttctt agagaactg 19  
<210> 470  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-128-275.mis1  
<400> 470  
ccctaccta catgtgtag 19  
<210> 471  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-128-313.mis1  
<400> 471  
tctctagaca gatatacat 19  
<210> 472  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 99-128-60.mis1  
<400> 472  
cactgtgacc caggcgctag cgt 23  
<210> 473  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-12907-295.mis1  
<400> 473  
tatggcatta tatctccac 19  
<210> 474  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>

<221> primer\_bind  
<222> 1..19  
<223> microsequencing oligo for 99-130-58.mis1  
<400> 474  
caaaagagct tcaaaaata 19  
<210> 475  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-134-362.mis1  
<400> 475  
acactcatgt tagttagat 19  
<210> 476  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> microsequencing oligo for 99-140-130.mis1  
<400> 476  
caaaagcagc tacagacca 19  
<210> 477  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> microsequencing oligo for 99-1462-238.mis1  
<400> 477  
ttcaagggtta gtaactcat 19  
<210> 478  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-147-181.mis1  
<400> 478  
catgaaaaag agcatgata 19  
<210> 479  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-1474-156.mis1  
<400> 479  
tactcataag ttaaattatt 19  
<210> 480  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind

<222> 1..19  
<223> potential microsequencing oligo for 99-1474-359.mis1  
<400> 480 19  
aaaatcaaatt tattgtacc  
<210> 481  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> microsequencing oligo for 99-1479-158.mis1  
<400> 481 19  
aaaatccact tgtaatcgc  
<210> 482  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-1479-379.mis1  
<400> 482 19  
agctgtgtac tgaggtcag  
<210> 483  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-148-129.mis1  
<400> 483 19  
tatctataca aataatttt  
<210> 484  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-148-132.mis1  
<400> 484 19  
ctatacaaatt aattttgaa  
<210> 485  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-148-139.mis1  
<400> 485 19  
aataatttttg aatttaata  
<210> 486  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19

<223> potential microsequencing oligo for 99-148-140.mis1  
<400> 486  
ataattttga atttaatac 19  
<210> 487  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-148-182.mis1  
<400> 487 19  
tggtgatatg ggcaactgt  
<210> 488  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-148-366.mis1  
<400> 488 19  
tgtcaaaggt ctctccctg  
<210> 489  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-148-76.mis1  
<400> 489 19  
agaatgcctt cctgaatta  
<210> 490  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-1480-290.mis1  
<400> 490 19  
ccatcttcac cacaacccc  
<210> 491  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-1481-285.mis1  
<400> 491 19  
ataacctgtt ttgcttctc  
<210> 492  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-1484-101.mis1

<400> 492  
agatcaaata taagcatgt 19  
<210> 493  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> microsequencing oligo for 99-1484-328.mis1  
<400> 493  
acgtggtcat gaggagttt 19  
<210> 494  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-1485-251.mis1  
<400> 494  
gccttgatat atgctcca 19  
<210> 495  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> microsequencing oligo for 99-1490-381.mis1  
<400> 495  
cagtggaaat accatgtca 19  
<210> 496  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-1493-280.mis1  
<400> 496  
gacagagtat tgttggagg 19  
<210> 497  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-151-94.mis1  
<400> 497  
agatcattga taaggaaat 19  
<210> 498  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> microsequencing oligo for 99-211-291.mis1  
<400> 498

```

ttatatcaga ctgaccttc
<210> 499
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> potential microsequencing oligo for 99-213-37.mis1
<400> 499
cttccggctg caggactgt
<210> 500
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> potential microsequencing oligo for 99-221-442.mis1
<400> 500
tttgtagata tgcattggga
<210> 501
<211> 23
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..23
<223> microsequencing oligo for 99-222-109.mis1
<400> 501
caggtgagga gtgctggatt ggc
<210> 502
<211> 23
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..23
<223> microsequencing oligo for 4-14-107.mis2
<400> 502
ctatcaggca ttgcctggt tgc
<210> 503
<211> 23
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..23
<223> microsequencing oligo for 4-14-317.mis2
<400> 503
tcatgacctg tgcccacctc ttt
<210> 504
<211> 23
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..23
<223> microsequencing oligo for 4-14-35.mis2
<400> 504
tctctgcaga cagcttctgc ctg

```



<210> 505  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-20-149.mis2  
<400> 505  
agcaggcaat aaaccaaga 19  
<210> 506  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-20-77.mis2  
<400> 506  
gtgttctcag acaacaaag 19  
<210> 507  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-22-174.mis2  
<400> 507  
aaattaacat ttttgaaca 19  
<210> 508  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-22-176.mis2  
<400> 508  
acaaattaac atttttgaa 19  
<210> 509  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-26-60.mis2  
<400> 509  
aagtcgctcc tcggcctgct aac 23  
<210> 510  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-26-72.mis2  
<400> 510  
accctttaaa gtcgctcct 19  
<210> 511

<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-3-130.mis2  
<400> 511  
agttaataacc aatttaagct tta 23  
<210> 512  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-38-63.mis2  
<400> 512  
aaaaaaaaaag ttttagcctc 19  
<210> 513  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-38-83.mis2  
<400> 513  
atattctcaa cagcattgcc aaa 23  
<210> 514  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-4-152.mis2  
<400> 514  
tgtttatata taggataac 19  
<210> 515  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-4-187.mis2  
<400> 515  
tttttttttt ttttttttt 19  
<210> 516  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-4-288.mis2  
<400> 516  
tgaaatcaaa acataggta 19  
<210> 517  
<211> 19

<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-42-304.mis2  
<400> 517  
aaaaaacccct gaaaataag 19  
<210> 518  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-42-401.mis2  
<400> 518  
tctgtgggtt taaactttg 19  
<210> 519  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-43-328.mis2  
<400> 519  
actggctctg tgggtttaa 19  
<210> 520  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-43-70.mis2  
<400> 520  
ttgtgtgtg tcccatggt 19  
<210> 521  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-50-209.mis2  
<400> 521  
cataaagcct tcagtttca 19  
<210> 522  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-50-293.mis2  
<400> 522  
tcaatgtttt aaactgtcc 19  
<210> 523  
<211> 19  
<212> DNA

<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-50-323.mis2  
<400> 523  
atcgaaccct tttgtagta 19  
<210> 524  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-50-329.mis2  
<400> 524  
gcctaaatcg aaccctttt 19  
<210> 525  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-50-330.mis2  
<400> 525  
agcctaaatc gaacccttt 19  
<210> 526  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-52-163.mis2  
<400> 526  
aatagatgtg taaaattct 19  
<210> 527  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-52-88.mis2  
<400> 527  
caccttgtgt attttttaa 19  
<210> 528  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-53-258.mis2  
<400> 528  
ttaggttaaa atttgagtga gaa 23  
<210> 529  
<211> 19  
<212> DNA  
<213> Homo Sapiens

```

<220>
<221> primer_bind
<222> 1..19
<223> potential microsequencing oligo for 4-54-283.mis2
<400> 529
taagccatcg attgtatca
<210> 530
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> potential microsequencing oligo for 4-54-388.mis2
<400> 530
gtcttggcgc tgcagcgtg
<210> 531
<211> 23
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..23
<223> microsequencing oligo for 4-55-70.mis2
<400> 531
taaagatgta tacgatagag agt
<210> 532
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> potential microsequencing oligo for 4-55-95.mis2
<400> 532
gtcttggcgc tgcagcgtg
<210> 533
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> potential microsequencing oligo for 4-56-159.mis2
<400> 533
ttgactgtaa catggagac
<210> 534
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> potential microsequencing oligo for 4-56-213.mis2
<400> 534
tatcaaactc ctctgaagg
<210> 535
<211> 19
<212> DNA
<213> Homo Sapiens
<220>

```

<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-58-289.mis2  
<400> 535  
aggtaaagta gtcacccct 19  
<210> 536  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-58-318.mis2  
<400> 536  
gaagaaataa acttgcaaa 19  
<210> 537  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-60-266.mis2  
<400> 537  
agaaatactg aaactttat 19  
<210> 538  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-60-293.mis2  
<400> 538  
aggacttcct gctggcttc 19  
<210> 539  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 4-84-241.mis2  
<400> 539  
ttctgaagaa ctgaattatt cac 23  
<210> 540  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-84-262.mis2  
<400> 540  
tgagatcatg ttgctgctt 19  
<210> 541  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind

<222> 1..23  
<223> microsequencing oligo for 4-86-206.mis2  
<400> 541  
aatgttaacg tgtagatgcc att 23  
<210> 542  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-86-309.mis2  
<400> 542  
gctctctggg tcctcactc 19  
<210> 543  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-88-349.mis2  
<400> 543  
aagaacttgg aaaatctca 19  
<210> 544  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 4-89-87.mis2  
<400> 544  
ttctcaacac aaaaactat 19  
<210> 545  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-123-184.mis2  
<400> 545  
cccagcagaa ctcttggcc 19  
<210> 546  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-128-202.mis2  
<400> 546  
gtatgtatgt gtgtgtgtt 19  
<210> 547  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19

<223> potential microsequencing oligo for 99-128-275.mis2  
<400> 547  
acatatatgc atacatttg 19  
<210> 548  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-128-313.mis2  
<400> 548  
tctatgccaa atagaatct 19  
<210> 549  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-128-60.mis2  
<400> 549  
ggagtgtcac tgtaagagg 19  
<210> 550  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> microsequencing oligo for 99-12907-295.mis2  
<400> 550  
ttgtacatca ggtctgccc 19  
<210> 551  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 99-130-58.mis2  
<400> 551  
ctcgccatat gcacactcct gaa 23  
<210> 552  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> microsequencing oligo for 99-134-362.mis2  
<400> 552  
tctttgtaat aggaataat 19  
<210> 553  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> microsequencing oligo for 99-140-130.mis2



<400> 553  
catgctcaat tgtttacat 19  
<210> 554  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-1462-238.mis2  
<400> 554  
ctgaagcaga aacacagca 19  
<210> 555  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 99-147-181.mis2  
<400> 555  
tataaagatt taagtttttc ttt 23  
<210> 556  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> microsequencing oligo for 99-1474-156.mis2  
<400> 556  
ccatatttct tcttggtat 19  
<210> 557  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-1474-359.mis2  
<400> 557  
catctgatat tagggaatt 19  
<210> 558  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-1479-158.mis2  
<400> 558  
aatatacact ccaattagc 19  
<210> 559  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-1479-379.mis2  
<400> 559

```

ctgtaccatg agctgcttc 19
<210> 560
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> potential microsequencing oligo for 99-148-129.mis2
<400> 560
cagccctatg tattaaatt 19
<210> 561
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> potential microsequencing oligo for 99-148-132.mis2
<400> 561
ttgcagccct atgtattaa 19
<210> 562
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> potential microsequencing oligo for 99-148-139.mis2
<400> 562
ccttggtttg cagccctat 19
<210> 563
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> potential microsequencing oligo for 99-148-140.mis2
<400> 563
accttggttt gcagcccta 19
<210> 564
<211> 23
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..23
<223> microsequencing oligo for 99-148-182.mis2
<400> 564
gaatgctttg ggaccatcca aca 23
<210> 565
<211> 19
<212> DNA
<213> Homo Sapiens
<220>
<221> primer_bind
<222> 1..19
<223> potential microsequencing oligo for 99-148-366.mis2
<400> 565
gaggcggcag ccgtgagca 19

```

<210> 566  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-148-76.mis2  
<400> 566  
tatgaagcca tcaagagta 19  
<210> 567  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> microsequencing oligo for 99-1480-290.mis2  
<400> 567  
aaaaggatca gtggttgcc 19  
<210> 568  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> microsequencing oligo for 99-1481-285.mis2  
<400> 568  
taccatcttg aggttagag 19  
<210> 569  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-1484-101.mis2  
<400> 569  
agattttaag gagaggagt 19  
<210> 570  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-1484-328.mis2  
<400> 570  
tctgaaaact gaatccctt 19  
<210> 571  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> microsequencing oligo for 99-1485-251.mis2  
<400> 571  
aggggacatt cttggttct 19  
<210> 572

<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-1490-381.mis2  
<400> 572  
agatgcacag tagcgtagc 19  
<210> 573  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> microsequencing oligo for 99-1493-280.mis2  
<400> 573  
acaagcagcc aaaccccat 19  
<210> 574  
<211> 23  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..23  
<223> microsequencing oligo for 99-151-94.mis2  
<400> 574  
atatagattt tgaaatttta gaa 23  
<210> 575  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-211-291.mis2  
<400> 575  
cattgacctg ttgaaaaca 19  
<210> 576  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-213-37.mis2  
<400> 576  
tcagacactg gagtccctcc 19  
<210> 577  
<211> 19  
<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-221-442.mis2  
<400> 577  
gtctggctag gtcattggaa 19  
<210> 578  
<211> 19

<212> DNA  
<213> Homo Sapiens  
<220>  
<221> primer\_bind  
<222> 1..19  
<223> potential microsequencing oligo for 99-222-109.mis2  
<400> 578  
ctgaagaaat tcatatcgt